

Peculiarity of the mammal associations from the Upper Pleistocene (Dobrogea, Romania)

Alexandru Petculescu*, Emanoil Știucă

Institute of Speleology, Bucharest, Romania

Available online 4 September 2007

Abstract

Paleontological studies of the mammals from the Upper Pleistocene at Dobrogea are distinct in comparison with other mammal associations from Romania. The paleoenvironment from the last glacial cycle (Wurm) is reconstructed on the basis of the mammal cave fossils discovered in caves at Dobrogea. Research in karst deposits from the last glacial cycle is based on systematic surveying of the most important paleontological sites.

© 2007 Elsevier Ltd and INQUA. All rights reserved.

1. Introduction

Located between the Danube River and the Black Sea, the territory of Dobrogea is distinguished by many climatic and faunal peculiarities compared with other regions of southern Romania. Paleontological investigations, carried out since 1956 in the karst deposits located in the Jurassic limestones of central Dobrogea, revealed a rich mammal association.

The paleontological studies of the mammals from the Upper Pleistocene of the Dobrogea region reveal distinct particularity in comparison with other mammal associations from Romania (Burghele et al., 1995). The research in karst deposits from the last glacial cycle is based on systematic surveying of the most important paleontological sites, including La Adam Cave, Cheia Cave, Casian Cave and Gaura Vulpii Fissure (Fig. 1).

2. Mammals

The mammal remains, especially herbivores, were accumulated in caves by Upper and Middle Paleolithic hunters. These communities within caves are represented on different levels by pebble tools and bones.

Caves, used as shelters, were used not only by primitive humans but also by the large carnivores such as cave bear (*Ursus spelaeus*) and cave hyena (*Crocota spelaea*). In the absence of humans, the carnivores used these places for shelter, birth, and nurseries. Thus, La Adam cave, during the middle phases of the last glacial cycle, was used by the hyenas which brought numerous remains there (Samson and Rădulescu, 1959; Dumitrescu et al., 1963; Terzea, 1972). Thanks to this “artificial selection”, the number of the bones is not significant, but the faunal association, generally speaking, is relatively well represented.

The small mammal remains are due to the existence, next to the entrance of the caves, of nests of carnivorous birds. The remains were accumulated in chronological positions and represent a mirror of the mammalian associations outside, especially for the small mammals.

3. Faunas from the last glacial cycle from Dobrogea

During the Upper Pleistocene, Dobrogea was characterized by a cold climate with a strong continental influence and warm periods (interstadials). Northern species including *Mamuthus primigenius*, *Coelodonta antiquitatis*, *Rangifer tarandus*, *Alopex lagopus*, *Stenocranius gregalis* and *Microtus oeconomus* are present, with steppe elements including bovids, *Saiga tatarica*, cervids (*Equus* cf. *transilvanicus*, *Equus spelaeus* spp., *Equus* sp., *Hydruntinus hydruntinus*), *Vulpes corsac*, and small mammals (*Allactaga*, *Lagurus* and

*Corresponding author.

E-mail address: romanianspeleology@k.ro (A. Petculescu).




Fig. 1. Location map. (1) Bordeiu de Piatră rockshelter, (2) Cheia Cave, (3) Casian Cave, (4) Gaura Vulpilor Cave, (5) Gura Dobrogei Cave, (6) Gura Dobrogei-4 rockshelter, (7) “La Adam” Cave.

Eolagurus genus) (Table 1). The yellow lemming (*Eolagurus luteus*), now living in central Asia, is represented in Dobrogea as the westernmost point of the range of a large group (Rădulescu and Samson, 1995). Other Nordic species, such as *Lemmus* and *Dicrostonyx*, are confined to northern Romania (Markova, 1982; Ştiucă et al., 2002).

The relative warming intervals in Dobrogea are represented by species such as *Magaloceros giganteus* and *Cervus elaphus*. In the first part of MIS 3 (local interstadials Adam I and Adam II), *Bos primigenius* was identified. In the second part of the same stage (Vistorna I) *Martes* sp. and *Panthera pardus* were found in La Adam Cave. MIS 2 is characterized by the existence of wild boar (*Sus scrofa*) associated with *Magaloceros* and *Hydruntinus*.

4. Particularity of the small mammal associations

The massive presence of the small mammal remains on different levels indicates some morphological and dimensional characteristics. Different subspecies can be separated statistically, especially the steppe lemmings and *S. gregalis*, as characteristic of distinct periods.

Lagurus lagurus dobrogeicus and *S. gregalis* spp. from MIS 4 and the first part of MIS 3, are large. *S. gregalis* are very similar to the Siberian form.

Lagurus lagurus thracicus and *Stenocranius gregalis anglicus* from the last part of MIS 3 and MIS 2 are small. *S. gregalis anglicus* specimens are very close to the modern

species from central Asia and the Ural Mountains. Some species are characteristic only for a clearly delimited period of time (e.g. *Allocrietulus eversmanni*, *Sciropoda telum*).


5. Chronology of the fossil deposits

The position of the fossiliferous levels from the caves was correlated with the fossil soils from the entrance zone. A good example is the karstic deposits from La Adam cave, which grade into loess deposits and fossil soil near the entrance of the cave. This allows correlation with the well known loess stratigraphy of Dobrogea (Samson and Rădulescu, 1959; Conea, 1970; Samson, 1971). The thickness of the deposits from La Adam cave is around 15m. This section begins with the last part of the Riss, followed by a strong erosional period with red soil remains correlated with the Riss-Wurm interglacial, and a succession of layers with very rich small and large mammal remains, associated with Upper and Middle Paleolithic human industry (Leroi-Gourhan, 1969).

6. Conclusions

The similarity between faunal associations from Crimea and Dobrogea during the last glacial cycle occurred when the level of the Black Sea was 80–90 m below the present sea level, and Dobrogea was directly connected to the Crimea. These regions represent a bridge between the

Table 1
Biochronology and chronostratigraphy of Dobrogea



faunal associations from Eastern Europe and the Balkan Peninsula.

Evidence includes:

- dominance of *U. spelaeus* in the first stage of the last glacial cycle;
- presence of the steppe lemmings (*Lagurus* and *Eolagurus*) in both regions;
- presence of *Scirtopoda telum* in both regions;
- association between *S. tatarica* and *Hydruntinus*, common in Dobrogea and Crimea;
- presence of the carnivores *V. corsac* and *A. lagopus* in both regions;

- presence of *Ovis orientalis* remains in caves from Dobrogea and Crimea;
- Dobrogea was a very distinct region in the middle of the migration route between eastern Europe and central Asia and southeastern Europe in the last glacial cycle.

References

- Burghel, A., Rădulescu, C., Samson, P.-M., Ştiucă, E., 1995. La première faune interglaciaire (Eemien ?) de micromammifères d'Olténie. Travaux de l'Institut de Spéologie Emile Racovitza 33, 101–106.
- Conca, A., 1970. Quaternary formations from Dobrogea. Education Academy R. S. Romania, 234pp.

- Dumitrescu, M., Samson, P., Terzea, E., Rădulescu, C., Ghica, M., 1963. La Adam cave, Pleistocene site. *Lucrarile Institutului de Speologie "Emil Racoviță I–II*, 229–284.
- Leroi-Gourhan, A., 1969. Dénomination des oscillations wurmiennes. *Bulletin de l'AFEQ* 4.
- Markova, A.K., 1982. Pleistocene Rodents of the Russian Plain (in Russian). Nauka, Moscow, 184pp.
- Rădulescu, C., Samson, P.-M., 1995. On some middle late pleistocene rare small mammal elements from the Karst deposits of central Dobrogea (Romania). *Theoretical and Applied Karstology* 8, 163–173.
- Samson, P., 1971. La stratigraphie du Quaternaire supérieur en Roumanie. VIIIe Congrès INQUA, Paris 1969. *Études sur le Quaternaire dans le Monde*.
- Samson, P., Rădulescu, C., 1959. Beiträge zur Kenntnis der Chronologie de "Jüngerer Losses" in der Dobrudscha. *Eiszeitalter u. Gegenwart* 10, 129–204.
- Ştiucă, E., Petculescu, A., Arghir, R., 2002. Nouvelles données sur les faunes de microvertébrés du Pleistocène supérieur de Dobrogea, avec regard spécial sur la zone Movile (dep. de Constanta, Roumanie). *Travaux de l'Institut de Spéologie "Emile Racovitza"*, Bucaresti, 41–42, pp. 111–121.
- Terzea, E., 1972. Remarques sur la morphologie dentaire et la répartition de *Microtus nivalis* Martins (Rodentia, Mammalia) dans le Pléistocène de Roumanie. *Travaux de l'Institut de Spéologie Emile Racovitza XI*, 271–298.