

- tocopherol and plastoquinol metabolic pathways at the plastoglobule. *Biochem J* 2010, **425**:389-399.
13. Kruk J, Karpinski S: An HPLC-based method of estimation of the total redox state of plastoquinone in chloroplasts, the size of the photochemically active plastoquinone-pool and its redox state in thylakoids of *Arabidopsis*. *BBA-Bioenergetics* 2006, **1757**:1669-1675.
 14. Yoshida K, Shibata M, Terashima I, Noguchi K: Simultaneous determination of *in vivo* plastoquinone and ubiquinone redox states by HPLC-based analysis. *Plant Cell Physiol* 2010, **51**:836-841.
 15. Byrdwell WC: Atmospheric pressure chemical ionization mass spectrometry for analysis of lipids. *Lipids* 2001, **36**:327-346.
 16. Swartz ME: UPLC (TM): An introduction and review. *J Liq Chromatogr Relat Technol* 2005, **28**:1253-1263.
 17. Nguyen DTT, Guillaume D, Rudaz S, Veuthey JL: Fast analysis in liquid chromatography using small particle size and high pressure. *J Sep Sci* 2006, **29**:1836-1848.
 18. Plumb R, Castro-Perez J, Granger J, Beattie I, Joncour K, Wright A: Ultra-performance liquid chromatography coupled to quadrupole-orthogonal time-of-flight mass spectrometry. *Rapid Commun Mass Sp* 2004, **18**:2331-2337.
 19. Lauridsen C, Leonard SW, Griffin DA, Liebler DC, McClure TD, Traber MG: Quantitative analysis by liquid chromatography tandem mass spectrometry of deuterium-labeled and unlabeled vitamin E in biological samples. *Anal Biochem* 2001, **289**:89-95.
 20. Lanina SA, Toledo P, Sampels S, Kamal-Eldin A, Jastrebova JA: Comparison of reversed-phase liquid chromatography-mass spectrometry with electrospray and atmospheric pressure chemical ionization for analysis of dietary tocopherols. *J Chromatogr A* 2007, **1157**:159-170.
 21. Kurilich AC, Britz SJ, Clevidence BA, Novotny JA: Isotopic labeling and LC-APCI-MS quantification for investigating absorption of carotenoids and phyloquinone from kale (*Brassica oleracea*). *J Agric Food Chem* 2003, **51**:4877-4883.
 22. Suhara Y, Kamao M, Tsugawa N, Okano T: Method for the determination of vitamin K homologues in human plasma using high-performance liquid chromatography-tandem mass spectrometry. *Anal Chem* 2005, **77**:757-763.
 23. Maeda H, Song W, Sage TL, DellaPenna D: Tocopherols play a crucial role in low-temperature adaptation and phloem loading in *Arabidopsis*. *Plant Cell* 2006, **18**:2710-2732.
 24. Szymanska R, Kruk J: Plastoquinol is the main prenyl lipid synthesized during acclimation to high light conditions in *Arabidopsis* and is converted to plastoquinone by tocopherol cyclase. *Plant Cell Physiol* 2010, **51**:537-545.
 25. Kruk J: Charge-transfer complexes of plastoquinone and alpha-tocopherol quinone *in vitro*. *Biophys Chem* 1988, **30**:143-149.
 26. Hiltbrunner A, Bauer J, Alvarez-Huerta M, Kessler F: Protein translocon at the *Arabidopsis* outer chloroplast membrane. *Biochem Cell Biol* 2001, **79**:629-635.