



Pavlovian Conditioning of Larval *Drosophila*: An Illustrated, Multilingual, Hands-On Manual for Odor-Taste Associative Learning in Maggots

Birgit Michels¹, Timo Saumweber¹, Roland Biernacki², Jeanette Thum², Rupert D. V. Glasgow¹, Michael Schleyer¹, Yi-chun Chen¹, Claire Eschbach³, Reinhard F. Stocker⁴, Naoko Toshima¹, Teiichi Tanimura⁵, Matthieu Louis⁶, Gonzalo Arias-Gil⁷, Manuela Marescotti⁸, Fabio Benfenati⁹ and Bertram Gerber^{1, 10, 11*}

¹ Department of Genetics, Leibniz Institute for Neurobiology, Magdeburg, Germany, ² Department Neurobiology and Genetics, Julius Maximilians University, Würzburg, Germany, ³ HHMI Janelia Research Campus, Ashburn, VA, USA, ⁴ Department of Biology, Université de Fribourg, Fribourg, Switzerland, ⁵ Department of Biology, Kyushu University, Fukuoka, Japan, ⁶ Department of Molecular, Cellular, and Developmental Biology, University of California, Santa Barbara, Santa Barbara, CA, USA, ⁷ Department Systems Physiology, Leibniz Institute for Neurobiology Magdeburg, Magdeburg, Germany, ⁸ Brainwave Discovery Ltd., Edinburgh, UK, ⁹ Italian Institute of Technology, Center for Synaptic Neuroscience and Technology, Genova, Italy, ¹⁰ Institute of Biology, Otto von Guericke University, Magdeburg, Germany, ¹¹ Center for Behavioral Brain Sciences, Magdeburg, Germany

OPEN ACCESS

Edited by:

Martin Giurfa,
Research Center on Animal Cognition
(CNRS), France

Reviewed by:

Jean-François Ferveur,
Centre des Sciences du Goût et de
l'Alimentation (CSGA) (CNRS), France
Simon G. Sprecher,
University of Fribourg, Switzerland

*Correspondence:

Bertram Gerber
bertram.gerber@lin-magdeburg.de

Received: 07 December 2016

Accepted: 27 February 2017

Published: 19 April 2017

Citation:

Michels B, Saumweber T, Biernacki R, Thum J, Glasgow RDV, Schleyer M, Chen Y-c, Eschbach C, Stocker RF, Toshima N, Tanimura T, Louis M, Arias-Gil G, Marescotti M, Benfenati F and Gerber B (2017) Pavlovian Conditioning of Larval *Drosophila*: An Illustrated, Multilingual, Hands-On Manual for Odor-Taste Associative Learning in Maggots. *Front. Behav. Neurosci.* 11:45. doi: 10.3389/fnbeh.2017.00045

Larval *Drosophila* offer a study case for behavioral neurogenetics that is simple enough to be experimentally tractable, yet complex enough to be worth the effort. We provide a detailed, hands-on manual for Pavlovian odor-reward learning in these animals. Given the versatility of *Drosophila* for genetic analyses, combined with the evolutionarily shared genetic heritage with humans, the paradigm has utility not only in behavioral neurogenetics and experimental psychology, but for translational biomedicine as well. Together with the upcoming total synaptic connectome of the *Drosophila* nervous system and the possibilities of single-cell-specific transgene expression, it offers enticing opportunities for research. Indeed, the paradigm has already been adopted by a number of labs and is robust enough to be used for teaching in classroom settings. This has given rise to a demand for a detailed, hands-on manual directed at newcomers and/or at laboratory novices, and this is what we here provide.

The paradigm and the present manual have a unique set of features:

- The paradigm is cheap, easy, and robust;
- The manual is detailed enough for newcomers or laboratory novices;
- It briefly covers the essential scientific context;
- It includes sheets for scoring, data analysis, and display;
- It is multilingual: in addition to an English version we provide German, French, Japanese, Spanish and Italian language versions as well.

The present manual can thus foster science education at an earlier age and enable research by a broader community than has been the case to date.

Keywords: olfaction, taste, cognition, memory, reinforcement, association

Predictive, associative learning enables animals to decipher many aspects of the causal structure of the world and to behave accordingly (Dickinson, 2001). It is therefore a ubiquitous faculty across the animal kingdom. Indeed, following the pioneering work of Ebbinghaus, Pavlov, and Thorndike, research has uncovered remarkable conservation in the mechanisms of learning and memory (Kandel et al., 2014). Because of the feasibility of genetic screens combined with robust behavioral protocols, *Drosophila* has been one of the workhorses for these endeavors (Benzer, 1967; Dudai et al., 1976; Heisenberg et al., 1985; Tully and Quinn, 1985; reviews include Heisenberg, 2003; Gerber et al., 2014; Guven-Ozkan and Davis, 2014; Harris and Littleton, 2015; Oswald and Waddell, 2015; Gerber and Aso, in press). The field received a further boost when versatile methods for transgene expression were introduced (Rubin and Spradling, 1982; O’Kane and Gehring, 1987; Brand and Perrimon, 1993), opening up the possibility for experimental manipulation with cellular specificity at the single-neuron level (Pfeiffer et al., 2010; Jenett et al., 2012; Aso et al., 2014a,b). These and related techniques (reviews include Venken et al., 2011; Sivanantharajah and Zhang, 2015) now make it relatively straightforward to express any transgene, in any cell or group of cells, at any time. Thus, *Drosophila* has become a model system for understanding learning and memory not “only” at the molecular level, but also for understanding the function of molecules within behaviorally meaningful circuitry—as envisioned by Hotta and Benzer (1970).

With a slight delay (befitting their shuffling gait, as we hesitate to add), *Drosophila* larvae entered the stage as the subjects of behavioral neurogenetics (e.g., Aceves-Piña and Quinn, 1979; Rodrigues, 1980), receiving renewed attention since the mid-1990s (Stocker, 1994; Cobb, 1999; Sokolowski, 2001; Gerber and Stocker, 2007; Gomez-Marin and Louis, 2012; Keene and Sprecher, 2012; Diegelmann et al., 2013). Larvae possess 10 times fewer neurons than adult flies, and in many cases appear to lack cellular redundancy altogether. Even so, they feature fundamental adult-like circuit motifs (e.g., in the olfactory pathways: Vosshall and Stocker, 2007; Stocker, 2008) and exhibit fundamental faculties of behavior, including learning and memory (see below). Last but not least, a synapse-by-synapse connectome of the larval nervous system seems within reach, and driver strains for transgenic manipulation can now be established to cover the neurons of the larva, one at a time (Li et al., 2014; Ohyama et al., 2015; Berck et al., 2016; Fushiki et al., 2016; Jovanic et al., 2016; Schlegel et al., 2016; Schneider-Mizell et al., 2016; Zwart et al., 2016). Taken together, the possibilities for research into the behavioral neurogenetics of larval *Drosophila* appear enticing, given the combination of analytical power, ease, elegance, and completeness.

The current contribution deals with Pavlovian odor-reward learning in larval *Drosophila* (Scherer et al., 2003; Neuser et al., 2005; **Figure 1**). In brief, the larvae are free to move about an agarose-filled Petri dish; the agarose substrate can either be supplemented with sugar reward, or can be used as plain substrate, not containing reward. An odor A (gray cloud in **Figure 1**) is presented together with a reward-supplemented substrate (+; indicated by green color in **Figure 1**). Then the




FIGURE 1 | (A) Principle of the behavioral paradigm. In a Petri-dish assay, different groups of larvae receive odor A (gray cloud) paired with a sugar reward (green circle), alternated with presentations of another odor B (white cloud) without a reward (A+/B training); a second group of larvae is trained reciprocally (A/B+). Then, for both groups the preference of the animals

(Continued)

FIGURE 1 | Continued

between odors A and B is measured as the number of animals located on the A-side minus the number of animals located on the B-side, divided by the total number of animals (including the ones located on the middle stripe). The Performance Index is calculated as the difference in preference between the A+/B versus A/B+ trained groups of larvae (divided by 2 to yield scores between -1 and 1). The Performance Index thus represents associative memory, averaging-out effects of innate odor preference, odor exposure, reward exposure, or handling. Note that the sequence of trials is alternated across repetitions of the experiment (i.e., B/A+ and B+/A). Various sugars, aspartic acid, or low-concentration salt can alternatively be used as a taste reward; as taste punishment, quinine, or high-concentration salt can be used.

(B) Example data from a non-academic setting. For the Preference scores (left) and the associative Performance Indices (right) the box plots show the median as the middle line, the 25/75% quantiles as box boundaries, and the 10/90% quantiles as whiskers. For the Preference scores *refers to $P < 0.05$ in a Mann-Whitney U -test ($N = 16, 16$); for the associative Performance Indices based on these Preference scores *refers to $P < 0.05$ in a one-sample sign-test. If the odor pairs, or the concentrations of the odors in a pair, are chosen such that one of them is more strongly attractive than the other, the Preference scores of both reciprocally trained groups will be shifted along the y-axis, i.e., will be “asymmetrically” different from zero. This does not affect the interpretation of the Performance Index as reflecting associative memory, however, because the Performance Index is based on the difference in Preference scores between the reciprocally trained groups (for more detail see **Supplemental Material 1**).

(C) A class of 8th grade high school students performing odor-reward learning in larval *Drosophila* in a 1-day course at the Gymnasium Stettisches Institut, Augsburg, Germany. The histogram at the bottom shows the median Performance Indices from 11 such experiments in various non-academic settings, with sample sizes in the range of $N = 12$ – 20 each.

(D) Side-view of a 3D print of the larval body (top; image courtesy of R. Blumenstein, LIN) and schematic overview of the internal organs of a larva (bottom; modified from Demerec and Kaufmann, 1972).

(E) Simplified circuit diagram showing the processing of odor and taste reward. AL, antennal lobe; MBINs, mushroom body input neurons; LP, lateral protocerebrum; MB-KC, mushroom body Kenyon cells; MBONs, mushroom body output neurons; OSN, olfactory sensory neurons; PN, projection neurons. SEZ, subesophageal zone. The pink color indicates an MBIN activated by reward; the light pink color indicates an MBIN activated by punishment. The star indicates presynaptic plasticity in the MB-KC to MBON connection; the ~symbol indicates that the pathway from the MBONs toward motor control is susceptible to modulation, including modulation by the testing situation. For more details, see text. Images taken from Gerber et al. (2010) **(C)** and Demerec and Kaufmann (1972) **(D)**. The following copyright holders kindly granted permission to use these figures: Cold Spring Harbor Laboratory Press **(C)** and The Carnegie Institution **(D)**.

larvae are transferred to a second Petri dish, this time with the plain substrate, and exposed to a different odor B (white cloud in **Figure 1**). After repeating this A+/B procedure two more times, the animals are transferred to a test Petri dish and are offered a choice between the two odors. A second set of larvae is trained reciprocally (A/B+) and is likewise tested for its preference between the two odors. If the larvae systematically prefer the previously rewarded odor relative to the previously non-rewarded odor, the conclusion is that an odor-sugar associative memory has been formed. In other words, the odor-reward association established in training guides the larvae's search for reward during the test (Gerber and Hendel, 2006; Schleyer et al., 2011, 2015a,b).

The working hypothesis as to how this type of learning comes about has recently been reviewed (Diegelmann et al., 2013) and is largely concordant with what has been suggested for adult flies

(Heisenberg, 2003; Gerber et al., 2014; Guven-Ozkan and Davis, 2014; Harris and Littleton, 2015; Oswald and Waddell, 2015; Gerber and Aso, in press) and other insects such as the honey bee (Tedjakumala and Giurfa, 2013; Menzel, 2014). In brief, larval olfactory sensory neurons are located in the dorsal organ and project to the antennal lobe. Downstream of the antennal lobe, the olfactory processing stream splits: one collateral of the projection neurons targets the lateral protocerebrum, which features premotor centers for innate olfactory behavior. The other collateral takes a “detour” to the mushroom bodies. According to the ligand profiles of the olfactory sensory neurons, the cellular properties and the connectivity within this system, including local circuitry within the antennal lobe, odors can thus be coded across these ascending olfactory pathways.

Gustatory pathways originate from multiple larval cephalic sense organs, bypass the brain, and target the subesophageal zone and premotor centers (Apostolopoulou et al., 2015). Taste pathways are thus linked relatively closely to the motor system. Notably, a “detour” branch also splits off from the gustatory pathway. From the subesophageal zone this sends information about the reinforcing value of the food toward the brain. Through an as yet unknown number of synaptic steps, this activates octopaminergic as well as dopaminergic input neurons signaling toward the Kenyon cells of the mushroom body (Schroll et al., 2006; Rohwedder et al., 2016; regarding adult *Drosophila*, reviews include Heisenberg, 2003; Gerber et al., 2014; Guven-Ozkan and Davis, 2014; Oswald and Waddell, 2015; Gerber and Aso, in press; see also Hammer, 1993; Kreissl et al., 1994 on the bee).

Within the mushroom body Kenyon cells, a coincidence can thus be detected between olfactory input in terms of an odor-specific subset of activated Kenyon cells, and an internal aminergic reinforcement signal. This coincidence modulates the synapse between the odor-activated set of mushroom body Kenyon cells and their output neurons, by processes taking place presynaptically within the respective Kenyon cells. If a trained odor is subsequently encountered, it is via this odor-specific set of modulated synapses that the balance is shifted between mushroom body output neurons favoring approach and mushroom body output neurons mediating avoidance. By analogy with what has been observed in adult *Drosophila* (for reviews see Oswald and Waddell, 2015; Gerber and Aso, in press), learned approach may come about by a weakening of synapses from Kenyon cells to those output neurons that are sufficient for avoidance, resulting in net relative attraction. Note that the pathway from the mushroom body output neurons carrying learned valence signals toward motor control comprises an as yet unknown number of synaptic steps and is susceptible to modulation, including modulation by the testing situation (Gerber and Hendel, 2006; Schleyer et al., 2011, 2015a,b).

Since its introduction this paradigm has made significant advances possible, including the first application of Channelrhodopsin-2 in a brain (Schroll et al., 2006), and the discovery of memories specific to the kind of reward (fructose vs. amino acid) and the kind of punishment (quinine versus high-concentration salt; Schleyer et al., 2015a). It has been adopted by a number of labs, including new groups entering the field of learning and memory. Indeed, the paradigm is robust

enough to be routinely used for undergraduate teaching and in classroom settings. This has given rise to a demand for a detailed, hands-on manual directed at newcomers in the field of behavioral science and/or at laboratory novices, and this is what we here provide (**Supplemental Materials 1–16**). The paradigm and the presented manual have a unique set of features:

- The paradigm is cheap and easy to carry out, and can be performed in classroom settings under “degraded” experimental conditions;
- The manual is richly illustrated and detailed enough to allow newcomers or laboratory novices, even at high school level, to perform the experiment;
- It features brief “introduction” and “outlook” sections covering the scientific context and guidelines for the display and the analysis of the data;
- It includes data sheets for scoring, and customized excel sheets for data analysis and display;
- Possibly most importantly for use in schools, we provide not only an English version (**Supplemental Materials 1–3**), but German (**Supplemental Materials 4–6**), French (**Supplemental Materials 7–9**), Japanese (**Supplemental Material 10**) Spanish (**Supplemental Materials 11–13**), and Italian (**Supplemental Materials 14–16**) language versions as well.

The current contribution can thus foster science education at an earlier age and enable research by a broader community than has been the case to date (Gerber et al., 2010, 2013; Apostolopoulou et al., 2013). The paradigm allows experimental access to a fascinating aspect of nervous system function: the adaptive balance between robustness and flexibility of behavior. Given the versatility of *Drosophila* for genetic analyses, combined with their evolutionarily shared genetic heritage with humans, the paradigm has utility not only in behavioral science, genetics, neurobiology, and experimental psychology, but for translational biomedicine as well.

ETHICS STATEMENT

Procedures comply with applicable law for experimentation with invertebrates of the State of Sachsen-Anhalt and the Federal Republic of Germany.

AUTHOR CONTRIBUTIONS

BG: Authored manuscript, co-authored Supplement 1–6. BM, TS, RB, JT, RG, MS, YC: Authored Supplement 1–6, co-authored manuscript. CE, RS, ML: Authored Supplement 7–9, co-authored manuscript. NT, TT: Authored Supplement 10. GA, RG: Authored Supplement 11–13. MM, FB: Authored Supplements 14–16.

ACKNOWLEDGMENTS

The development of this paradigm was made possible by a grant from the Volkswagen Foundation (to BG, then located at the Université de Fribourg and hosted by RS). The authors

acknowledge institutional support by the Leibniz Institut für Neurobiologie (LIN) Magdeburg, Wissenschaftsgemeinschaft Gottfried Wilhelm Leibniz (WGL), Universität Magdeburg, Universität Würzburg, HHMI Janelia Research Campus, and the Université de Fribourg. Project funding from the Deutsche Forschungsgemeinschaft (DFG) (CRC 779 Motivated Behavior; GE1091/4-1), the Bundesministerium für Bildung und Forschung (BMBF; Bernstein Focus Insect Inspired Robotics), and the European Commission (FP7-ICT) [Miniature Insect Model for Active Learning (MINIMAL)] is gratefully acknowledged. We thank the teachers and students of the Gymnasium Stettensches Institut, Augsburg, and of the Domgymnasium, Magdeburg, Germany, Petra Skiebe-Corrette, NatLab, Berlin, Germany, as well as Janna Klein and Kirsten Tiedemann, Lübecker offenes Labor (LoLa), Lübeck, Germany, for inspiring hours of experimentation and user comments on earlier versions of this manual, Reinhard Blumenstein (LIN) for providing images, and Tomoko Ohyama, HHMI Janelia Research Campus, Ashburn, USA, for comments on Supplement 10.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fnbeh.2017.00045/full#supplementary-material>

Supplemental Materials 1–3 | A manual for odor-reward learning in larval *Drosophila* (Supplemental Material 1), example of a table for data analysis (Supplemental Material 2), and an empty table for entering and analyzing one’s own data (Supplemental Material 3), in the English language.

Versions of this manual in the German, French, Japanese, Spanish, and Italian languages can be found in **Supplemental Materials 4–6, 7–9, 10, 11–13, 14–16**, respectively.

Supplemental Materials 4–6 | A manual for odor-reward learning in larval *Drosophila* (Supplemental Material 4), example of a table for data analysis (Supplemental Material 5), and an empty table for entering and analyzing one’s own data (Supplemental Material 6), in the German language.

Versions of this manual in the English, French, Japanese, Spanish, and Italian languages can be found in **Supplemental Materials 1–3, 7–9, 10, 11–13, 14–16**, respectively.

Supplemental Materials 7–9 | A manual for odor-reward learning in larval *Drosophila* (Supplemental Material 7), example of a table for data analysis (Supplemental Material 8), and an empty table for entering and analyzing one’s own data (Supplemental Material 9), in the French language.

Versions of this manual in the English, German, Japanese, Spanish, and Italian languages can be found in **Supplemental Materials 1–3, 4–6, 10, 11–13, 14–16**, respectively.

Supplemental Material 10 | A manual for odor-reward learning in larval *Drosophila* in the Japanese language.

Versions of this manual in the English, German, French, Spanish, and Italian languages can be found in **Supplemental Materials 1–3, 4–6, 7–9, 11–13, 14–16**, respectively. For a table for data analysis and an empty table for entering and analyzing one’s own data, please use the document in the English language (**Supplemental Materials 2, 3**, respectively).

Supplemental Materials 11–13 | A manual for odor-reward learning in larval *Drosophila* (Supplemental Material 11), example of a table for data analysis (Supplemental Material 12), and an empty table for entering and analyzing one’s own data (Supplemental Material 13), in the Spanish language.

Versions of this manual in the English, German, French, Japanese, and

Italian languages can be found in **Supplemental Materials 1–3, 4–6, 7–9, 10, 14–16**, respectively.

Supplemental Materials 14–16 | A manual for odor-reward learning in larval *Drosophila* (Supplemental Material 14), example of a table for data

analysis (Supplemental Material 15), and an empty table for entering and analyzing one's own data (Supplemental Material 16), in the Italian language. Versions of this manual in the English, German, French, Japanese, and Spanish languages can be found in **Supplemental Materials 1–3, 4–6, 7–9, 10, and 11–13**, respectively.

REFERENCES

- Aceves-Piña, E. O., and Quinn, W. G. (1979). Learning in normal and mutant *Drosophila* larvae. *Science* 206, 93–96. doi: 10.1126/science.206.4414.93
- Apostolopoulou, A. A., Rist, A., and Thum, A. S. (2015). Taste processing in *Drosophila* larvae. *Front. Integr. Neurosci.* 9:50. doi: 10.3389/fnint.2015.00050
- Apostolopoulou, A. A., Widmann, A., Rohwedder, A., Pfitzenmaier, J. E., and Thum, A. S. (2013). Appetitive associative olfactory learning in *Drosophila* larvae. *J. Vis. Exp.* 72:4334. doi: 10.3791/4334
- Aso, Y., Hattori, D., Yu, Y., Johnston, R. M., Iyer, N. A., Ngo, T. T.-B., et al. (2014a). The neuronal architecture of the mushroom body provides a logic for associative learning. *eLife* 3:e04577. doi: 10.7554/eLife.04577
- Aso, Y., Sitaraman, D., Ichinose, T., Kaun, K. R., Vogt, K., Belliart-Guérin, G., et al. (2014b). Mushroom body output neurons encode valence and guide memory-based action selection in *Drosophila*. *eLife* 3:e04580. doi: 10.7554/eLife.04580
- Benzer, S. (1967). Behavioral mutants of *Drosophila* isolated by countercurrent distribution. *Proc. Natl. Acad. Sci. U.S.A.* 58, 1112–1119. doi: 10.1073/pnas.58.3.1112
- Berck, M. E., Khandelwal, A., Claus, L., Hernandez-Nunez, L., Si, G., Tabone, C. J., et al. (2016). The wiring diagram of a glomerular olfactory system. *eLife* 5:e14859. doi: 10.7554/eLife.14859
- Brand, A. H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* 118, 401–415.
- Cobb, M. (1999). What and how do maggots smell? *Biol. Rev.* 74, 425–459.
- Demerec, M., and Kaufmann, B. P. (1972). *Drosophila Guide: Introduction to the Genetics and Cytology of *Drosophila Melanogaster**. Washington, DC: Carnegie Institution of Washington.
- Dickinson, A. (2001). The 28th Bartlett Memorial Lecture. Causal learning: an associative analysis. *Q. J. Exp. Psychol. B* 54, 3–25. doi: 10.1080/02724990042000010
- Diegelmann, S., Klagges, B., Michels, B., Schleyer, M., and Gerber, B. (2013). Maggot learning and Synapsin function. *J. Exp. Biol.* 216, 939–951. doi: 10.1242/jeb.076208
- Dudai, Y., Jan, Y. N., Byers, D., Quinn, W. G., and Benzer, S. (1976). *dunce*, a mutant of *Drosophila* deficient in learning. *Proc. Natl. Acad. Sci. U.S.A.* 73, 1684–1688. doi: 10.1073/pnas.73.5.1684
- Fushiki, A., Zwart, M. F., Kohsaka, H., Fetter, R. D., Cardona, A., and Nose, A. (2016). A circuit mechanism for the propagation of waves of muscle contraction in *Drosophila*. *eLife* 5:e13253. doi: 10.7554/eLife.13253
- Gerber, B., and Aso, Y. (in press). “Localization, diversity and behavioral expression of associative engrams in *Drosophila*,” in *Learning Theory and Behavior*, ed R. Menzel (Oxford: Elsevier).
- Gerber, B., and Hendel, T. (2006). Outcome expectations drive learned behaviour in larval *Drosophila*. *Proc. Biol. Sci.* 273, 2965–2968. doi: 10.1098/rspb.2006.3673
- Gerber, B., and Stocker, R. F. (2007). The *Drosophila* larva as a model for studying chemosensation and chemosensory learning: a review. *Chem. Senses* 32, 65–89. doi: 10.1093/chemse/bjl030
- Gerber, B., Biernacki, R., and Thum, J. (2010). “Odor-taste learning in larval *Drosophila*,” in *Drosophila Neurobiology: A Laboratory Manual*, eds B. Zhang M. R. Freeman, and S. Waddell (Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press), 443–455.
- Gerber, B., Biernacki, R., and Thum, J. (2013). Odor-taste learning assays in *Drosophila* larvae. *Cold Spring Harb. Protoc.* 2013.pdb.prot071639. doi: 10.1101/pdb.prot071639
- Gerber, B., Yarali, A., Diegelmann, S., Wotjak, C. T., Pauli, P., and Fendt, M. (2014). Pain-relief learning in flies, rats, and man: basic research and applied perspectives. *Learn. Mem.* 21, 232–252. doi: 10.1101/lm.032995.113
- Gomez-Marin, A., and Louis, M. (2012). Active sensation during orientation behavior in the *Drosophila* larva: more sense than luck. *Curr. Opin. Neurobiol.* 22, 208–215. doi: 10.1016/j.conb.2011.11.008
- Guven-Ozkan, T., and Davis, R. L. (2014). Functional neuroanatomy of *Drosophila* olfactory memory formation. *Learn. Mem.* 21, 519–526. doi: 10.1101/lm.034363.114
- Hammer, M. (1993). An identified neuron mediates the unconditioned stimulus in associative olfactory learning in honeybees. *Nature* 366, 59–63. doi: 10.1038/366059a0
- Harris, K. P., and Littleton, J. T. (2015). Transmission, development, and plasticity of synapses. *Genetics* 201, 345–375. doi: 10.1534/genetics.115.176529
- Heisenberg, M. (2003). Mushroom body memoir: from maps to models. *Nat. Rev. Neurosci.* 4, 266–275. doi: 10.1038/nrn1074
- Heisenberg, M., Borst, A., Wagner, S., and Byers, D. (1985). *Drosophila* mushroom body mutants are deficient in olfactory learning. *J. Neurogenet.* 2, 1–30. doi: 10.3109/01677068509100140
- Hotta, Y., and Benzer, S. (1970). Genetic dissection of the *Drosophila* nervous system by means of mosaics. *Proc. Natl. Acad. Sci. U.S.A.* 67, 1156–1163. doi: 10.1073/pnas.67.3.1156
- Jenett, A., Rubin, G. M., Ngo, T. T., Shepherd, D., Murphy, C., Dionne, H., et al. (2012). A GAL4-driver line resource for *Drosophila* neurobiology. *Cell Rep.* 2, 991–1001. doi: 10.1016/j.celrep.2012.09.011
- Jovanic, T., Schneider-Mizell, C. M., Shao, M., Masson, J. B., Denisov, G., Fetter, R. D., et al. (2016). Competitive disinhibition mediates behavioral choice and sequences in *Drosophila*. *Cell* 167, 858–870. doi: 10.1016/j.cell.2016.09.009
- Kandel, E. R., Dudai, Y., and Mayford, M. R. (2014). The molecular and systems biology of memory. *Cell* 157, 163–186. doi: 10.1016/j.cell.2014.03.001
- Keene, A. C., and Sprecher, S. G. (2012). Seeing the light: photobehavior in fruit fly larvae. *Trends Neurosci.* 35, 104–110. doi: 10.1016/j.tins.2011.11.003
- Kreissl, S., Eichmüller, S., Bicker, G., Rapus, J., and Eckert, M. (1994). Octopamine-like immunoreactivity in the brain and subesophageal ganglion of the honeybee. *J. Comp. Neurol.* 348, 583–595. doi: 10.1002/cne.903480408
- Li, H. H., Kroll, J. R., Lennox, S. M., Ogundeyi, O., Jeter, J., Depasquale, G., et al. (2014). A GAL4 driver resource for developmental and behavioral studies on the larval CNS of *Drosophila*. *Cell Rep.* 8, 897–908. doi: 10.1016/j.celrep.2014.06.065
- Menzel, R. (2014). The insect mushroom body, an experience-dependent recoding device. *J. Physiol. Paris* 108, 84–95. doi: 10.1016/j.jphysparis.2014.07.004
- Neuser, K., Husse, J., Stock, P., and Gerber, B. (2005). Appetitive olfactory learning in *Drosophila* larvae: effects of repetition, reward strength, age, gender, assay type and memory span. *Anim. Behav.* 69, 891–898. doi: 10.1016/j.anbehav.2004.06.013
- O’Kane, C. J., and Gehring, W. J. (1987). Detection in situ of genomic regulatory elements in *Drosophila*. *Proc. Natl. Acad. Sci. U.S.A.* 84, 9123–9127. doi: 10.1073/pnas.84.24.9123
- Ohshima, T., Schneider-Mizell, C. M., Fetter, R. D., Aleman, J. V., Franconville, R., Rivera-Alba, M., et al. (2015). A multilevel multimodal circuit enhances action selection in *Drosophila*. *Nature* 520, 633–639. doi: 10.1038/nature14297
- Oswald, D., and Waddell, S. (2015). Olfactory learning skews mushroom body output pathways to steer behavioral choice in *Drosophila*. *Curr. Opin. Neurobiol.* 35, 178–184. doi: 10.1016/j.conb.2015.10.002
- Pfeiffer, B. D., Ngo, T. T., Hibbard, K. L., Murphy, C., Jenett, A., Truman, J. W., et al. (2010). Refinement of tools for targeted gene expression in *Drosophila*. *Genetics* 186, 735–755. doi: 10.1534/genetics.110.119917
- Rodrigues, V. (1980). Olfactory behavior of *Drosophila melanogaster*. *Basic Life Sci.* 16, 361–371. doi: 10.1007/978-1-4684-7968-3_26
- Rohwedder, A., Wenz, N. L., Stehle, B., Huser, A., Yamagata, N., Zlatić, M., et al. (2016). Four individually identified paired dopamine neurons signal reward in larval *Drosophila*. *Curr. Biol.* 26, 661–669. doi: 10.1016/j.cub.2016.01.012

- Rubin, G. M., and Spradling, A. C. (1982). Genetic transformation of *Drosophila* with transposable element vectors. *Science* 218, 348–353. doi: 10.1126/science.6289436
- Scherer, S., Stocker, R. F., and Gerber, B. (2003). Olfactory learning in individually assayed *Drosophila* larvae. *Learn. Mem.* 10, 217–225. doi: 10.1101/lm.57903
- Schlegel, P., Texada, M. J., Miroshnikov, A., Schoofs, A., Hückesfeld, S., Peters, M., et al. (2016). Synaptic transmission parallels neuromodulation in a central food-intake circuit. *eLife* 5:e16799. doi: 10.7554/eLife.16799
- Schleyer, M., Miura, D., Tanimura, T., and Gerber, B. (2015a). Learning the specific quality of taste reinforcement in larval *Drosophila*. *eLife* 4:e04711. doi: 10.7554/eLife.04711
- Schleyer, M., Reid, S. F., Pamir, E., Saumweber, T., Paisios, E., Davies, A., et al. (2015b). The impact of odor-reward memory on chemotaxis in larval *Drosophila*. *Learn. Mem.* 22, 267–277. doi: 10.1101/lm.037978.114
- Schleyer, M., Saumweber, T., Nahrendorf, W., Fischer, B., von Alpen, D., Pauls, D., et al. (2011). A behavior-based circuit model of how outcome expectations organize learned behavior in larval *Drosophila*. *Learn. Mem.* 18, 639–653. doi: 10.1101/lm.2163411
- Schneider-Mizell, C. M., Gerhard, S., Longair, M., Kazimiers, T., Li, F., Zwart, M. F., et al. (2016). Quantitative neuroanatomy for connectomics in *Drosophila*. *eLife* 5:e12059. doi: 10.7554/eLife.12059
- Schroll, C., Riemensperger, T., Bucher, D., Ehmer, J., Voller, T., Erbguth, K., et al. (2006). Light-induced activation of distinct modulatory neurons triggers appetitive or aversive learning in *Drosophila* larvae. *Curr. Biol.* 16, 1741–1747. doi: 10.1016/j.cub.2006.07.023
- Sivanantharajah, L., and Zhang, B. (2015). Current techniques for high-resolution mapping of behavioral circuits in *Drosophila*. *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* 201, 895–909. doi: 10.1007/s00359-015-1010-y
- Sokolowski, M. B. (2001). *Drosophila*: genetics meets behaviour. *Nat. Rev. Genet.* 2, 879–890. doi: 10.1038/35098592
- Stocker, R. F. (1994). The organization of the chemosensory system in *Drosophila melanogaster*: a review. *Cell Tissue Res.* 275, 3–26. doi: 10.1007/BF00305372
- Stocker, R. F. (2008). Design of the larval chemosensory system. *Adv. Exp. Med. Biol.* 628, 69–81. doi: 10.1007/978-0-387-78261-4_5
- Tedjakumala, S. R., and Giurfa, M. (2013). Rules and mechanisms of punishment learning in honey bees: the aversive conditioning of the sting extension response. *J. Exp. Biol.* 216, 2985–2997. doi: 10.1242/jeb.086629
- Tully, T., and Quinn, W. G. (1985). Classical conditioning and retention in normal and mutant *Drosophila melanogaster*. *J. Comp. Physiol. A* 157, 263–277. doi: 10.1007/BF01350033
- Venken, K. J., Simpson, J. H., and Bellen, H. J. (2011). Genetic manipulation of genes and cells in the nervous system of the fruit fly. *Neuron* 72, 202–230. doi: 10.1016/j.neuron.2011.09.021
- Vosshall, L. B., and Stocker, R. F. (2007). Molecular architecture of smell and taste in *Drosophila*. *Annu. Rev. Neurosci.* 30, 505–533. doi: 10.1146/annurev.neuro.30.051606.094306
- Zwart, M. F., Pulver, S. R., Truman, J. W., Fushiki, A., Fetter, R. D., Cardona, A., et al. (2016). Selective inhibition mediates the sequential recruitment of motor pools. *Neuron* 91, 615–628. doi: 10.1016/j.neuron.2016.06.031

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Michels, Saumweber, Biernacki, Thum, Glasgow, Schleyer, Chen, Eschbach, Stocker, Toshima, Tanimura, Louis, Arias-Gil, Marescotti, Benfenati and Gerber. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.