

## Eumitrioid *Usnea* species (*Parmeliaceae*, lichenized Ascomycota) in tropical South America and the Galapagos

Camille TRUONG and Philippe CLERC

**Abstract:** Eumitrioid *Usnea* species are characterized by a tubular central axis throughout the entire thallus. Four species were identified in South America: *Usnea baileyi* is a (sub-) tropical species occurring both in continental South America and the Galapagos; *U. perplectata* has a scattered distribution worldwide and occurs in continental South America, mainly on the eastern range; *U. flaveola* and the newly described species *U. subflaveola* are both so far endemic to the Neotropical Andes. Distinct chemotypes, as well as anatomical characters, such as the thickness ratio of the medulla or the presence of medullar pigmentation, are diagnostic in delimiting these species. A detailed description of the species is provided together with an identification key.

**Key words:** chemistry, endemism, lichens, morphology, Neotropics, taxonomy

*Accepted for publication 9 December 2012*

---

### Introduction

*Usnea* Adans. is a fruticose lichen genus from the family *Parmeliaceae* (Lecanoromycetes), comprising more than 350 species (Clerc 1998) widely distributed in polar, temperate and tropical regions. The genus is readily distinguished by the shrubby to pendulous thallus, branches holding a central axis and the presence of usnic acid in the cortex. At the species level, the extreme plasticity of morphological characters, in response to environmental parameters, considerably challenges the delimitation of species (Clerc 1998).

Eumitrioid *Usnea* species are characterized by a tubular central axis throughout the entire thallus, sometimes filled with loose internal hyphae. These species were circumscribed as the genus *Eumitria* (Stirton 1882), then under the genus *Usnea* subgenus *Eumitria* (Motyka 1936; Ohmura 2001, 2002). Articus (2004) attempted to re-elevate *Eumi-*

*tria* to the genus level based on phylogenetic lineages, but other authors agreed to keep *Eumitria* as a subgenus (Ohmura & Kanda 2004; Wirtz *et al.* 2006), since the backbone of the phylogeny of *Usnea s. lat.* remained unresolved and phenotypic characters used to circumscribe the lineages were ambiguous. For example, only one species with a tubular central axis (*U. baileyi*) was included in the phylogeny and it clustered in a highly supported clade with a species with a solid central axis (*U. pectinata*). More eumitrioid species need to be added to the phylogeny to understand the circumscription and phylogenetic significance of *Eumitria*. Until that time, we prefer to use the term ‘eumitrioid’ for these species with a tubular central axis throughout the whole thallus, until the monophyly of these species is tested.

Motyka (1936) described several eumitrioid species that are now recognized as morphological variants of the same species, and the diversity of eumitrioid species has been drastically reduced over the years (Swinscow & Krog 1974; Rogers & Stevens 1988; Ohmura 2001). Comprehensive revisions of *Eumitria* species have been completed in Central and East Africa (Swinscow & Krog 1974, 1986; Krog 1994), Australia

---

C. Truong and P. Clerc: Conservatoire et Jardin botaniques de la Ville de Genève, CP 60, 1292 Chambésy, Switzerland; Laboratoire de systématique végétale et biodiversité, Faculté des sciences de l'Université de Genève, CP 60, 1292 Chambésy, Switzerland.  
Email: camille.truong@ville-ge.ch

TABLE 1. *Distribution per country and altitudinal ranges of the species in tropical South America and the Galapagos.*

Species	<i>n</i>	BOL*	BRA	COL	ECU	GAL	PAR	PER	VEN	Altitude† (SAM)	Altitude† (GAL)
<i>U. baileyi</i>	125	+	+	+	–	+	–	+	+	0–2300	150–1100
<i>U. flaveola</i>	9	+	–	–	–	–	–	+	–	1100–2500	–
<i>U. perplectata</i>	21	+	+	–	–	–	+	–	+	0–1500	–
<i>U. subflaveola</i> **	16	–	–	–	–	–	–	+	+	2000–3100	–

\* BOL, Bolivia; BRA, Brazil; COL, Colombia; ECU, Ecuador (continental); GAL, Galapagos; PAR, Paraguay; PER, Peru; VEN, Venezuela; SAM, continental South America; *n*, number of specimens studied; +, presence; \*\* newly described species; †altitudes in [m].

(Rogers & Stevens 1988; Stevens 1999, 2004) and East Asia (Ohmura 2001, 2012). Today, *c.* 8 species are known to occur in East Africa (Swinscow & Krog 1974), one also occurring in Japan (Ohmura 2001, 2012), and two in Australia (Rogers & Stevens 1988) and in North America (Brodo *et al.* 2001; Clerc 2008). South America exhibits a high diversity of *Usnea* species and ongoing taxonomical investigations have been started recently in continental South America and the Galapagos (Rodriguez *et al.* 2011; Truong *et al.* 2011; Truong & Clerc 2012). This study is a further step towards the complete revision of *Usnea* in tropical South America, treating the species with a tubular central axis.

Materials & Methods

This study is based mainly on material collected by the authors in Bolivia, Ecuador (including the Galapagos) and Peru. For a detailed description of the collection sites see Truong *et al.* (2011). In addition, herbarium specimens from the collections of A. Spielmann (SP) and K. Kalb (private hb.) in Brazil, H. Sipman (B) in Colombia and M. E. Hale, Jr. (US) in Venezuela were examined, as well as the Galapagos collections from CDS and COLO. Specimens and types from the following herbaria were included: BM, G, LBL, S and TUR. Table 1 indicates the number of specimens studied for each species, their distribution per country and their altitudinal range. The complete list of specimens examined, including geographical coordinates and ecology, is available upon request to the author.

Morphology of specimens was examined using a stereomicroscope Leica MS5. For a description of characters used in the taxonomy of *Usnea*, see Clerc (1987a, 1998, 2011), Herrera-Campos *et al.* (1998) and Ohmura (2001). The shape of branches and branch segments in longitudinal and cross-section is a difficult character to understand and illustrations to clarify this are available

in Clerc (2011, figs 1–5). Thickness of cortex/medulla/axis was measured in longitudinal sections of branches at  $\times 40$  magnification (on the largest branch above the trunk). The percentage ratio of cortex/medulla/axis of the total branch diameter (CMA) was calculated according to Clerc (1987b). The tubular part of the axis had been used to characterize species (Motyka 1936), but it was not expressed numerically in a formula to enable comparisons among species. Therefore we measured the tubular part of the axis (TBA) and developed a formula to calculate its percentage over the total axis diameter (Fig. 1).

In the description of species, CMA and TBA values are given with their standard deviations and CMA values follow the categories described by Clerc (2011). A synthesis of CMA and TBA values for each species is presented in Fig. 2. Spores were observed under a Leica DM 2000 microscope and at least 10 spores per specimen were measured at  $\times 1000$  magnification.

Chemical analyses were performed by thin-layer chromatography (TLC) in solvents A, B and C, following the method of Culberson & Ammann (1979), with solvent B modified according to Culberson & Johnson (1982). A summary of chemotypes detected for each species is presented in Table 2, with major metabolites highlighted. Metabolites were considered major if their presence was constant within the specimens studied of a given species.

The Species

*Usnea baileyi* (Stirt.) Zahlbr.

*Denkschr. Kaiserl. Akad. Wiss., Math.-Naturwiss. Kl.* **83**: 182 (1909).—*Eumitria baileyi* Stirt., *Scott. Natur.* **6**: 100 (1882); type: Australia, Queensland, near Brisbane, *Bailey* 164 (BM!—lectotype). % C/M/A/TBA: 3.5/3/87/85. Chemistry: usnic, eumitrins A<sub>2</sub> and B, zeorin, salazinic and norstictic acids (Ohmura 2001).

*Usnea antillarum* (Vain.) Zahlbr., *Cat. Lich. Univ.* **6**: 536 (1930).—*Eumitria antillarum* Vain., *Ann. Acad. Sci. Fem., Ser. A* **6** (7): 9 (1915); type: Antilles, Ins. St. Thomas, Signalhill, 1876, *Eggers* (TUR V00328!—holotype, LBL!—isotype). % C/M/A/TBA: 3.5/1.5/90/68. Chemistry: usnic, norstictic and salazinic acids and eumitrins (Rogers & Stevens 1988).




FIG. 1. Measurements and calculations of % cortex/medulla/axis (CMA), ratio A/M and % tubular axis (TBA) in longitudinal section of branch (at the widest branch diameter above first ramification). In colour online.




FIG. 2. % cortex/medulla/axis of total branch diameter and % tubular axis of total axis diameter. Standard deviations as wide bars; extreme values as lines; numbers in parentheses = number of specimens measured.

TABLE 2. Secondary metabolites (columns) detected by TLC in the medulla of eumitrioid species.

Species	<i>n</i>	DIF*	EUM	NOR	SAL	ZEO	UT8
<i>U. baileyi</i>	62	–	+	±	±	+	–
<i>U. flaveola</i>	9	–	–	–	–	–	+
<i>U. perplectata</i>	20	+	–	–	–	–	–
<i>U. subflaveola</i>	16	–	–	±	–	–	+

\*DIF, diffractaic; EUM, eumitrins; NOR, norstictic; SAL, salazinic acid; ZEO, zeorin; UT8, unidentified tri-terpenoid UT8 (RF classes A/B/C: 1-2/4-5/2, UV+ white after charring); *n*, number of specimens studied; +, presence constant within species; ±, presence variable among specimens within a species; grey boxes indicate major metabolites, in which presence was constant within the specimens studied of a given species.

*Usnea implicita* (Stirt.) Zahlbr., *Denkschr. Kaiserl. Akad. Wiss., Math.-Naturwiss. Kl.* **83**: 182 (1909).—*Eumitria implicita* Stirt., *Scott. Natur.* **6**: 100 (1882); type: Madeira, W. Funchal, *Payne* (BM!—lectotype). % C/M/A/TBA: 5.5/2.5/84/63. Chemistry: usnic and norstictic acids, eumitrins A<sub>1</sub>, A<sub>2</sub> and B, zeorin (Ohmura 2001).

*Usnea inanis* Motyka syn. nov., *Lich. Gen. Usnea Stud. Monogr. Pars Syst.* **1**: 58 (1936); type: Brazil, Matto Grosso, Serra da Chapada, Buriti, 1894, *Malme* (S!—holotype, LBL!—isotype). % C/M/A/TBA: 4.5/3.5/85/77. Chemistry: usnic, eumitrin A, zeorin and norstictic acid.

(Figs 3A & B, 4A–C)

**Nomenclatural notes.** Three specimens are represented on the herbarium sheet containing the holotype specimen of *U. inanis* Motyka: we marked them as (a), (b) and (c). Thallus (a) was designated by Motyka’s handwriting as the holotype and corresponds well morphologically and chemically to *U. baileyi* (eumitrins), whereas thallus (c) corresponds to *U. perplectata* (diffractaic acid). An isotype specimen found in LBL also corresponds to *U. baileyi*.

Only names that have been cited for South America are included in the above list of synonyms. For a complete list of synonyms, see former publications (Swinscow & Krog 1974; Rogers & Stevens 1988; Ohmura 2001).

**Diagnostic notes.** For a detailed description of this taxon, see Ohmura (2001). *Usnea baileyi* is a sorediate species characterized by a stiff, erect-shrubby to subpendulous thallus, branches ± irregular, often sinuous towards the apices, with slightly to strongly flattened

segments (Fig. 4B). The base is concolorous or blackish on the first mm below the first ramification (Fig. 4A). Branches are abundantly covered by minute soralia growing from the cortex *ad initio* or from low tubercles (Fig. 3A). Soralia are slightly raised, remaining plane, crowded on terminal branches, usually without fusing, with numerous short isidiomorphs (Fig. 3B). The cortex is very thin (3.0–5.5%), matt on the thallus surface but sometimes shinier in section. The medulla is extremely thin (1.5–3.0%), almost invisible except for the pink-red pigmentation throughout (Fig. 4C). The axis is extremely large (83.0–91.5%), with a large tubular section (67–83%) filled with loose hyphae. *Usnea baileyi* is characterized by the presence of eumitrins and zeorin detected by TLC in the medulla.

**Variation.** The density of ramifications, fibrils and isidiomorphs varies among individuals. Rarely, soralia may fuse in an irregular ‘soralium’ on terminal branches, from the aggregation of several minute soralia. In addition to eumitrins and zeorin, norstictic acid (*n* = 44) and rarely salazinic acid (*n* = 5) were also detected by TLC in the medulla of South American specimens and, as a consequence, the K reaction in the medulla is variable.

**Differentiation.** *Usnea perplectata* differs in the branch anatomy in longitudinal section (Fig. 2): it has a thinner axis than *U. baileyi* and a larger medulla, with a distinctly periaxial pigmentation. In addition, it has a somewhat thicker cortex and a thinner tubu-




FIG. 3. A & B, *Usnea baileyi* (Truong 1446); A, tubercles turning into minute and slightly raised soralia; B, soralia circular and plane, without a cortical margin, crowded on terminal branches but without fusing. C & D, *Usnea perplectata*; C, minute to slightly elongated soralia bursting along cracks and ridges of branches (Ferrero 1062); D, soralia of irregular outline, plane to slightly capitate, without a cortical margin, crowded and fusing on terminal branches (Bach 373). E & F, *Usnea subflaveola*; E, fiberclles abundantly distributed on the branches (Truong 1783); F, soralia distinctly stipitate, circular and often capitate at maturity, with a distinct cortical margin, crowded on terminal branches but without fusing (Truong 1707). Scales: A–F = 500  $\mu$ m. In colour online.




FIG. 4. A–C, *Usnea baileyi*; A, trunk blackish (Truong 1446); B, flattened branches abundantly covered by minute soralia (Clerc 08-319); C, thin cortex; extremely thin medulla, pigmented throughout; very large axis with large tubular section (Clerc 08-319). D–F, *Usnea perplectata*; D, trunk concolorous with thin annular cracks (Rodríguez 0261B); E, ridged branches abundantly covered by soralia of irregular outline (Ferrero 1062); F, cortex thicker than (C); medulla thicker than (C) with periaxial pigmentation; large axis with large tubular section (Bach 373). Scales: A–F = 500  $\mu$ m. In colour online.

lar section of axis, although these values overlap in the two species. *Usnea perplectata* also has slightly ridged branches (flattened in *U. baileyi*), with few to numerous fiberclles, especially visible on basal branches. Diffraction acid is detected by TLC in the medulla instead of eumitrins in *U. baileyi*.

**Ecology and distribution.** This is primarily a corticolous species, growing on a wide variety of trees and shrubs (including *Rhizophora*

and cacti), occasionally also lignicolous (fencepost) or saxicolous. It is very frequent in the Galapagos, occurring from the arid to the humid zone (150–1100 m). In continental South America, it is one of the few *Usnea* species occurring at low altitudes (0–2300 m). It is found in open pastures, humid evergreen forests and mangroves, but also in the semi-arid climate of seasonally dry tropical forests and savannas of Brazil, Colombia and Venezuela. In Australia, this species was

reported to be restricted to warm and moist places (Rogers & Stevens 1988). From these observations we can deduce that *U. baileyi* has a rather wide ecological range from humid to arid places. This species has a world-wide distribution in (sub-) tropical regions of Asia (Ohmura 2001), Australia (Rogers & Stevens 1988), East Africa (Swinscow & Krog 1974) and North and South America (Brodo *et al.* 2001; Clerc 2008).

*Selected specimens examined.* **Bolivia:** Beni: 20 km al SW del km 12 Yucumo – Rurrenabaque, 900 m, 1997, *Bach et al.* (B).—**Brazil:** Minas Gerais: Bei Lagoa Dourada etwa 50 km nordöstlich von São João del Rei, 1000 m, 1978, *Kalb & Plöbst* (private hb.). *Rio Grande do Sul:* Herveiras, 570 m, 2004, *Spielmann & Putzke* 6308 (SP). *São Paulo:* near Praia de Pernibe, near Itanhaëna, 0 m, 1979, *Sipman* 13501 (B).—**Colombia:** Amazonas: Araracuara, on S-side of river Caquetá, 350 m, 1988, *Sipman & Duivenvoorden* 27856 (B). *Cauca:* Los Robles, campus of Fundacion Universitária, 1750 m, 1986, *Sipman et al.* 32738 (B). *Cundinamarca:* Laguna de Pedro Palo, 2000 m, 1984, *Aguirre & Sipman* 5640 (B).—**Ecuador:** Galapagos Islands: Isabela, mirador el Mango, 161 m, 2008, *Truong* 1191 (G, CDS); Pinta, along the trail up to the summit from the S-coast, 369 m, 2007, *Bungartz* 5850 (CDS); Pinzón, on the W-slope of the highest mountain, 310 m, 2006, *Aptroot* 64105 (CDS); San Cristobal, Cerro Colorado summit, 159 m, 2008, *Truong* 1420 (G, CDS); Santa Cruz, dirt road to Mina Granillo, 547 m, 2008, *Truong* 1302 (G, CDS); Santiago, Cerro Gavilan, 680 m, 2006, *Bungartz* 4686 (CDS).—**Peru:** Cusco: Valle de Lacco, 1928, *Herrera* 2066 (US).—**Venezuela:** Amazonas: en la cumbre de la laja de Carestia, bajo Sanariapo, 150 m, 1942, *Williams* 16038 (US). *Aragua:* Arriba de Guanitas, parque nacional, 950 m, 1939, *Williams* 12213 (US). *Distrito Federal:* Silla de Caracas, 1700 m, 1951, *Vareschi* 57 (LBL). *Lara:* Morán, bordeando a las cabeceras del río Tocuyo, 1900 m, 1974, *Steyermark & Carreño Espinoza* 111081 (US). *Merida:* La Carbonera, sector San Rafael del Macho, 2200 m, 1976, *Hale* 46631 (US). *Miranda:* Los Guayabitos, 1400 m, 1955, *Vareschi* 3935 (LBL). *Sucre:* Península de Paria, Cerro de Río Arriba, 700 m, 1966, *Steyermark y Rabe* 96276 (US).

## Usnea flaveola Motyka

*Lich. Gen. Usnea Stud. Monogr. Pars Syst.* 1: 52 (1936); type: Peru, Sandia, 2100 m, 1902, *Weberbauer* (LBL!—holotype, isotype). % C/M/A/TBA: 7/7.5/71.5/75. Chemistry: usnic acid and unknown triterpenoids UT8.

(Fig. 5D–F)

*Nomenclatural note.* Motyka (1936) mentions that the holotype specimen of *U. flaveola* was deposited in B, but we found it within his collections in LBL.

*Thallus* stiff, erect-shrubby to subpendulous (to c. 10 cm long); *ramifications* ± anisotomic-dichotomous; *trunk* concolorous with the branches, without annulations or rarely with very thin annular cracks (Fig. 5D); *branches* tapering to irregular; *branch segments* terete to flattened or ridged, sometimes almost angular; *lateral branches* not constricted at ramification; *foveoles* and *maculae* absent; *papillae* and *tubercles* absent or scarce (not to be confounded with young fibrils); *fibrils* slender (to 5 mm long), abundantly and regularly distributed on the branches, looking like cylindrical papillae crowded on the cortex surface at a juvenile stage; *fibercles* usually numerous (Fig. 5E), especially on basal branches; *soralia* absent; *cortex* shiny in section, moderately thin to moderately thick (6.5–10.5%); *medulla* white (Fig. 5F), compact and thin (3.5–9.0%); *axis* thick (66.5–74.5%) with a moderate tubular section (52–73%) loosely filled with thin hyphae and often also chondroid strands of hyphae, axis rarely solid (not tubular) on thin terminal branches.

*Apothecia* usually present and often numerous, (sub-) terminal, large at maturity (to c. 15 mm diam.), with few to numerous long cilia; *spores* hyaline, obovoid, 8.5–10.5 × 5–7 µm (*n* = 45).

*Pycnidia* rare, usually visible when apothecia are absent, appearing as small hemispherical protuberances on terminal branches.

*Diagnostic notes.* *Usnea flaveola* is a non-sorediate species (usually with apothecia) characterized by an erect to subpendulous thallus, with somewhat irregular branches abundantly covered by fibrils and fibercles. Clerc & Herrera-Campos (1997) described fibercles as stipitate scars left by the breaking away of fibrils, probably functioning as sterile propagules, sometimes exposing the central axis of the fallen fibril. The axis is large with a moderately large tubular section and the medulla is thin and white (without pigmentation), reacting K– (unidentified triterpenoids UT8: RF classes A/B/C: 1-2/4-5/2, UV+ white after charring). Triterpenoids seem to be frequent in Neotropical *Usnea* specimens and are useful for identifying several species, for example *U. rubricornuta*




FIG. 5. A–C, *Usnea subflaveola*; A, trunk concolorous, sometimes with thin annular cracks; branch segments almost angular (Truong 2783); B, tapering branches abundantly covered by fiberclles (Truong 1783); C, cortex moderately thin; medulla thin, without pigmentation; axis large, with a moderately large tubular section (Truong 1707). D–F, *Usnea flaveola*; D, trunk concolorous, without annular cracks (Truong 3140); E, branches covered by fibrils and fiberclles, holding apothecia (Bach 325); F, cortex moderately thin; medulla thinner than (C), without pigmentation; axis large, with a moderately large tubular section (Truong 3140). Scales: A–F = 500  $\mu$ m. In colour online.

Truong & P. Clerc and *U. steineri* Zahlbr. (Truong *et al.* 2011).

**Variation.** The irregularity of branch diameter, as well as the shape of branch segments (terete to flattened or ridged, sometimes even angulate) vary mostly with the development stage of the thallus. The abundance of fibrils and fiberclles also vary among individuals; nevertheless, fiberclles are always present and characteristic of this species.

**Differentiation.** The tubular axis, the absence of medullar pigmentation and the absence of soralia readily distinguish this species. It is the only non-sorediate eumitrioid species occurring in South America so far (see also *U. trullifera* Motyka in the paragraph treating doubtful species).

**Ecology and distribution.** This is mostly a corticolous species, occasionally also found on rocks. It occurs in humid and somewhat

open places within evergreen forest, mountain cloud forest or in the vicinity of the forest. It is so far endemic to the Neotropical Andes, at moderate altitudes (1100–2500 m).

*Selected specimens examined.* **Bolivia:** La Paz: 15 km de Charazani hacia Apolo, 2400 m, 1997, *Bach et al.* 252 (B). **Cochabamba:** Parque Nacional Carrasco, Sehuencas hasta Monte Puncu, 2500 m, 2007, *Truong* 3140 (G).—**Peru:** *Junin:* Distrito de San Ramon, carretera arriba de la catarata Tirol, 1139 m, 2007, *Truong* 2676 (G).

### **Usnea perplectata** Motyka

*Lich. Gen. Usnea Stud. Monogr. Pars Syst.* 1: 55 (1936); type: Paraguay, Gran Chaco, ad Rio Negro e ad Capernicians, 1883, *Malme* (S!—holotype). % C/M/A/TBA: 6.5/16/55/41. Chemistry: usnic, diffractaic, traces of 4-O-demethylbarbatic acids and associated compounds.

*Usnea elata* Motyka *syn. nov.*, *Lich. Gen. Usnea Stud. Monogr. Pars Syst.* 1: 57 (1936); type: [Congo], Mpala, ad litus occid. Lacus Tanganjika, 1898, *Guilleme* (TUR!—holotype). C/M/A/TBA: unknown (small fragment of branch). Chemistry: usnic and diffractaic acids (Rogers & Stevens 1988).

*Usnea subcavata* Motyka *syn. nov.*, *Lich. Gen. Usnea Stud. Monogr. Pars Syst.* 1: 57 (1936); type: Brazil, Sao Paulo, prope S. Amaro in circuito urbis S. Paulo, 800 m, 1901, *Schiffner* (W—holotype, LBL!—isotype). % C/M/A/TBA: 4.5/8.5/73.5/67. Chemistry: usnic, diffractaic, traces of barbatic, 4-O-demethylbarbatic acids and associated compounds.

(Figs 3C & D, 4D–F)

*Nomenclatural note.* Clerc (2008) synonymized *U. perplectata* into *U. baileyi*. However, the current study treats *U. perplectata* as a distinct species.

*Thallus* relatively stiff, subpendulous to pendulous (to c. 25 cm long); *ramifications* ± anisotomic-dichotomous; *trunk* concolorous with the branches or rarely blackish on the first mm (below first ramification), usually with thin annulations extending along the basal branches (Fig. 4D); *branches* of slightly to strongly irregular diameter; *branch segments* terete to slightly ridged (Fig. 4E); *lateral branches* not constricted at ramification; *foveoles* and *maculae* absent; *papillae* absent or scarce; *tubercles* few to abundant on main branches; *fibrils* slender (to 5 mm long), scattered on the thallus; *fibercles* few to abundant on basal branches; *soralia* growing from the cortex *ad initio* (cracks, along ridges of branches, Fig. 3C) or from tubercles, at once slightly raised, usually remaining minute to

slightly enlarging (less than the branch half-diameter), often elongated (along ridges of branches) or of irregular outline, without a cortical margin, plane to slightly capitate at maturity, often crowded on terminal branches and aggregating in irregular ‘soralium’ (Fig. 3D), with few to numerous isidiomorphs, rarely growing into isidiofibrils; *cortex* shiny in section, thin to moderately thin (4.5–7.5%); *medulla* compact and thin (6.5–13.5%), with a pink to orange, periaxial pigmentation (Fig. 4F); *axis* large (59.5–75.5%), with a moderately large tubular section (44–78%) loosely filled with thin hyphae that are often yellow pigmented and reacting C+ yellow-orange.

*Apothecia* rare (*n* = 1), lateral.

*Pycnidia* not seen.

*Diagnostic notes.* This species is characterized by a stiff thallus with slightly ridged branches and minute to slightly elongated soralia, crowded and aggregating on terminal branches, with few to numerous isidiomorphs. The medulla is thin and compact, with a characteristic pink to orange pigmentation, located at the periphery of the axis. Diffractaic acid was detected by TLC in the medulla.

*Variation.* The density of tubercles, fibrils, fibercles and isidiomorphs varies among individuals. The trunk is usually concolorous and distinctly annulated, but it is rarely brownish to blackish (below first ramification) and without annulations.

*Differentiation.* See under *U. baileyi*.

*Ecology and distribution.* This is mostly a corticolous species, occasionally also found on rocks. It occurs in continental South America, mainly on the Eastern range. As *U. baileyi*, it is one of the few *Usnea* species occurring at low altitudes (0–1500 m). It is found in humid places, within evergreen forest, coastal forest (*restinga*) or in the vicinity of the forest, but also in drier places such as the Brazilian *cerrado*. It also occurs in North America (Brodo *et al.* 2001), East Africa and Australia (*sub U. elata*).

*Selected specimens examined.* **Argentina:** Corrientes: Capital San Cayetano, 1976, *Ferraro* 1062 (G). **Formosa:** Dpto. Lahisi, ruta nacional 11, 2007, *Rodriguez* 281B

(G). *Misiones*: Bei den Wasserfällen Victoria am Rio Iguazu, 1933, *Hosseus* (LBL).—**Bolivia**: *Beni*: 16 km al SW del km 12 Yucumo–Rurrenabaque, 750 m, 1997, *Bach et al.* 373 (B).—**Brazil**: *Mato Grosso*: Serra dos Coroados, Naturschutzgebiet der evangelischen Schule von Buriti, 600 m, 1980, *Kalb* (private hb.). *Santa Catarina*: Ilha de Santa Catarina, Praia do Forte, 3 m, 1980, *Kalb* (private hb.). *São Paulo*: Reserva biológica de Mogi-Guaçu, fazenda Campininha, 610 m, 2008, *Jungbluth* 1755 (SP).—**Tanzania**: Sud-Pale Gebirge, bei Vudu, 1800 m, 1958, *Schütz* (LBL).—**Paraguay**: In vicinity urbis Assuntion, *Aminsris* (LBL).—**USA**: *Florida*: Standford, 1924, *Rapp* (LBL).—**Venezuela**: *Aragua*: Choroni, Peñon Blanco, 1200 m, 1954, *Vareschi* 3603 (LBL). *Zulia*: Selva de Geonema, 1500 m, 1954, *Vareschi* 3172 (LBL).

***Usnea subflaveola* Truong & P. Clerc  
sp. nov.**

Mycobank No.: MB 564913

Thallus sorediate, fibrils and fiberclles abundant, medulla unpigmented, reacting K– (triterpenoids), axis tubular to the apices.

Type: Venezuela, Merida, La Carbonera, Merida–La Azulita road, 2200 m, remnants of forest in open pasture, fence post, 1976, *Hale* 47542 (USM—holotype; G—isotype). % C/M/A/TBA: 6.5/18.5/54. Chemistry: usnic and norstictic acids, unidentified triterpenoids UT8.

(Figs 3E & F, 5A–C, 6)

*Thallus* subpendulous to pendulous (to c. 30 cm long), rarely erect-shrubby; *ramifications* mostly anisotomic-dichotomous; *trunk* concolorous with the branches, without annulations or rarely with thin annular cracks (Fig. 5A); *branches* tapering to slightly irregular; *branch segments* terete to slightly ridged or angular, sometimes slightly inflated; *lateral branches* not constricted at ramification; *foveoles* and *maculae* absent; *papillae* and *tubercles* absent or scarce (not to be confounded with young fibrils); *fibrils* slender (to 5 mm long), scattered (mainly on basal branches) to abundantly distributed on the branches; *fiberclles* usually numerous (Fig. 3E & 5B), especially on basal branches; *soralia* developing at the top of fiberclles thus being distinctly stipitate, minute to circular, circular, enlarging almost to the branch diameter on terminal branches, convex or rarely slightly excavate at maturity, often crowded on terminal branches but remaining well-delimited with a thin cortical margin (Fig. 3F), with numerous isidiomorphs, some-

times growing into isidiofibrils; *cortex* shiny in section, moderately thin to moderately thick (6.5–9.5%); *medulla* white (Fig. 5C), compact and thin (8.0–15.5%); axis large (52.0–69.5%) with a moderate tubular section (47–62%), loosely filled with thin hyphae and sometimes chondroid strands of hyphae, axis rarely solid (not tubular) on small terminal branches.

*Apothecia* and *pycnidia* not seen.

*Diagnostic notes.* This is a sorediate species characterized by a (sub)pendulous thallus, with tapering to slightly irregular branches abundantly covered by fibrils and fiberclles, from which arise stipitate soralia, often crowded on terminal branches but remaining well-delimited with a distinct cortical margin. The medulla is thin and white (without pigmentation), reacting K– (unidentified triterpenoids UT8), rarely faintly K+ (norstictic acid).

*Variation.* The abundance of fibrils and fiberclles may vary among individuals, but fiberclles are conspicuously present, at least towards the basal branches. The development of soralia and isidiomorphs varies among individuals or in response to infestation by parasitic fungi, as is often the case in *Usnea*. In addition to UT8, norstictic acid ( $n = 6$ ) was sometimes detected by TLC in the medulla.

*Differentiation.* The absence of medullar pigmentation and the presence of circular soralia growing from fiberclles readily distinguish this species. Apart from the presence of soralia, it shares with *U. flaveola* a similar morphology (presence of fiberclles, absence of medullar pigmentation) and the same chemotype. CMA values show slight differences between the two species, *U. subflaveola* having a thicker medulla and a thinner axis than *U. flaveola*, although these values overlap in the two species (Fig. 2). The corroboration of molecular tools will be needed to distinguish these two taxa.

In East Africa, *U. cristata* Motyka is another sorediate eumitrioid species without medullar pigmentation. The holotype specimen of *U. cristata* deposited in W is apparently missing (Swinscow & Krog 1974), but



FIG. 6. *Usnea subflaveola*; A, holotype; B, isotype. Scale = 1 cm. In colour online.

we found an isotype in Motyka's collections in LBL. It doesn't correspond to *U. subflaveola*: in *U. cristata*, branches are deformed, with strongly ridged segments and the presence of foveoles. Soralia arise irregularly along the ridges of branches or from thick fiberclles. They remain minute to slightly elongated, with an irregular outline and numerous isidiomorphs.

**Ecology and distribution.** This is mostly a corticolous species, occasionally also found on dead wood (fence post). It occurs in humid places within mountain cloud forest or in open places in the vicinity of the forest, such as deforested zones of *matorral*, remnants of forest within pastures or isolated trees. It is so far endemic to the Neotropical Andes, at a moderately high altitude (2000–3100 m).

**Selected specimens examined.** **Peru:** *Cajamarca:* Ruinas de Kuelap, 3100 m, 2007, *Truong* 1870 (G). *Huancayo:*

Bosque de Carpish, 2424 m, 2007, *Truong* 2738 (G). *Pasco:* Parque Nacional Yanachaga-Chemillén, camino hasta el refugio El Cedro, 2309 m, 2007, *Truong* 2444 (G).—**Venezuela:** *Merida:* La Carbonera, sector San Rafael del Macho, 2200 m, 1976, *Hale* 46603 (US). *Tachira:* Slopes of Pico Banberas, páramo de Tama, 2600 m, 1975, *Hale & Lopez-Figueiras* 45208 (US).

## Doubtful Species

### *Usnea trullifera* Nyl.

*Lich. Gen. Usnea Stud. Monogr. Pars Syst.* 1: 65 (1936); type: Tahiti, 1875, *Moseley* (BM)—holotype). % C/M/A/TBA: 4/3.5/85/71. Chemistry: usnic, salazinic and norstictic acids, zeorin.

This is a non-sorediate species (usually with apothecia), characterized by the presence of high tubercles abundantly covering the branches and often eroded at the top. CMA values are similar to *U. baileyi*, with a very thin and pale pink to orange pigmented medulla. In his description, Motyka (1936)

mentions specimens from Tahiti, East Africa and Brazil, most of which were hosted in B and are probably lost today. We found one Brazilian specimen identified by Motyka in LBL. It is composed of a small fragment of branch, holding apothecia (without soralia), but lacking the trunk and basal branches. Apart from the presence of apothecia, this specimen resembles *U. baileyi*, but it is too small to be accurately identified. Despite studying more than 100 South American specimens of *U. baileyi*, we have never ob-

served apothecia in this species. In addition, protocetraric acid and zeorin were detected by TLC in the medulla of this Brazilian specimen. Protocetraric acid has been detected in Asiatic *U. baileyi* specimens (Ohmura 2001), but we have never found it in South American specimens.

*Selected specimens examined.* **Brazil:** Mato Grosso: Caracas, 1908, *Hoehne* (LBL).—**Cameroon:** Kamerun-berg, 1800 m, 1958, *Knorr* (LBL).—**Tahiti:** 1847, *Vesco* (LBL).—**Uganda:** Kasatoro Forest Kigezi, 1800–2500 m, 1947, *Dale* 39 (LBL).

### Key to eumitrioid *Usnea* species from tropical South America and the Galapagos

- 1 Medulla with a pink, orange or red pigmentation. . . . . 2  
Medulla white (not pigmented) . . . . . 3
- 2(1) Medulla extremely thin (<5%); pigmentation pink-red throughout the entire medulla; branch segments terete to flattened . . . . . ***U. baileyi***  
Medulla thicker (>5%); pigmentation pink to orange, distinctly periaxial; branch segments terete to ridged . . . . . ***U. perplectata***
- 3(1) Soralia present, developing at the top of fiberclles, distinctly stipitate, often crowded on terminal branches, remaining well-delimited with a distinct cortical margin . . . . . ***U. subflaveola***  
Soralia absent and apothecia usually present, if not pycnidia often visible as hemispherical protuberances on terminal branches; fibrils and fiberclles abundantly distributed on the branches . . . . . ***U. flaveola***

Fieldwork was supported by the Conservatoire et Jardin Botaniques de la Ville de Genève, a grant “Augustin Lombard” from the Société de Physique et d’Histoire Naturelle de Genève and a grant from the Amazon Conservation Association. Fieldwork was realized with the collaboration of the herbaria CDS, LPB, QCNE, USM and the Wayqecha Biological Station (Peru). F. Bungartz, M. Herrera-Campos, C. Aldana Munguia, A. Cuba Villena, M. F. Collaguazo, Y. Manami, A. Ramirez-Ordaza, P. Rodriguez, L. Salcedo Valdez and A. X. Shuguli participated in the field trips. The Ministries of Environment of Bolivia, Peru, Ecuador and the Galapagos provided collection permits. The staff in national parks kindly supported our investigation by providing guides and/or lodging. A. Tehler and W. Mulenko gave us a warm welcome during our visits to S and LBL, and curators of the cited herbaria kindly sent specimens on loan. Florian Visentin helped with TLC analysis and M. Price corrected the English.

### REFERENCES

Articus, K. (2004) *Neuropogon* and the phylogeny of *Usnea* s.l. (*Parmeliaceae*, lichenized Ascomycetes). *Taxon* **53**: 925–934.

- Brodo, I. M., Sharnoff, S. D. & Sharnoff, S. (2001) *Lichens of North America*. New Haven: Yale University Press.
- Clerc, P. (1987a) On the morphology of soralia in the genus *Usnea*. *Bibliotheca Lichenologica* **25**: 99–102.
- Clerc, P. (1987b) Systematics of the *Usnea fragiliscens* aggregate and its distribution in Scandinavia. *Nordic Journal of Botany* **7**: 479–495.
- Clerc, P. (1998) Species concepts in the genus *Usnea* (lichenized Ascomycetes). *Lichenologist* **30**: 321–340.
- Clerc, P. (2008) *Usnea*. In *Lichen Flora of the Greater Sonoran Desert Region* Vol. 3 (T. H. Nash III, C. Gries & F. Bungartz, eds.): 302–335. Tempe, Arizona: Lichens Unlimited, Arizona State University.
- Clerc, P. (2011) *Usnea*. In *Nordic Lichen Flora* Vol. 4 (A. Thell & R. Moberg, eds): 107–127. Uddevalla: Nordic Lichen Society.
- Clerc, P. & Herrera-Campos, M. A. (1997) Saxicolous species of *Usnea* subgenus *Usnea* (lichenized Ascomycetes) in North America. *Bryologist* **100**: 281–301.
- Culbertson, C. F. & Ammann, K. (1979) Standardmethode zur Dünnschichtchromatographie von Flechtensubstanzen. *Herzogia* **5**: 1–24.

- Culberson, C. F. & Johnson, A. (1982) Substitution of methyl tert-butyl ether for diethyl ether in the standardized thin-layer chromatographic method for lichen products. *Journal of Chromatography* **238**: 483–487.
- Herrera-Campos, M. A., Clerc, P. & Nash III, T. H. (1998) Pendulous species of *Usnea* from the temperate forests in Mexico. *Bryologist* **101**: 303–329.
- Krog, H. (1994) New observations on *Usnea* subgenus *Eumitria* in eastern and central Africa. In *Proceedings of the XIII plenary meeting AETFAT* Vol. 2 (J. Seyani & A. Chikuni, eds): 813–821. Malawi: National Herbarium and Botanic Gardens.
- Motyka, J. (1936) *Lichenum Generis Usnea Studium Monographicum. Pars Systematica* (Vol. 1). Leopoli: privately printed.
- Ohmura, Y. (2001) Taxonomic study of the genus *Usnea* (lichenized Ascomycetes) in Japan and Taiwan. *Journal of the Hattori Botanical Laboratory* **90**: 1–96.
- Ohmura, Y. (2002) Phylogenetic evaluation of infrageneric groups of the genus *Usnea* based on ITS regions in rDNA. *Journal of the Hattori Botanical Laboratory* **92**: 231–243.
- Ohmura, Y. (2012) A synopsis of the lichen genus *Usnea* (Parmeliaceae, Ascomycota) in Taiwan. *Memoirs of the National Museum of Nature and Science* **48**: 91–137.
- Ohmura, Y. & Kanda, H. (2004) Taxonomic status of section *Neuropogon* in the genus *Usnea* elucidated by morphological comparisons and ITS rDNA sequences. *Lichenologist* **36**: 217–225.
- Rodriguez, J. M., Estrabou, C., Truong, C. & Clerc, P. (2011) The saxicolous species of the genus *Usnea* subgenus *Usnea* (Parmeliaceae) in Argentina and Uruguay. *Bryologist* **114**: 504–525.
- Rogers, R. W. & Stevens, G. N. (1988) The *Usnea baileyi* complex (Parmeliaceae, lichenised Ascomycetes) in Australia. *Australian Systematic Botany* **1**: 355–361.
- Stevens, G. N. (1999) A revision of the lichen family *Usneaceae* in Australia. *Bibliotheca Lichenologica* **72**: 1–128.
- Stevens, G. N. (2004) *Usneaceae*. In *Flora of Australia* Vol. 56A, Lichens 4 (P. M. McCarthy & K. Mallett, eds): 78–98 & 107–115. Melbourne: ABR/CSIRO.
- Stirton, J. (1882) Notes on the genus *Usnea* with descriptions of new species. *Scottish Naturalist* **6**: 292–297.
- Swinscow, T. D. V. & Krog, H. (1974) *Usnea* subgenus *Eumitria* in East Africa. *Norwegian Journal of Botany* **21**: 165–185.
- Swinscow, T. D. V. & Krog, H. (1986) *Usnea antiqua* sp. nov. described from Tanzania. *Lichenologist* **18**: 293–295.
- Truong, C. & Clerc, P. (2012) The lichen genus *Usnea* (Parmeliaceae) in tropical South America: species with a pigmented medulla, reacting C+ yellow. *Lichenologist* **44**: 1–13.
- Truong, C., Bungartz, F. & Clerc, P. (2011) The lichen genus *Usnea* (Parmeliaceae) in the tropical Andes and the Galapagos: species with a red-orange cortical or subcortical pigmentation. *Bryologist* **114**: 477–503.
- Wirtz, N., Printzen, C., Sancho, L. & Lumbsch, H. T. (2006) The phylogeny and classification of *Neuropogon* and *Usnea* (Parmeliaceae, Ascomycota) revisited. *Taxon* **55**: 367–376.