

# Two novel mutations: 5108delAG and 5816insG in the NF1 gene detected by SSCP analysis

Jun Zhong, Roland Spiegel\*, Eugen Boltshauser<sup>1</sup> and Werner Schmid

Institute of Medical Genetics, University of Zurich, Raemistrasse 74, CH-8001 Zurich and <sup>1</sup>Division of Paediatric Neurology, Department of Paediatrics, University of Zurich, Switzerland

Received May 26, 1993; Revised and Accepted July 14, 1993

Neurofibromatosis type I (NF1) is one of the most common autosomal dominant disorders in man and is characterized by *café-au-lait* spots, multiple neurofibromas, iris hamartomas (Lisch nodules), orthopedic problems, learning disabilities, and an increased risk of malignancy. The NF1 gene consists of 49 exons, and extends over a genomic distance of approximately 300 kb on human chromosome 17q11.2. It encodes for neurofibromin, a 2818 amino acid protein which is capable of down-regulating *ras* by stimulating its intrinsic GTPase activity (1–4). The mutation rate for NF1 is high (1 in 10 000), and about half of all cases are due to new mutations (5). In order to characterize the individual mutations in the NF1 gene, we investigated genomic DNA samples from 82 Swiss NF1 families (including 31 sporadic cases) by single strand conformation polymorphism (SSCP) analysis. Here we describe two novel mutations found in exons 28 and 31 of the NF1 gene. The other 80 families investigated for these two exons showed no SSCP alterations.

Exon 28 and exon 31 were amplified by the polymerase chain reaction (PCR) using primers described by Upadhyaya *et al.* (5) and Cawthon *et al.* (2) on a Perkin Elmer 9600 thermal cycler. For exon 28, PCR was performed at 94°C for 1 min followed by 30 cycles at 94°C 30 sec, 52°C 30 sec, and 72°C 30 sec, with final elongation at 72°C for 10 min. For exon 31, the amplification conditions were: 94°C for 1 min, then 30 cycles at 94°C 30 sec, 50°C 15 sec, 72°C 15 sec, with final elongation at 72°C for 10 min. SSCP was performed following the method of Yap *et al.* (6): Briefly, 2–10 µl of PCR product was adjusted to a total volume of 10 µl with water, incubated at 42°C for 10 min after addition of 1 µl 0.5 M NaOH, 10 mM EDTA. 1 µl of formamide containing 0.5% bromophenol blue and 0.5% xylene cyanol was added just before loading. Electrophoresis was carried out on 5–6% polyacrylamide gels either with 10% glycerol at 20°C or without glycerol at 4°C. The gels were run at 400–500 volts (20–25 V/cm) for 4 hours. Bands were visualized by silver staining. Double-stranded PCR products purified by agarose gel electrophoresis were sequenced using Sequenase version 2.0 (USB) under the conditions described by Casanova *et al.* (7).

Every affected individual in NF1 family 15 showed one aberrant single strand band and one aberrant double strand band in exon 28 by SSCP analysis. Direct sequencing of the PCR-product revealed an 'AG' deletion at codon 1703 (5108delAG). This deletion leads to a shift of the reading frame, creates a stop codon 93 bases downstream, and thus should result in a truncated gene product of 1733 amino acids (see Figure 1). One aberrant

single strand band segregating with the disease phenotype in NF1 family 72 was detected in exon 31 by SSCP analysis. Sequence analysis revealed a guanine insertion at codon 1939 (5816insG), which also leads to a frameshift resulting in a stop codon 7 bases downstream, predicting a truncated protein of 1941 amino acids (see Figure 2). This mutation abolishes a CviR I restriction site.

The adult patients 49 and 51 of NF1 family 15 suffer from a very severe dermal form of NF1 with extensive involvement of trunk, neck and face. Individual 47 of the same family developed multiple dermal neurofibromas before puberty, and has left-sided tibia pseudoarthrosis and one plexiform neurofibroma causing extreme hypertrophy of the right foot. All patients present with mild to severe funnel chest. Common clinical features shared by the patients of NF1 family 72 include multiple *café-au-lait* spots, and dermal and subcutaneous neurofibromas. In addition, patient 332 and her affected brother, who has not been investigated by molecular studies, suffer from severe scoliosis. In patients 328 and 326, no spine deformity has been found.

Only a few mutations have been previously identified in the NF1 gene. Of the mutations presented here, 5108delAG is associated with very severe dermal manifestations and funnel chest in NF1 family 15, and 5816insG with severe scoliosis in 2 out of 4 affected members from NF1 family 72. Interestingly, three other previously reported families carrying nonsense mutations in exon 31 also exhibited scoliosis (8, 9), suggesting a possible correlation between this clinical feature and these particular mutations. It will be important to identify a larger number of mutations in NF1 patients in order to clarify the phenotypic consequences of mutations in specific gene regions. The identification of these mutations is also helpful for genetic counselling and diagnosis of affected individuals and asymptomatic gene carriers in the respective families.

## ACKNOWLEDGEMENTS

We would like to thank Dr Albert Schinzel and Dr Wendy Robinson for helpful comments on this manuscript.

## REFERENCES

1. Viskochil, D., Buchberg, A.M., Xu, G., Cawthon, R., Stevens, J., Wolff, R.K., Culver, M., Carey, J.C., Copeland, N.G., Jenkins, N.A., White, R. and O'Connell, P. (1990) *Cell* 62, 187–192.
2. Cawthon, R.M., Weiss, R., Xu, G., Viskochil, D., Culver, M., Stevens, J., Robertson, M., Dunn, D., Gesteland, R., O'Connell, P., and White, R. (1990) *Cell* 62, 193–201.

\* To whom correspondence should be addressed




Figure 1. a) SSCP analysis of exon 28 in NF1 family 15 and controls. The arrow indicates the aberrant single-stranded (ss) DNA band, the arrow head shows the aberrant double-stranded (ds) DNA band. b) Partial sequence of exon 28 in patient 51 and control. The position of the 'AG' deletion is indicated by an arrow. The samples were sequenced from the 3' direction. c) Nucleotide and amino acid sequence information showing the normal (N) and mutated (A) sequence. The deletion of nucleotides 5108 and 5109 leads to a shift of the reading frame resulting in 30 altered amino acids and a stop codon at amino acid 1734.




Figure 2. a) SSCP analysis of exon 31 in NF1 family 72 and controls. The aberrant single-stranded (ss) DNA band is indicated by an arrow. b) Partial sequence of exon 31 showing a guanine insertion in patient 328. The normal sequence is shown at the right. The samples were sequenced from the 5' direction. c) Comparison of the normal (N) and mutated (A) sequence shows that the guanine insertion at nucleotide 5816 leads to alterations of three amino acids and creates a stop codon at amino acid 1942.

- Wallace, M.R., Marchuk, D.A., Andersen, L.B., Letcher, R., Odeh, H.M., Saulino, A.M., Fountain, J.W., Brereton, A., Nicholson, J., Mitchell, A.L., Brownstein, B.H., and Collins, F.S. (1990) *Science* 249, 181-186.
- Marchuk, D.A., Saulino, A.M., Tavakoli, R., Swaroop, M., Wallace, M.R., Andersen, L.B., Mitchell, A.L., Gutmann, D.H., Boguski, M. and Collins, F.S. (1991) *Genomics* 11, 931-940.
- Upadhyaya, M., Shen, M., Cherryson, A., Farnham, J., Maynard, J., Huson, S.M. and Harper, P.S. (1992) *Hum. Mol. Genet.* 1, 735-740.
- Yap, E.P.H. and McGee, J. (1992) *Trends Genet.* 8, 49.
- Casanova, J.-L., Pannetier, C., Jaulin, C. and Kourilsky, P. (1990) *Nucleic Acids Res.* 18, 4028.
- Alnsworth, P.J., Rodenhiser, D.L. and Costa, M.T. (1993) *Hum. Genet.* 91, 151-156.
- Estivill, X., Lazzaro, C., Casals, T. and Ravello, A. (1991) *Hum. Genet.* 88, 185-188.