

## The geometric realization of Wall obstructions by nilpotent and simple spaces

By GUIDO MISLIN

*ETH-Z, 8092 Zürich*

(Received 12 March 1979)

*Introduction.* Let  $\pi$  denote a finite group. It is well known that every element of the projective class group  $K_0\mathbb{Z}\pi$  may be realized as Wall obstruction of a finitely dominated complex with fundamental group  $\pi$  (cf. (13)). We will study two subgroups  $N_0\mathbb{Z}\pi$  and  $N\mathbb{Z}\pi$  of  $K_0\mathbb{Z}\pi$ , which are closely related to the Wall obstruction of nilpotent spaces. If the group  $\pi$  is nilpotent and if  $S$  denotes the set of elements  $x \in K_0\mathbb{Z}\pi$  which occur as Wall obstructions of nilpotent spaces, then

$$N_0\mathbb{Z}\pi \subset S \subset N\mathbb{Z}\pi.$$

It turns out that in many instances one has  $N_0\mathbb{Z}\pi = N\mathbb{Z}\pi$  (cf. Section 3) and one obtains hence new information on  $S$ . The main theorem (2.4) provides a systematic way of constructing finitely dominated nilpotent (or even simple) spaces with non-vanishing Wall obstructions.

1. *The groups  $T\mathbb{Z}\pi$  and  $N\mathbb{Z}\pi$ .* If  $\pi$  denotes a finite group then one defines  $T\mathbb{Z}\pi \subset K_0\mathbb{Z}\pi$  to be the subgroup consisting of all elements of the form  $[(k, N)] - [\mathbb{Z}\pi]$ , where  $N = \Sigma x$ ,  $x \in \pi$ , and  $(k, N)$  is the projective ideal in  $\mathbb{Z}\pi$  generated by  $N$  and an integer  $k$  prime to  $\text{card}(\pi)$ . The group  $T\mathbb{Z}\pi$  is known to be trivial if  $\pi$  is cyclic (9). On the other hand  $T\mathbb{Z}\pi \neq 0$  if  $\pi$  contains a noncyclic subgroup of odd order (11).  $T\mathbb{Z}\pi$  is completely known for  $\pi$  a  $p$ -group (10).

It is convenient to think of  $K_0\mathbb{Z}\pi$  to be generated by  $\pi$ -modules  $M$  of type  $FP$  and to write  $[M]$  for the element  $\Sigma(-1)^i [P_i] \in K_0\mathbb{Z}\pi$ , if

$$0 \rightarrow P_n \rightarrow P_{n-1} \rightarrow \dots \rightarrow P_0 \rightarrow M \rightarrow 0$$

is a projective resolution of finite type. For instance, if  $k$  is prime to  $\text{card}(\pi)$  one has an exact sequence  $0 \rightarrow \mathbb{Z}\pi \rightarrow (k, N) \rightarrow \mathbb{Z}/k \rightarrow 0$  and hence

$$[(k, N)] - [\mathbb{Z}\pi] = [\mathbb{Z}/k] \in K_0\mathbb{Z}\pi,$$

where  $\mathbb{Z}/k$  is considered as a trivial  $\pi$ -module (cf. (4)). If  $\pi \neq \{1\}$  then every trivial  $\pi$ -module of type  $FP$  is necessarily finite and of order prime to  $\text{card}(\pi)$ . We can then identify  $T\mathbb{Z}\pi$  with the subgroup of  $K_0\mathbb{Z}\pi$  consisting of all elements representable in the form  $[M]$ , where  $M$  is a trivial  $\pi$ -module of type  $FP$ .

In view of the applications we have in mind, we will define more general subgroups  $N_i\mathbb{Z}\pi \subset K_0\mathbb{Z}\pi$  in a similar way.

*Definition 1.1.* Let  $\pi$  be a finite non-trivial group. Then  $N_i\mathbb{Z}\pi \subset K_0\mathbb{Z}\pi$  is the sub-group consisting of all elements of the form  $\Sigma(-1)^k [P_k]$ , where  $P = \{P_k\}$  is a

projective complex of finite type whose homology groups  $H_k(P)$  are all nilpotent  $\pi$ -modules and for which  $H_j(P) = 0$  for  $j > i$ . Furthermore,  $N\mathbb{Z}\pi = \cup N_i\mathbb{Z}\pi$  and for  $\pi = \{1\}$  we define  $N\mathbb{Z}\pi = N_i\mathbb{Z}\pi = 0$  for all  $i$ .

To see that  $N_i\mathbb{Z}\pi$  is indeed a subgroup for  $\pi \neq \{1\}$ , it suffices to check that all elements of  $N_i\mathbb{Z}\pi$  are of finite order ( $N_i\mathbb{Z}\pi$  is obviously closed under addition). But this amounts to showing that  $\Sigma(-1)^k \text{rank}(P_k) = 0$  for  $P = \{P_k\}$  as in 1.1. But this follows immediately from the isomorphism

$$H_k(P) \otimes \mathbb{Q} \xrightarrow{\cong} H_k(P) \otimes_{\pi} \mathbb{Q}$$

which holds, since  $H_k(P)$  is a nilpotent  $\pi$ -module, and from the equalities

$$\Sigma(-1)^k \text{rank}(P_k) = \Sigma(-1)^k \dim_{\mathbb{Q}}(H_k(P) \otimes_{\pi} \mathbb{Q}) = \frac{1}{|\pi|} \Sigma(-1)^k \dim_{\mathbb{Q}}(H_k(P) \otimes \mathbb{Q}).$$

By definition  $N_0\mathbb{Z}\pi$  consists of elements  $x = [M]$ , where  $M$  is a nilpotent  $\pi$ -module of type  $FP$ . If  $P$  is as in 1.1 and  $H_j(P) = 0$  for  $j > 0$  then  $H_0(P) = M$  is a finite module, since

$$\Sigma(-1)^k \text{rank}(P_k) = \dim_{\mathbb{Q}}(M \otimes_{\pi} \mathbb{Q}) = 0.$$

Clearly, this  $M$  is also cohomologically trivial, since it is of type  $FP$ .

**COROLLARY 1.2.** *The subgroup  $N_0\mathbb{Z}\pi \subset K_0\mathbb{Z}\pi$  consists of all elements  $x = [M] \in K_0\mathbb{Z}\pi$ , where  $M$  is a finite, nilpotent, and cohomologically trivial  $\pi$ -module.*

This is clear from the above, since a finite  $M$  which is cohomologically trivial is of type  $FP$  (and even of projective dimension  $\leq 1$  by (8)).

In particular we see that  $T\mathbb{Z}\pi \subset N_0\mathbb{Z}\pi \subset N\mathbb{Z}\pi$ . The following example will illustrate that in general however  $T\mathbb{Z}\pi \neq N_0\mathbb{Z}\pi$ .

**LEMMA 1.3.** *If  $\pi$  is cyclic of order 15, then  $N_0\mathbb{Z}\pi$  is of order two.*

*Proof.* Choose a map  $\pi \rightarrow \text{Aut}(\mathbb{Z}/9)$  which maps on to the subgroup of order 3. This defines a nilpotent  $\pi$ -module  $M$  with underlying abelian group  $\mathbb{Z}/9$ .  $M$  is a nilpotent, cohomologically trivial  $\pi$ -module and  $M$  generates  $\tilde{K}_0\mathbb{Z}\pi \cong \mathbb{Z}/2$  (cf. (5), Lemma 2.8). Hence  $N_0\mathbb{Z}\pi$  is of order two.

*Remark.* Let  $D\mathbb{Z}\pi$  denote the kernel of the map  $K_0\mathbb{Z}\pi \rightarrow K_0\overline{\mathbb{Z}\pi}$ , induced by including  $\mathbb{Z}\pi$  into a maximal  $\mathbb{Z}$ -order  $\overline{\mathbb{Z}\pi}$  in  $\mathbb{Q}\pi$ . If  $\pi$  is nilpotent, then

$$N\mathbb{Z}\pi \subset D\mathbb{Z}\pi.$$

This is proved in (12) for  $\pi$  cyclic and in (7) for a general nilpotent  $\pi$ . An example is given in (7) to show that in general  $N\mathbb{Z}\pi \neq D\mathbb{Z}\pi$ , even if  $\pi$  is cyclic.

**2. The realization theorem.** All spaces we consider are supposed to be pointed connected  $CW$ -complexes;  $\tilde{X}$  denotes the universal covering space of a space  $X$ . As usual a homology class is called spherical if it lies in the image of the Hurewicz homomorphism. First, we will describe a particular way of killing certain spherical classes.

**LEMMA 2.1.** *Let  $X$  be an  $n$ -dimensional  $CW$ -complex and let  $P \subset H_n \tilde{X}$  denote a projective  $\pi_1 X$ -module consisting of spherical classes. Denote by  $\phi: L \rightarrow H_n \tilde{X}$  a map from a*

free  $\pi_1 X$ -module  $L$  with basis  $\{b_\alpha, \alpha \in I\}$ , such that  $\phi(L) = P$ . Then one can form a new complex

$$X' = X \cup \left( \coprod_{\alpha \in I} e_\alpha^{n+1} \right)$$

of dimension  $n+1$  such that

(1) there is a commutative diagram

$$\begin{array}{ccc} \pi_{n+1}(\tilde{X}', \tilde{X}) & \xrightarrow{\partial} & \pi_n \tilde{X} \xrightarrow{Hu} H_n \tilde{X} \\ \phi_L \uparrow \cong & & \nearrow \phi \\ L & & \end{array}$$

(2)

$$H_i \tilde{X}' \cong \begin{cases} H_i \tilde{X} & \text{if } i \neq n, n+1 \\ (H_i \tilde{X})/P & \text{if } i = n \\ \text{Ker } \phi & \text{if } i = n+1. \end{cases}$$

(3)  $H_{n+1} \tilde{X}'$  is projective and spherical (i.e. it consists entirely of spherical classes).

*Proof.* Since  $P$  is projective, one can find  $\bar{P} \subset \pi_n \tilde{X}$  such that  $\bar{P}$  is mapped isomorphically onto  $P$  by the Hurewicz homomorphism. Hence we can choose  $\bar{\phi}: L \rightarrow \bar{P}$  to obtain a commutative diagram

$$\begin{array}{ccc} & & \bar{P} \subset \pi_n \tilde{X} \\ & \nearrow \bar{\phi} & \\ L & & \downarrow h \quad \downarrow Hu \\ & \searrow \phi & P \subset H_n \tilde{X} \end{array}$$

Denote by  $\text{pr}: \tilde{X} \rightarrow X$  the projection. We attach  $(n+1)$ -cells to  $X$  using the maps  $\text{pr}(\bar{\phi} b_\alpha)$ ,  $\alpha \in I$ , and obtain  $X' = X \cup (\coprod e_\alpha^{n+1})$ . It is immediate that  $\phi$  lifts to an isomorphism  $\phi_L$  giving rise to diagram (1). We consider now the diagram obtained by mapping the homotopy sequence of  $(\tilde{X}', \tilde{X})$  into the homology sequence of this pair:

$$\begin{array}{ccccccc} \pi_{n+1} \tilde{X}' & \xrightarrow{\alpha} & \pi_{n+1}(\tilde{X}', \tilde{X}) & \xrightarrow{\partial_n} & \pi_n \tilde{X} & & \\ \downarrow & & \downarrow \cong & & \downarrow & & \\ H_{n+1} \tilde{X}' & \xrightarrow{\beta} & H_{n+1}(\tilde{X}', \tilde{X}) & \xrightarrow{\partial_H} & H_n \tilde{X} & \twoheadrightarrow & H_n \tilde{X}' \end{array}$$

Then  $\text{im } \partial_n = \bar{P}$  and hence  $\text{im } \partial_H = P$ . Therefore  $H_n \tilde{X}' \cong (H_n \tilde{X})/P$ . Note also that  $\alpha(\pi_{n+1} \tilde{X}')$  is mapped isomorphically onto  $\beta(H_{n+1} \tilde{X}') \cong H_{n+1} \tilde{X}'$ , since  $\bar{P}$  is mapped isomorphically onto  $P$ . Hence  $H_{n+1} \tilde{X}' \cong \text{Ker } \partial_n \cong \text{Ker } \phi$  and  $\pi_{n+1} \tilde{X}' \rightarrow H_{n+1} \tilde{X}'$  is onto. Therefore (2) and (3) hold.

**THEOREM 2.2.** Let  $X$  be a connected CW-complex of dimension  $n > 1$  and let  $M$  be a  $\pi_1 X$ -module of cohomological dimension  $\leq 1$ . Then there is a space  $Y$  obtained from  $X$  by attaching cells of dimension  $\geq n$ , such that

$$H_i \tilde{Y} = \begin{cases} H_i \tilde{X} & \text{if } i \neq n \\ (H_n \tilde{X}) \oplus M & \text{if } i = n. \end{cases}$$

Furthermore, if  $X$  is finitely dominated and  $M$  of type  $FP$ , then  $Y$  can be chosen to be finitely dominated; the reduced Wall obstructions of  $X$  and  $Y$  are then related by

$$\tilde{w} Y = \tilde{w} X + (-1)^n [M] \in \tilde{K}_0 \mathbb{Z} \pi.$$

*Proof.* Choose a free resolution

$$\cdots \rightarrow L_{n+i} \xrightarrow{\phi_{n+i}} L_{n+i-1} \rightarrow \cdots \rightarrow L_{n+1} \xrightarrow{\phi_{n+1}} L_n \twoheadrightarrow M.$$

(If  $M$  is a type  $FP$ , we choose a free resolution of finite type.) Since  $\text{proj. dim } M \leq 1$ ,  $\text{im } (\phi_{n+i})$  is projective for all  $i \geq 1$ . We construct  $Y$  inductively as follows. Let  $Y^n = X \vee B$ , where  $B$  is a bouquet of  $n$ -spheres corresponding to a basis of  $L_n$ . Then  $H_n \tilde{Y}^n \cong (H_n \tilde{X}) \oplus L_n$  and  $H_n \tilde{Y}^n$  contains a spherical projective submodule  $P$  isomorphic to  $\text{im } (\phi_{n+1})$ . Attaching  $(n+1)$ -cells to  $Y^n$  with respect to the map  $L_{n+1} \rightarrow H_n \tilde{Y}^n$  corresponding to  $\phi_{n+1}$ , we obtain by the previous Lemma a new space  $Y^{n+1}$  with

$$H_i \tilde{Y}^{n+1} = \begin{cases} H_i \tilde{X} & \text{if } i \neq n, n+1 \\ (H_n \tilde{X}) \oplus M & \text{if } i = n \\ \text{Ker } \phi_{n+1} & \text{if } i = n+1. \end{cases}$$

Since  $\text{Ker } \phi_{n+1} \cong H_{n+1} \tilde{Y}^{n+1}$  is projective and spherical (Lemma 2.1) we can kill this group using  $L_{n+2} \rightarrow H_{n+1} \tilde{Y}^{n+1}$ . By repeating this construction we obtain spaces  $Y^{n+k}$ ,  $k \geq 1$ , and we can form  $Y = \cup Y^{n+k}$ . By construction,  $\tilde{Y}$  has the homology groups claimed in the theorem. Furthermore, the cellular chain complex of  $\tilde{Y}$  is isomorphic to the complex

$$\cdots \rightarrow L_{n+1} \rightarrow L_{n+i-1} \rightarrow \cdots \rightarrow L_n \oplus C_n \tilde{X} \rightarrow C_{n-1} \tilde{X} \rightarrow \cdots,$$

where  $C \tilde{X}$  is the cellular chain complex of  $\tilde{X}$ . Since the complex

$$\cdots \rightarrow L_{n+i} \rightarrow L_{n+i-1} \rightarrow \cdots \rightarrow L_{n+1} \rightarrow \text{im } \phi_{n+1}$$

is contractible, it follows that  $Y$  is a retract of  $Y^{n+1}$ . Hence  $Y$  is finitely dominated, if  $X$  is finitely dominated and  $M$  of type  $FP$ . From the definition of the Wall obstruction it is immediate that

$$\begin{aligned} \tilde{w}(Y) &= \sum_{i=0}^n (-1)^i [\bar{C}_i X] + (-1)^{n+1} [\text{im } \phi_{n+1}] \\ &= \tilde{w}(X) + (-1)^n [M], \end{aligned}$$

where  $\bar{C} \tilde{X}$  is a chain complex of type  $FP$ , chain homotopy equivalent to  $C \tilde{X}$ .

Before we apply this Theorem to the construction of certain nilpotent spaces, we need the following elementary lemma.

**LEMMA 2.3.** *Let  $\pi$  be a finite group. Then there exists a finite complex  $X$  with  $\pi_1 X \cong \pi$  and Euler characteristic  $\chi(X) = 0$ , such that all covering transformations  $t: \tilde{X} \rightarrow \tilde{X}$  are homotopic to the identity.*

*Proof.* Choose an embedding  $\pi \subset SU(k)$ . Then  $X = SU(k)/\pi$  has the desired properties.

Note that  $X = SU(k)/\pi$  is nilpotent, if  $\pi$  is a nilpotent group, and it is a simple space, in case  $\pi$  is abelian.

We can now prove our main theorem.

**THEOREM 2.4.** *Let  $\pi$  be a finite nilpotent group and let  $x \in N_0\mathbb{Z}\pi \subset K_0\mathbb{Z}\pi$ . Then there exists a finitely dominated nilpotent space  $Y$  with fundamental group  $\pi$  and Wall obstruction  $w(Y) = x$ . If  $x$  lies in  $T\mathbb{Z}\pi$  and  $\pi$  is abelian, then  $Y$  may be chosen simple.*

*Proof.* Let  $x = [M]$  with  $M$  a nilpotent  $\pi$ -module (trivial  $\pi$ -action in case  $x \in T\mathbb{Z}\pi$ ). Choose  $X = SU(k)/\pi$  as in the previous lemma (we may assume that  $\dim X$  is even). Then, according to Theorem 2.2 we can construct a finitely dominated space  $Y$  with  $\pi_1 Y \cong \pi_1 X$  and

$$H_i \tilde{Y} = \begin{cases} H_i \tilde{X} & \text{if } i \neq \dim X \\ (H_i \tilde{X}) \oplus M & \text{if } i = \dim X. \end{cases}$$

It follows that  $wY = [M] = x$ , since  $X$  is finite with Euler characteristic 0. Moreover,  $Y$  is nilpotent since its fundamental group is nilpotent and since  $H_i \tilde{Y}$  is nilpotent for all  $i$ . In order to see that  $Y$  is simple in case  $\pi$  is abelian and  $x \in T\mathbb{Z}\pi$ , we prove the stronger result stating that, if  $M$  is a trivial  $\pi$ -module, then all covering transformations  $t: \tilde{Y} \rightarrow \tilde{Y}$  are homotopic to the identity. By the 'Hasse-Principle' for free maps (3) it suffices to show that the localizations  $t_p: \tilde{Y}_p \rightarrow \tilde{Y}_p$  are homotopic to the identity for all primes  $p$ . If  $p$  does not divide the order of  $M$ , then the inclusion  $X \subset Y$  induces  $H_i \tilde{X}_p \cong H_i \tilde{Y}_p$  and hence  $X_p \simeq Y_p$  (the induced map of fundamental groups is certainly an isomorphism). Therefore  $t_p \simeq \text{Id } \tilde{Y}_p$  since the corresponding result is true for  $X$  by construction. If  $p$  divides the order of the trivial  $\pi$ -module  $M$ , then  $p$  is necessarily prime to the order of  $\pi$ , since otherwise  $M$  would not be cohomologically trivial. It follows therefore that the projection  $\tilde{Y} \rightarrow Y$  induces a homotopy equivalence  $\tilde{Y}_p \simeq Y_p$  if  $p$  divides the order of  $M$ ; clearly this implies that  $t_p \simeq \text{Id } \tilde{Y}_p$  and hence the global map  $t$  is homotopic to the identity (3).

*Remark.* If in Theorem 2.4 the assumption that  $\pi$  be nilpotent is dropped, one can still construct the finitely dominated space  $Y$  with  $w(Y) = x \in N_0\mathbb{Z}\pi$ . The space  $Y$  will however in general only be *homologically nilpotent* in the sense that  $\pi_1 Y$  operates nilpotently on  $H_i \tilde{Y}$  for all  $i$ .

Theorem 2.4 enables us to construct examples of the following types:

**COROLLARY 2.5.** (a) *There exists a finitely dominated simple space with non-vanishing Wall obstruction.*

(b) *There exists a finitely dominated nilpotent space with cyclic fundamental group and non-vanishing Wall obstruction.*

*Proof.* For (a) choose any abelian group  $\pi$  with  $T\mathbb{Z}\pi \neq 0$  (e.g.  $(\mathbb{Z}/p) \times (\mathbb{Z}/p) \times (\mathbb{Z}/p)$ ,  $p$  any prime) and apply Theorem 2.4. Similarly, for (b) we can choose any cyclic group  $\pi$  with  $N_0\mathbb{Z}\pi \neq 0$  (e.g.  $\mathbb{Z}/15$ , cf. Lemma 1.3) and we obtain such an example by Theorem 2.4.

3. *Some computations involving  $N\mathbb{Z}\pi$ .* One can consider  $N_0\mathbb{Z}\pi$  as a lower bound for the elements in  $K_0\mathbb{Z}\pi$  which occur as Wall obstructions of finitely dominated homologically nilpotent space. Similarly,  $N\mathbb{Z}\pi$  provides an upper bound for this set. The

following examples show that for many groups one has  $N_0\mathbb{Z}\pi = N\mathbb{Z}\pi$  and, indeed, we don't know of an example with  $N_0\mathbb{Z}\pi \neq N\mathbb{Z}\pi$ .

Our main tool will be a homomorphism

$$T\colon U(\mathbb{Z}[1/n]) \rightarrow K_0\mathbb{Z}\pi/T\mathbb{Z}\pi$$

which was defined in (5) for groups  $\pi$  with cyclic Sylow subgroups ( $U(\mathbb{Z}[1/n])$  the units in  $\mathbb{Z}[1/n]$  and  $n = \text{card}(\pi)$ ). For the following computation we will assume that  $\pi$  is of square-free order  $n$  (hence  $T$  is defined). If  $N = \Sigma x, x \in \pi$ , the projection  $\mathbb{Z}\pi \rightarrow \mathbb{Z}\pi/N$  induces an injective map

$$\overline{pr}_*\colon K_0\mathbb{Z}\pi/T\mathbb{Z}\pi \hookrightarrow K_0(\mathbb{Z}\pi/N).$$

It is convenient to describe  $T$  by considering  $K_0(\mathbb{Z}\pi/N)$  as the range of  $T$ . If  $p$  is a prime dividing  $n = \text{card}(\pi)$  then the trivial  $\pi$ -module  $\mathbb{Z}/p$  considered as a  $\mathbb{Z}\pi/N$ -module, is of type  $FP$  with respect to the ring  $\mathbb{Z}\pi/N$  and

$$\overline{pr}_*T(p) = [\mathbb{Z}/p] \in K_0(\mathbb{Z}\pi/N).$$

Furthermore,  $T(-1) = 0$  (for details see (5)).

The connexion with  $N\mathbb{Z}\pi$  is given by the next lemma.

**LEMMA 3.1.** *Let  $\pi$  denote a group of square-free order  $n$  and let  $x \in N\mathbb{Z}\pi$ . Let  $P = \{P_i\}$  be a projective  $\pi$ -complex with  $x = \Sigma (-1)^i [P_i]$  and  $H_iP$  nilpotent for all  $i$ . Then*

$$\rho(P) = \text{card } H_{\text{ev}}(\mathbb{Z}\pi/N \otimes_\pi P) / \text{card } H_{\text{odd}}(\mathbb{Z}\pi/N \otimes_\pi P)$$

is a unit in  $\mathbb{Z}[1/n]$  and if  $\bar{x}$  denotes the image of  $x$  in  $N\mathbb{Z}\pi/T\mathbb{Z}\pi$  then

$$\bar{x} = T\rho(P) \in K_0\mathbb{Z}\pi/T\mathbb{Z}\pi.$$

In particular one has  $N\mathbb{Z}\pi/T\mathbb{Z}\pi \subset \text{im}(T)$ .

*Proof.* This result was proved in ((5), Section 3) in case  $x = wX$ , the Wall obstruction of a homologically nilpotent space  $X$ . The same proof works for an arbitrary  $x \in N\mathbb{Z}\pi$ .

**COROLLARY 3.2.** *Let  $\pi$  be of order  $p$  or  $2p$ ,  $p$  an arbitrary prime. Then  $N\mathbb{Z}\pi = 0$ .*

*Proof.* If  $\text{card}(\pi) = p$  with  $p$  an arbitrary prime or if  $\text{card}(\pi) = 2p$ ,  $p$  an odd prime, then  $\text{im}(T) = 0$  by ((5), Theorem 2.5). Hence  $N\mathbb{Z}\pi = T\mathbb{Z}\pi$  in these cases. But in both cases one has  $T\mathbb{Z}\pi = 0$  (cf. (11)). It remains to consider the case  $\text{card}(\pi) = 4$ . But it is well known that  $\tilde{K}_0\mathbb{Z}\pi = 0$  if  $\text{card}(\pi) = 4$ . Hence the result follows.

**THEOREM 3.3.** *If  $\pi$  is a cyclic group of square-free order, then*

$$N_0\mathbb{Z}\pi = N\mathbb{Z}\pi = \text{im}(T).$$

*Proof.* Note that  $T\mathbb{Z}\pi = 0$  for  $\pi$  cyclic. Hence  $N\mathbb{Z}\pi \subset \text{im}(T)$  by Lemma 3.1 and it suffices therefore to show that  $\text{im}(T) \subset N_0\mathbb{Z}\pi$ . If  $p$  is a prime dividing  $\text{card}(\pi)$  then  $\overline{pr}_*T(p) = [\mathbb{Z}/p] \in K_0(\mathbb{Z}\pi/N)$ . It remains to prove that there exist  $x \in N_0\mathbb{Z}\pi$  with  $\overline{pr}_*x = [\mathbb{Z}/p]$ . If  $x = [M]$ ,  $M$  a nilpotent  $\pi$ -module of projective dimension  $\leq 1$ , and if  $0 \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$  is a resolution of type  $FP$ , then by definition

$$pr_*x = [\mathbb{Z}\pi/N \otimes_\pi P_0] - [\mathbb{Z}\pi/N \otimes_\pi P_1].$$

But since  $M$  is cohomologically trivial, one has

$$\mathrm{Tor}_1^{\pi}(\mathbb{Z}\pi/N, M) \cong \mathrm{Ker}(M/IM \xrightarrow{N} M) = \hat{H}^{-1}(\pi, M) = 0$$

and therefore the sequence

$$0 \rightarrow \mathbb{Z}\pi/N \otimes_{\pi} P_1 \rightarrow \mathbb{Z}\pi/N \otimes_{\pi} P_0 \rightarrow \mathbb{Z}\pi/N \otimes_{\pi} M \rightarrow 0$$

is exact. Hence we can write

$$\overline{pr}_*[M] = [\mathbb{Z}\pi/N \otimes_{\pi} M].$$

We will now construct for all prime divisors of  $\mathrm{card}(\pi)$  nilpotent  $\pi$ -modules of type  $FP$  with  $\mathbb{Z}\pi/N \otimes_{\pi} M \cong \mathbb{Z}/p$ . First consider the case of an odd prime  $p$ . Let  $\mathbb{Z}/p$  act in a non-trivial way on  $\mathbb{Z}/p^2$  and define a  $\pi$ -action on  $\mathbb{Z}/p^2$  using a surjection  $\pi \rightarrow \mathbb{Z}/p$ . One verifies easily that the resulting  $\pi$ -module  $M$  is nilpotent and cohomologically trivial. Furthermore,  $\overline{pr}_*[M] = [\mathbb{Z}\pi/N \otimes_{\pi} M] = [M/NM] = [\mathbb{Z}/p]$ . If  $p = 2$ , one can use  $\mathbb{Z}/8$  as underlying abelian group for  $M$ , which one equips with a  $\mathbb{Z}/2$ -action by mapping  $a$  into  $5a$ ,  $a \in \mathbb{Z}/8$ , and defining a  $\pi$ -module structure by means of a surjection  $\pi \rightarrow \mathbb{Z}/2$ . Again one verifies that  $\overline{pr}_*[M] = [\mathbb{Z}/2]$ . Hence  $\mathrm{im}(T) \subset N_0\mathbb{Z}\pi$  and the result follows.

For the groups of Theorem 3.3 we can obtain an upper bound for the order and the exponent of  $N\mathbb{Z}\pi$  in terms of the Euler  $\phi$ -function  $\phi(n) = \mathrm{card}(U(\mathbb{Z}/n))$  and the function  $e(n) = (\text{exponent of } U(\mathbb{Z}/n))$ .

**THEOREM 3.4.** *If  $\pi$  is a cyclic group of square-free order  $n$ , then the order of*

$$N_0\mathbb{Z}\pi = N\mathbb{Z}\pi$$

*divides  $\phi(n)/e(n)$  and its exponent divides  $e(n)$ .*

*Proof.* Let  $p$  be a prime which divides  $n$  and let  $\bar{\pi} \subset \pi$  be a subgroup of index  $p$ . The  $[\mathbb{Z}/p] \in T\mathbb{Z}\bar{\pi} \subset K_0\mathbb{Z}\bar{\pi}$  is mapped to  $[\mathbb{Z}\pi \otimes_{\pi} \mathbb{Z}/p] = [\mathbb{Z}/p[\pi/\bar{\pi}]] \in K_0\mathbb{Z}\pi$  by the map induced by  $\bar{\pi} \subset \pi$  (one uses that  $\mathrm{Tor}_{\frac{1}{p}}^{\pi}(\mathbb{Z}\pi, \mathbb{Z}/p) = 0$ ). But if  $M = \mathbb{Z}/p[\pi/\bar{\pi}]$  then  $\mathbb{Z}\pi/N \otimes_{\pi} M = M/NM$  is a nilpotent  $\pi$ -module of cardinality  $p^{p-1}$  and hence  $\overline{pr}_*[M] = (p-1)\overline{pr}_*T(p) \in K_0(\mathbb{Z}\pi/N)$ . Since  $T\mathbb{Z}\bar{\pi} = 0$  ( $\bar{\pi}$  is cyclic),  $[M] = 0$  and hence  $(p-1)T(p) = 0$ . We obtain therefore a factorization

$$\begin{array}{ccc} U(\mathbb{Z}[1/n]) & \xrightarrow{T} & K_0\mathbb{Z}\pi \\ \lambda \searrow & & \nearrow \bar{T} \\ & U(\mathbb{Z}/n) & \end{array}$$

where  $\lambda: U(\mathbb{Z}[1/n]) \rightarrow U(\mathbb{Z}/n) \cong \Pi(\mathbb{Z}/p-1)$  is defined by  $\lambda(-1) = 0$  and  $\lambda(p) = (0, \dots, 0, \bar{1}, 0, \dots, 0)$  if  $p$  divides  $n$ . The diagonal element  $\Delta = (\bar{1}, \dots, \bar{1})$  in  $U(\mathbb{Z}/n) \cong \Pi(\mathbb{Z}/p-1)$  is mapped to 0 by  $\bar{T}$ , since  $T(n) = 0$  (cf. Theorem 2.5 of (5)). Moreover,  $N_0\mathbb{Z}\pi = N\mathbb{Z}\pi = \mathrm{im}(T)$  by Theorem 3.3. Hence the exponent of  $N\mathbb{Z}\pi$  divides the exponent of  $U(\mathbb{Z}/n)$  and the order of  $N\mathbb{Z}\pi$  divides  $\phi(n)/e(n)$  which is the order of  $U(\mathbb{Z}/n)/\langle \Delta \rangle$ .

For example, if  $\pi$  is a cyclic group of order  $3p$ ,  $p$  a prime  $> 3$ , then  $\mathrm{card}(N\mathbb{Z}\pi) \leq 2$  since  $\phi(3p) = 2(p-1)$  and  $e(2p) = p-1$ .



As a final example, we want to compute  $N\mathbb{Z}\pi$  in case  $\pi = M(p, q)$ , the metacyclic group of square-free order  $pq$ ,  $p$  and  $q$  odd primes and  $q|p-1$ , defined by

$$M(p, q) = \langle x, y | x^p = y^q = 1, y^{-1}xy = x^r \rangle,$$

$r$  a primitive  $q$ th root of 1 mod  $p$ .

**THEOREM 3.5.** *Let  $\pi = M(p, q)$ . Then*

$$T\mathbb{Z}\pi = N_0\mathbb{Z}\pi = N\mathbb{Z}\pi \cong \mathbb{Z}/q.$$

*Proof.* It has been shown in (6) that if  $x \in K_0\mathbb{Z}\pi$  is the Wall obstruction of a homologically nilpotent space  $X$  with fundamental group  $M(p, q)$ , then  $x \in T\mathbb{Z}\pi$ . The same argument shows that for an arbitrary  $x \in N\mathbb{Z}\pi$  one has  $x \in T\mathbb{Z}\pi$  and hence  $N\mathbb{Z}\pi = T\mathbb{Z}\pi$ . Furthermore,  $T\mathbb{Z}\pi = \mathbb{Z}/q$  by (11).

One can combine the results of this section to obtain the following table for  $N\mathbb{Z}\pi$ , in case  $\pi$  is a group of small, square-free order.

**COROLLARY 3.6.** *Let  $\pi$  be a group of square-free order  $n < 30$ . Then*

$$N_0\mathbb{Z}\pi = N\mathbb{Z}\pi = \begin{cases} 0 & \text{if } n \neq 15, 21 \\ \mathbb{Z}/2 & \text{if } n = 15 \\ \mathbb{Z}/3 & \text{if } n = 21, \pi \text{ noncyclic} \\ \mathbb{Z}/2 \text{ or } 0 & \text{if } n = 21, \pi \text{ cyclic.} \end{cases}$$

## REFERENCES

- (1) FRÖHLICH, A. On the class group of integral group rings of finite abelian groups. I. *Mathematika* **16** (1969), 143–152.
- (2) FRÖHLICH, A., KEATING, M. E. and WILSON, S. M. J. The class groups of quaternion and dihedral 2-groups. *Mathematika* **21** (1974), 64–71.
- (3) HILTON, P., MISLIN, G., ROITBERG, J. and STEINER, R. On free maps and free homotopies into nilpotent spaces. *Springer Lecture Notes in Math. Vol. 673*, 1977.
- (4) MISLIN, G. Finitely dominated nilpotent spaces. *Ann. of Math.* **103** (1976), 547–556.
- (5) MISLIN, G. Groups with cyclic Sylow subgroups and finiteness conditions for certain complexes. *Comment. Math. Helv.* **52** (1977), 373–391.
- (6) MISLIN, G. Finitely dominated complexes with metacyclic fundamental groups. *Topology and Algebra, Proceedings of a Colloquium in honour of B. Eckmann*, Monographie No. 26, *L'Enseignement Mathématique* 1978, 233–235.
- (7) MISLIN, G. and VARADARAJAN, K. The finiteness obstruction for nilpotent spaces lie in  $D(\mathbb{Z}\pi)$ . *Inventiones math.* **53** (1979), 185–191.
- (8) RIM, D. S. Modules over finite groups. *Ann. of Math.* **63** (1959), 700–712.
- (9) SWAN, R. G. Periodic resolutions for finite groups. *Ann. of Math.* **72** (1960), 267–291.
- (10) TAYLOR, M. J. Locally free class groups of prime power order. *J. Algebra* **50** (1978), 463–487.
- (11) ULLOM, S. V. Nontrivial lower bounds for class groups of integral group rings. *Illinois J. of Math.* **20** (1976), 361–371.
- (12) VARADARAJAN, K. Finiteness obstructions for nilpotent spaces. *J. Pure and Appl. Algebra* **12** (1978), 137–146.
- (13) WALL, C. T. C. Finiteness conditions for CW-complexes. *Ann. of Math.* **81** (1965), 56–69.