

- 21 Chen L, Bracey AW, Radovancevic R, et al. Clopidogrel and bleeding in patients undergoing elective coronary artery bypass grafting. *J Thorac Cardiovasc Surg* 2004; **128**: 425–31
- 22 Ranucci M, Baryshnikova E, Soro G, Ballotta A, De Benedetti D, Conti D. Multiple electrode whole-blood aggregometry and bleeding in cardiac surgery patients receiving thienopyridines. *Ann Thorac Surg* 2011; **91**: 123–9
- 23 Vilahur G, Choi BG, Zafar MU, et al. Normalization of platelet reactivity in clopidogrel-treated subjects. *J Thromb Haemost* 2007; **5**: 82–90
- 24 Zafar MU, Santos-Gallego C, Vorchheimer DA, et al. Platelet function normalization after a prasugrel loading-dose: time-dependent effect of platelet supplementation. *J Thromb Haemost* 2013; **11**: 100–6
- 25 Storey RF, Bliden KP, Ecob R, et al. Earlier recovery of platelet function after discontinuation of treatment with ticagrelor compared with clopidogrel in patients with high antiplatelet responses. *J Thromb Haemost* 2011; **9**: 1730–7
- 26 Prüller F, Drexler C, Archan S, Macher S, Raggam RB, Mahla E. Low platelet reactivity is recovered by transfusion of stored platelets: a healthy volunteer in vivo study. *J Thromb Haemost* 2011; **9**: 1670–3
- 27 Douketis JD, Spyropoulos AC, Spencer FA, et al. Perioperative management of antithrombotic therapy: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest* 2012; **141**: e326S–50S
- 28 Gogarten W, Vandermeulen E, Van Aken H, et al. Regional anaesthesia and antithrombotic agents: recommendations of the European Society of Anaesthesiology. *Eur J Anaesthesiol* 2010; **27**: 999–1015
- 29 Horlocker TT. Regional anaesthesia in the patient receiving antithrombotic and antiplatelet therapy. *Br J Anaesth* 2011; **107**(Suppl. 1): i96–106
- 30 Jakubowski JA, Payne CD, Weerakkody GJ, et al. Dose-dependent inhibition of human platelet aggregation by prasugrel and its interaction with aspirin in healthy subjects. *J Cardiovasc Pharmacol* 2007; **49**: 167–73
- 31 Jackson SP. The growing complexity of platelet aggregation. *Blood* 2007; **109**: 5087–95
- 32 Hosokawa K, Ohnishi T, Fukasawa M, et al. A microchip flow-chamber system for quantitative assessment of the platelet thrombus formation process. *Microvasc Res* 2012; **83**: 154–61
- 33 Hosokawa K, Ohnishi T, Sameshima H, et al. Analysing responses to aspirin and clopidogrel by measuring platelet thrombus formation under arterial flow conditions. *Thromb Haemost* 2013; **109**: 102–11
- 34 Bevers EM, Comfurius P, Zwaal RF. Platelet procoagulant activity: physiological significance and mechanisms of exposure. *Blood Rev* 1991; **5**: 146–54
- 35 Kozek-Langenecker SA, Afshari A, Albaladejo P, et al. Management of severe perioperative bleeding: guidelines from the European Society of Anaesthesiology. *Eur J Anaesthesiol* 2013; **30**: 270–382
- 36 Gorlinger K, Shore-Lesserson L, Dirkmann D, Hanke AA, Rahe-Meyer N, Tanaka KA. Management of hemorrhage in cardiothoracic surgery. *J Cardiothorac Vasc Anesth* 2013; **27**: S20–34
- 37 Skolnick BE, Shenouda M, Khutoryansky NM, Pusateri AE, Gabriel D, Carr ME. Reversal of clopidogrel-induced bleeding with rFVIIa in healthy subjects: a randomized, placebo-controlled, double-blind, exploratory study. *Anesth Analg* 2011; **113**: 703–10
- 38 van der Linden J, Lindvall G, Sartipy U. Aprotinin decreases post-operative bleeding and number of transfusions in patients on clopidogrel undergoing coronary artery bypass graft surgery: a double-blind, placebo-controlled, randomized clinical trial. *Circulation* 2005; **112**: I276–80
- 39 Thiele T, Sumnig A, Hron G, et al. Platelet transfusion for reversal of dual antiplatelet therapy in patients requiring urgent surgery: a pilot study. *J Thromb Haemost* 2012; **10**: 968–71

British Journal of Anaesthesia **112** (5): 784–7 (2014)

Advance Access publication 27 February 2014 · doi:10.1093/bja/aeu004

EDITORIAL III

Fibrinogen concentrate: clinical reality and cautious Cochrane recommendation

S. Kozek-Langenecker^{1*}, D. Fries², D. R. Spahn³ and K. Zacharowski⁴

¹ Department of Anaesthesia and Intensive Care, Evangelical Hospital Vienna, Austria

² Department of General and Surgical Critical Care Medicine, Medical University Innsbruck, Austria

³ Institute of Anaesthesiology, University and University Hospital Zurich, Switzerland

⁴ Department of Anesthesiology, Intensive Care Medicine and Pain Therapy, University Hospital Frankfurt, Germany

* Corresponding author. E-mail: sibylle.kozek@aon.at

A recent systematic review, 'Fibrinogen concentrate in bleeding patients', published by The Cochrane Collaboration, aimed to assess the benefits and harms of fibrinogen concentrate compared with placebo or other treatments in bleeding patients.¹ We agree with the authors finding that fibrinogen concentrate reduces transfusion requirements for allogeneic blood products,

as this is in line with previously published work.^{2,3} In addition, the review did not identify any adverse events associated with fibrinogen concentrate; again, this is in line with previous findings.^{4,5} However, we feel that the review also made a number of incorrect or misleading assertions which merit further discussion and clarification.

Firstly, the authors suggest that the use of fibrinogen concentrate should be confined 'to a controlled clinical setting or trial', a suggestion that has been made previously by Stanworth and Hunt.⁶ However, if this principle were to be applied routinely, it would lead to the abolition of all blood bank products, including plasma and cryoprecipitate, until robust trial data became available.⁷ This suggestion also ignores the fact that the most widely available fibrinogen concentrate has held a licence since 1963 in Brazil, since 1966 in Europe, has been licensed in its current pasteurized form since 1985, and is currently licensed for treatment of both congenital and acquired fibrinogen deficiencies in Argentina, Austria, Brazil, Bulgaria, the Czech Republic, Germany, Hungary, Iran, Kuwait, the Netherlands, Portugal, Romania, Switzerland, Taiwan, Tunisia, and Turkey.⁸ Indeed, a number of recent guidelines recommend the use of fibrinogen concentrate in the bleeding patient, including the updated European trauma guidelines⁹ and the European Society of Anaesthesiology guidelines;¹⁰ both of these guidelines were published before the searches for the Cochrane analysis were completed. It should also be noted that fibrinogen concentrate has a licence for the treatment of congenital deficiencies in Australia, Canada, Israel, Mexico, New Zealand, Puerto Rico, the USA, and numerous EU countries. Any off-label use in these countries (e.g. in the setting of acquired bleeding) is at the discretion of the individual physician, taking into account the risk profile of the available alternative treatments.¹¹

The authors of the Cochrane review also recommend not using fibrinogen concentrate in the clinical setting until further evidence is available. However, they give no suitable recommendation of an alternative treatment. This negligence could in fact endanger patients, as the alternatives for fibrinogen supplementation, that is, fresh-frozen plasma (FFP) and cryoprecipitate, are associated with a number of adverse events.¹² Furthermore, the efficacy and safety of cryoprecipitate have never been tested in the setting of a randomized controlled trial (RCT). A recent systematic review of FFP found no consistent evidence or significant benefit for its prophylactic or therapeutic use across a range of settings in which it is indicated;¹³ FFP is also associated with increased morbidity.¹⁴ Cryoprecipitate has been withdrawn from a number of European countries in response to safety concerns,¹⁵ and in most countries in which it is still available, it is no longer used for hereditary bleeding disorders. In fact, as acknowledged in the Cochrane review, fibrinogen concentrate is now preferred to both cryoprecipitate and FFP in the hereditary setting, and so the continued use of these products for the treatment of acquired bleeding disorders represents a double standard.¹⁶ In 2009, the Irish Blood Transfusion Service took the decision to withdraw cryoprecipitate and replace it with fibrinogen concentrate, despite a lack of formal licensing.¹⁷ This decision was made to reduce the risk of pathogen transmission, and was supported by a lack of evidence suggesting any benefit of cryoprecipitate over fibrinogen concentrate.¹⁷ In addition, the World Health Organization recently urged member states to 'promote the availability of transfusion alternatives'.¹⁸ It is perhaps

unethical for the Cochrane group to suggest that a newer treatment with a growing body of evidence should be shunned in favour of older treatments which, despite being in use since World War II, have never been proven or evaluated according to the Cochrane methodology and are associated with a number of safety concerns. Tradition should not factor in the decision-making process in clinical practice, and, in any case, one might argue that fibrinogen concentrate is rapidly becoming the 'traditional' (standard of care) treatment for acquired hypofibrinogenaemia in a number of countries, such as Austria, Germany, and Switzerland.¹⁹

A number of RCTs are currently taking place or are planned to investigate the use of fibrinogen concentrate in settings such as cardiac surgery and postpartum haemorrhage. Of course, the evidence gathered from such trials can and should be used to inform clinical practice in the future.²⁰ However, there are already numerous publications reporting the regular use of fibrinogen concentrate, particularly in centres across Europe.^{17 21 22} There has also been a focus on the use of algorithms to guide individualized dosing of fibrinogen, often driven by point-of-care testing.^{21 22} As fibrinogen is the first coagulation factor to decrease to critically low levels during haemorrhage,²³ the use of fibrinogen concentrate allows targeted replacement of a specific part of the coagulation cascade. Relying on blanket therapies such as plasma or cryoprecipitate, with all their associated risks, is taking a step backwards when we should be moving forward into an era of targeted, personalized medicine. Indeed, the Research Ethics Committee of the Canton Zurich recently refused permission for the centre to participate in the REPLACE trial (a prospective, randomized, placebo-controlled study investigating fibrinogen concentrate to control bleeding during cardiac surgery), as they considered it unethical to expose patients in the control arm to unnecessary FFP transfusions when patients outside the context of the study were receiving fibrinogen concentrate according to the institutional algorithm-based coagulation management guidelines, by which their exposure to blood products is reduced.

Another point that warrants clarification is the authors' description of infusion times. The authors suggest that mixing of fibrinogen concentrate takes 15 min; in fact, the manufacturer suggests this takes only 5–10 min. The authors also underestimate the amount of time it takes to prepare FFP and cryoprecipitate; these products take time to be ordered, tested for blood group compatibility (where necessary), and prepared for administration. Not every hospital has access to equipment to allow them to thaw frozen products in <15 min, as suggested by the authors of the review, and thawing takes 45 min at room temperature.²⁴ Taking all these factors into account, the entire blood product preparation process can take 60–90 min.^{24 25} Pre-thawed or liquid (never frozen) plasmas can be used; however, these products are of limited availability, and are generally only accessible in a small number of large, tertiary care hospitals. The authors of this review are correct in their assertion that there is no recommended infusion rate for FFP; however, this is simply because there is a lack of clinical evidence from which an optimal infusion rate can be determined. The manufacturers of the

commercial plasma Octaplas recommend a rate of no more than $1 \text{ ml kg}^{-1} \text{ min}^{-1}$;²⁶ infusion at rates greater than this has been shown to result in citrate toxicity and hypocalcaemia.²⁶ Using the suggested rate, for a patient of 70 kg, and assuming a fibrinogen concentration for FFP of 2 g litre^{-1} ,²⁸ it would take more than 35 min (and over 2 litre of FFP) to transfuse a 5 g fibrinogen dose. For fibrinogen concentrate, an optimal infusion rate of 5 ml min^{-1} is quoted by the manufacturers, as determined from the results of pharmacokinetics studies and RCTs. In a controlled setting, infusion of both fibrinogen concentrate and blood products should take place as slowly as possible; however, in an emergency situation, rapid infusion may be desirable. Studies in cardiovascular surgery have shown that it is possible to infuse 1 g fibrinogen concentrate in $<20 \text{ s}$ (i.e. 250 ml min^{-1}).^{5 29}

Finally, despite the fact that none of the studies analysed in the Cochrane review addressed the cost–benefit issue, the authors of the Cochrane review chose to discuss the current cost of fibrinogen concentrate in one country, Denmark. However, the authors chose to quote the retail price of 657€, which includes both the margins added by the pharmacy and the VAT; it would be more appropriate to quote the pharmacy purchase price, which in Denmark is currently 479€.³⁰ The authors also failed to consider that costs will vary widely from region to region and are dependent upon a number of factors, such as the tax and healthcare systems of each individual country. Therefore, picking a price in just one European country to use as an example is of little benefit to the international readership of the Cochrane review. In addition, the authors neglected to discuss the equivalent costs of the alternatives, that is, FFP and cryoprecipitate. Without such a comparison, any stated costs are irrelevant and uninformative. In fact, determining the cost of allogeneics is a highly complex issue in need of detailed analysis, as the true cost of blood products is far higher than the direct acquisition cost of the product itself; indirect costs such as storage, preparation, thawing, processing, and compatibility testing also have to be taken into account.³¹ A previous discussion regarding the price of allogeneics in Sweden prompted the suggestion that health economics would benefit from the input of both clinicians and economists, as neither are in a position to fully understand all of the facts.³²

In conclusion, we feel that the Cochrane review contains information that may mislead readers. Systematic reviews such as this should provide clear, unambiguous recommendations for clinicians to follow, and these recommendations should be clearly justified, particularly when they relate to a licensed product with a broad label that is now the standard of care in several countries. If alternative treatments are suggested, this should be in the knowledge that they are a safe and efficacious option for patients, rather than simply relying on ‘tradition’ and potentially outdated treatments that have themselves never been subjected to the appropriate research scrutiny.

Declaration of interest

S.K.-L. has received payments and travel funding from Baxter, Biotest, CSL Behring, Novo Nordisk, Octapharma, and TEM

International. D.F. has received study funding, payments, and travel funding from Austrian National Bank, AOP Orphan, Astra Zeneca, Baxter, B.Braun, Biotest, CSL Behring, Fresenius Kabi, Glaxo, Haemoscope, Hemogem, Lilly, LFB, Mitsubishi Pharma, Novo Nordisk, Octapharma, and TEM International. D.R.S. has received study funding from CSL Behring, Vifor SA, Villars-sur-Glane, and payments and travel funding from Abbott, Baar, Amgen, AstraZeneca, Baxter, B. Braun, Boehringer Ingelheim, Bristol-Myers-Squibb, CSL Behring, Curacyte, Ethicon Biosurgery, Fresenius, Galenica, GlaxoSmithKline, Janssen-Cilag, Beers, Merck Sharp & Dohme, Novo Nordisk, Octapharma, Oxygen Biotherapeutics, TEM International, ratiopharm, Roche Pharma, Schering-Plough, and Vifor Pharma. K.Z. has received payments and travel funding from CSL Behring.

References

- 1 Wikkelso A, Lunde J, Johansen M, et al. Fibrinogen concentrate in bleeding patients. *Cochrane Database Syst Rev* 2013; **8**: CD008864
- 2 Rahe-Meyer N, Solomon C, Hanke A, et al. Effects of fibrinogen concentrate as first-line therapy during major aortic replacement surgery: a randomized, placebo-controlled trial. *Anesthesiology* 2013; **118**: 40–50
- 3 Fenger-Eriksen C, Jensen TM, Kristensen BS, et al. Fibrinogen substitution improves whole blood clot firmness after dilution with hydroxyethyl starch in bleeding patients undergoing radical cystectomy: a randomized, placebo-controlled clinical trial. *J Thromb Haemost* 2009; **7**: 795–802
- 4 Dickneite G, Pragst I, Joch C, Bergman GE. Animal model and clinical evidence indicating low thrombogenic potential of fibrinogen concentrate (Haemocomplettan P). *Blood Coagul Fibrinolysis* 2009; **20**: 535–40
- 5 Solomon C, Hagl C, Rahe-Meyer N. Time course of haemostatic effects of fibrinogen concentrate administration in aortic surgery. *Br J Anaesth* 2013; **110**: 947–56
- 6 Stanworth SJ, Hunt BJ. The desperate need for good-quality clinical trials to evaluate the optimal source and dose of fibrinogen in managing bleeding. *Crit Care* 2011; **15**: 1006
- 7 Kozek-Langenecker S, Sorensen B, Hess J, Spahn DR. Emotional or evidence-based medicine—is there a moral tragedy in haemostatic therapy? *Crit Care* 2011; **15**: 462
- 8 Levy JH, Welsby I, Goodnough LT. Fibrinogen as a therapeutic target for bleeding: a review of critical levels and replacement therapy. *Transfusion* Advance Access published on Oct 9, 2013, doi: 10.1111/trf.12431
- 9 Spahn DR, Bouillon B, Cerny V, et al. Management of bleeding and coagulopathy following major trauma: an updated European guideline. *Crit Care* 2013; **17**: R76
- 10 Kozek-Langenecker SA, Afshari A, Albaladejo P, et al. Management of severe perioperative bleeding: guidelines from the European Society of Anaesthesiology. *Eur J Anaesthesiol* 2013; **30**: 270–382
- 11 Gollop ND, Chilcott J, Benton A, et al. National audit of the use of fibrinogen concentrate to correct hypofibrinogenaemia. *Transfus Med* 2012; **22**: 350–5
- 12 Murad MH, Stubbs JR, Gandhi MJ, et al. The effect of plasma transfusion on morbidity and mortality: a systematic review and meta-analysis. *Transfusion* 2010; **50**: 1370–83
- 13 Yang L, Stanworth S, Hopewell S, Doree C, Murphy M. Is fresh-frozen plasma clinically effective? An update of a systematic review of randomized controlled trials. *Transfusion* 2012; **52**: 1673–86; quiz

- 14 Inaba K, Branco BC, Rhee P, et al. Impact of plasma transfusion in trauma patients who do not require massive transfusion. *J Am Coll Surg* 2010; **210**: 957–65
- 15 Sorensen B, Bevan D. A critical evaluation of cryoprecipitate for replacement of fibrinogen. *Br J Haematol* 2010; **149**: 834–43
- 16 Bevan DH. Cardiac bypass haemostasis: putting blood through the mill. *Br J Haematol* 1999; **104**: 208–19
- 17 Ahmed S, Harrity C, Johnson S, et al. The efficacy of fibrinogen concentrate compared with cryoprecipitate in major obstetric haemorrhage—an observational study. *Transfus Med* 2012; **22**: 344–9
- 18 World Health Organization (WHO). 63rd World Health Assembly. Availability, safety and quality of blood products. 2010. Available from http://apps.who.int/gb/ebwha/pdf_files/WHA63/A63_R12-en.pdf (accessed 22 October 2013)
- 19 Ranucci M, Solomon C. Supplementation of fibrinogen in acquired bleeding disorders: experience, evidence, guidelines, and licences. *Br J Anaesth* 2012; **109**: 135–7
- 20 Ranucci M. Fibrinogen supplementation in cardiac surgery: where are we now and where are we going? *J Cardiothorac Vasc Anesth* 2013; **27**: 1–4
- 21 Weber CF, Gorlinger K, Meininger D, et al. Point-of-care testing: a prospective, randomized clinical trial of efficacy in coagulopathic cardiac surgery patients. *Anesthesiology* 2012; **117**: 531–47
- 22 Schochl H, Nienaber U, Hofer G, et al. Goal-directed coagulation management of major trauma patients using thromboelastometry (ROTEM)-guided administration of fibrinogen concentrate and prothrombin complex concentrate. *Crit Care* 2010; **14**: R55
- 23 Hiippala ST, Myllylä GJ, Vahtera EM. Hemostatic factors and replacement of major blood loss with plasma-poor red cell concentrates. *Anesth Analg* 1995; **81**: 360–5
- 24 Goodnough LT, Spain DA, Maggio P. Logistics of transfusion support for patients with massive hemorrhage. *Curr Opin Anaesthesiol* 2013; **26**: 208–14
- 25 Goodnough LT, Shander A. How I treat warfarin-associated coagulopathy in patients with intracerebral hemorrhage. *Blood* 2011; **117**: 6091–9
- 26 Octapharma. Octaplas summary of product characteristics. Available from http://www.octapharma.co.uk/index.php?eID=tx_nawsecuredl&u=0&file=uploads/media/octaplasLG_SmPC_01.pdf&t=1393419280&hash=ddc2b6fbb6e096a0a733896dc09c0f220a7ee262 (accessed 25 February 2014)
- 27 Cote CJ, Drop LJ, Hoaglin DC, Daniels AL, Young ET. Ionized hypocalcemia after fresh frozen plasma administration to thermally injured children: effects of infusion rate, duration, and treatment with calcium chloride. *Anesth Analg* 1988; **67**: 152–60
- 28 Theusinger OM, Baulig W, Seifert B, et al. Relative concentrations of haemostatic factors and cytokines in solvent/detergent-treated and fresh-frozen plasma. *Br J Anaesth* 2011; **106**: 505–11
- 29 Solomon C, Pichlmaier U, Schoechl H, et al. Recovery of fibrinogen after administration of fibrinogen concentrate to patients with severe bleeding after cardiopulmonary bypass surgery. *Br J Anaesth* 2010; **104**: 555–62
- 30 Danish Medicine Agency. Medicine prices in Denmark. Available from <http://www.medicinpriser.dk/> (accessed 29 October 2013)
- 31 Shander A, Hofmann A, Ozawa S, et al. Activity-based costs of blood transfusions in surgical patients at four hospitals. *Transfusion* 2010; **50**: 753–65
- 32 Glennard AH, Persson U, Soderman C. Costs associated with blood transfusions in Sweden—the societal cost of autologous, allogeneic and perioperative RBC transfusion. *Transfus Med* 2005; **15**: 295–306

British Journal of Anaesthesia **112** (5): 787–90 (2014)

Advance Access publication 24 February 2014 · doi:10.1093/bja/aeu006

EDITORIAL IV

Surrogate measures, do they really describe anaesthetic state?

F. S. Servin^{1*} and V. Billard²

¹ Service d'anesthésie, APHP-Hôpital Bichat, 46, rue Henri Huchard, 75018 Paris, France

² Service d'anesthésie, Institut Gustave Roussy, 114, rue Édouard-Vaillant, 94805 Villejuif Cedex, France

* Corresponding author. E-mail: frederique.servin@bch.aphp.fr

Characterization of anaesthesia is tricky, in particular, to go beyond the usual definition of a 'state of non-responsiveness to various types of stimulations', and to find indicators useful for titrating drug administration. General anaesthesia is usually split into two main domains: unconsciousness (non-response to verbal stimulation and amnesia) and analgesia (non-response to noxious stimulations).^{1 2} Unconsciousness may be directly assessed at induction of anaesthesia (loss of verbal contact) and after recovery (absence of recall), but usually not during maintenance. The adequacy of the

balance between analgesia and stimulation can be clinically estimated through movement, haemodynamic changes, and autonomic nervous system responses (sweat and lacrimation, modification in pupil diameter); these clinical endpoints have limitations when used for titrating anaesthetic drugs in clinical practice, specifically in paralysed patients. Furthermore, as adequate anaesthesia is defined as non-responsiveness, clinical assessment cannot distinguish between adequate drug delivery and overdosing if overdosing does not induce adverse effects.