

A structural study of copper(II) carboxylates: Crystal structure and physical characterisation of $[\text{Cu}_2(2\text{-bromopropanoato})_4(\text{caffeine})_2]$

Fedor Valach,^{(1)*} Milan Melník,⁽²⁾ Gérald Bernardinelli,⁽³⁾ and Katharina M. Fromm⁽⁴⁾

Received August 30, 2005; accepted April 25, 2006
Published Online June 27, 2006

Based on the bond-valence sum model the continual transition between the bidentate and monodentate bonding mode of carboxylato group and minimum of Cu–O bond lengths were estimated. The dependence of Cu···Cu separation on Cu–N bond length in binuclear Cu(II) carboxylates with caffeine were derived and fitted to the observed data. The crystal structure of $[\text{Cu}_2(2\text{-bromopropanoato})_4(\text{caffeine})_2]$ has been determined by X-ray crystallography. The structure consists of centrosymmetric binuclear units where the two Cu atoms are coordinated by four disordered bridging 2-bromopropionates and two caffeine ligands at the apices of a bicapped square prism. Both 2-bromopropionates show disorder of their –Br and –CH₃ substituents. Stacking $\pi\cdots\pi$ interactions between the adjacent caffeine molecules link the complex units in 1-D networks. The binuclear structure of the studied compound is consistent with magnetic data and EPR spectrum.

KEY WORDS: Copper(II) carboxylates; halogenopropionates; caffeine; crystal structure; π - π stacking interaction.

Introduction

Copper(II) carboxylates are interesting from both points of views, the central atom and the ligand properties. This class of compounds may contain mononuclear, binuclear and polynuclear structural units.¹ The coordination sphere ge-

ometry of Cu(II) complexes, in contrast to the carboxylato group reveals a certain flexibility.² Correlations between interatomic distances, bond lengths or angles of binuclear Cu(II) complexes have been reported.^{1a,b,3} The bond-valence sum (BVS) model⁴ provides a framework for the explanation of such structural correlations implying the mechanisms that produce them.

Carboxylate groups may coordinate to a Cu(II) atom through one O atom or through both O atoms. The last bonding mode may occur as chelate ligand bonded to one metal center or as bridging group connecting two metal centers. To our knowledge, until now, several crystal structures of Cu(II) halogenated propionates were solved. $[\text{Cu}(\text{CF}_3\text{CF}_2\text{COO})_2(\text{NITPhCl})_2]$ (NIT-PhCl = 2-(4-chloro-phenyl)-4,4,5,5-tetramethyl-

⁽¹⁾ Slovak University of Technology, Faculty of Chemical and Food Technology, Institute of Physical Chemistry and Chemical Physics, Radlinského 9, SK-812 37 Bratislava, Slovak Republic.

⁽²⁾ Slovak University of Technology, Faculty of Chemical and Food Technology, Institute of Inorganic Chemistry, Technology and Materials, Radlinského 9, SK-812 37 Bratislava, Slovak Republic.

⁽³⁾ Laboratory of X-ray Crystallography, University of Geneva, 24, quai E. Anserment, CH-1211 Genève 4, Switzerland.

⁽⁴⁾ Department of Chemistry, University of Basel, Spitalstrasse 51, CH-4056 Basel, Switzerland.

* To whom correspondence should be addressed; e-mail: fedor.valach@stuba.sk

imidazoline-1-oxyl-3-oxide)⁵ (**1**) is mononuclear whereas $[\text{Cu}_2(\text{CF}_3\text{CF}_2\text{COO})_4(\text{NITmNO}_2)_2]$ ($\text{NITmNO}_2 = 2-(3\text{-nitrophenyl})-4,4,5,5\text{-tetramethylimidazoline-1-oxyl-3-oxide}$)⁶ (**2**) is a binuclear complex. The structures of $[\text{Cu}_2(\text{ClCH}_2\text{CH}_2\text{COO})_4(\text{Ph}_3\text{PO})_2]$ ($\text{Ph}_3\text{PO} = \text{triphenylphosphine oxide}$) (**3**),⁷ $[\text{Cu}_2(\text{CH}_3\text{CHClCOO})_4(\text{Ph}_3\text{PO})_2]$ ⁸ (**4**) and $[\text{Cu}_2(\text{ClCH}_2\text{CHClCOO})_4(4\text{-pic})_2]$ ($4\text{-pic} = 4\text{-picoline}$)⁹ (**5**) consist of binuclear complex units. $[\text{Cu}_3(\text{CH}_3\text{CHClCOO})_3(\text{Et}_2\text{LO})_2(\text{OH})(\text{H}_2\text{O})]_2$ ($\text{Et}_2\text{LO} = 2\text{-diethylaminoethanolate}$)¹⁰ (**6**) complex is hexanuclear. Recently, the structures of mononuclear complexes $[\text{Cu}(\text{CH}_3\text{CCl}_2\text{COO})_2(3\text{-mpyc})_2]$ ($3\text{-mpyc} = \text{methyl-3-pyridyl-carbamate}$) at room temperature¹¹ (**7**) and at 100 K¹² (**8**) and $[\text{Cu}_2(\text{CH}_3\text{CHBrCOO})_4(2\text{-pyme})_2]$ ($2\text{-pyme} = 2\text{-pyridylmethanol}$) at 208 K¹² (**9**) were reported. The structure of $[\text{Cu}(\text{CH}_3\text{CCl}_2\text{COO})_2(\text{dena})_2(\text{H}_2\text{O})_2 \cdot \text{Cu}(\text{CH}_3\text{CCl}_2\text{COO})_2(\text{dena})_2(\text{H}_2\text{O})]$ ($\text{dena} = \text{N,N-diethylnicotinamide}$)¹³ (**10**) consists of two chemically different units. The derivatives of propionic acid possess two quadribonded asymmetric carbon atoms¹⁴ in 2-(α -) and 3-(β -) positions. But only in the compounds **2**, **7** and **8** the positional disorder of propionate ligands was reported. $[\text{Cu}_2(\text{CH}_3\text{CHBrCOO})_4(\text{caffeine})_2]$ is the first example showing the chirality of halogenopropanoate carbon atom. A few crystal structures of copper(II) compounds with caffeine (3,7-dihydro-1,3,7-trimethyl-1,4-purine-2,6-dione) were resolved^{3b,15} in which the copper(II) atoms are five-coordinated. Eight of them^{3b,15a-g} comprise the complex units of the composition $[\text{Cu}_2\text{X}_4(\text{caffeine})_2]$ ($\text{X} = \text{carboxylate}$) with binuclear paddle-wheel cage structure. Four X anions function as bridging groups in a *syn-syn* arrangement. Nitrogen atom of caffeine assumes the apical position of the deformed tetragonal pyramid around copper(II) atom.

As a part of our investigations of halogenopropionates we present in this paper the crystal and molecular structure of the entitled compound. For Cu(II) carboxylates and described binuclear copper(II) complexes with caffeine the correlations of metal-ligand and metal-metal distances interpreted using BVS model is reported.

Experimental

Preparation of complex

The $[\text{Cu}_2(2\text{-bromopropionato})_4(\text{caffeine})_2]$ was prepared as described earlier¹⁶ with the racemic 2-bromopropionate acid. The studied compound was prepared by adding a methanolic solution of caffeine to a stirred methanolic solution of $\text{Cu}(\text{CH}_3\text{CHBrCOO})_2$ in an equimolar ratio. After heating to reflux, the solution was left to cool and stand at room temperature. The green product that precipitated was isolated and washed with cold ethanol and dried at ambient temperature. The crude product was recrystallized from hot ethanol to yield green crystals.

Anal. Found: Cu, 11.3; C, 30.0; H 3.9; N, 9.95%.

Calc. for $\text{Cu}(\text{CH}_3\text{CHBrCOO})_2(\text{caffeine})$; Cu, 11.31; C, 29.94; H, 3.95 and N, 9.98%.

Crystal structure determination

Intensity data were collected using a Stoe Ipds diffractometer at 200 K. The diffraction intensities were corrected for Lorentz and polarisation effects. Absorption correction was applied. The calculations were made by the programs of X-RED software.¹⁷ The structure was solved by direct methods using the program MULTAN-87,¹⁸ and refined by the full-matrix least-squares method (on F) using the program CRYLSQ¹⁹ (273 parameters). Hydrogen atoms of the caffeine moiety have been refined and blocked in the last cycles. The other hydrogen atoms have been refined with restraints on bond lengths and bond angles and blocked in the last cycles. The structure was drawn by the program ORTEP-3 for Windows.²⁰ Crystal data and conditions of data collections are reported in Table 1. Selected geometrical parameters of the crystal structure are reported in Table 2.

Magnetic and spectral measurements

Electronic spectrum ($9000\text{--}50\,000\text{ cm}^{-1}$) of the powdered sample in a Nujol mull was recorded

Table 1. Summary of Crystal Data, Intensity Measurement and Structure Refinement for [Cu₂(2-bromopropionato)₄(caffeine)₂]

CCDC deposition number	254424
Empirical formula	C ₂₈ H ₃₆ Br ₄ Cu ₂ N ₈ O ₁₂
Molecular weight	1123.4
Crystal size (mm)	0.274 × 0.172 × 0.045
Crystal system	Triclinic
Space group	<i>P</i> -1
<i>a</i> (Å)	8.1483(10)
<i>b</i> (Å)	9.8723(11)
<i>c</i> (Å)	13.2438(13)
α (°)	108.605(12)
β (°)	96.690(13)
γ (°)	101.492(14)
<i>V</i> (Å ³)	970.7(3)
<i>Z</i>	1
<i>D_x</i> (Mg m ⁻³)	1.922
Wavelength (Å)	0.71073
μ (Mo <i>K</i> α) (mm ⁻¹)	5.28
<i>T</i> _{min} , <i>T</i> _{max}	0.3510, 0.7991
(sinΘ/λ) _{max} (Å ⁻¹)	0.612
No. of measured data	12065
No. of independent data	3529
No. of data in the refinements	1825
Weighting scheme <i>p</i> ^a	0.0002
Maximum Δ/ σ	0.002
Max. and min. Δρ (e.Å ⁻³)	0.89, -0.66
<i>S</i>	1.13
<i>R</i> , <i>wR</i>	0.038, 0.037

$$^a w = 1/(\sigma^2(F_o) + p(F_o)^2).$$

at room temperature on a Specord 200. Electron paramagnetic resonance (EPR) spectrum was recorded on the spectrometer Bruker ESP 300 operating at X-band at room temperature, and equipped with the Bruker NMR gaussmeter ER 035 M and Hewlett Packard microwave frequency counter HP5350B.

Magnetic measurements in the temperature range of 80–300 K were carried out on a sample of the complex at a magnetic field of 5 kG, using a Quantum Design SQUID Magnetometer (type MPMS-XL5). Corrections for diamagnetism of the constituting atoms were calculated using the Pascal constants,²¹ the value of 60.10⁻⁶ cm³ mol⁻¹ was used as the temperature-independent paramagnetism of a copper(II) atom. The effective magnetic moments were calculated from the expression

$$\mu_{\text{eff}} = 2.83\sqrt{\chi_M^{\text{corr}}T} \quad [\text{B.M.}] \quad (1)$$

Table 2. Selected Bond Distances (Å) and Angles and (°) for [Cu₂(2-bromopropionato)₄(caffeine)₂] with e.s.d.'s in Parentheses

Cu–Cu ⁱ	2.694(1)	Cu–O(2a)	1.953(6)
Cu–O(1a)	1.968(6)	Cu–O(1b)	1.976(5)
Cu–N(3)	2.231(6)	Cu–O(2b ⁱ)	1.967(4)
C(1a)–C(2a)	1.51(1)	C(1b)–C(2b)	1.511(9)
C(2a)–Br(1a)	1.929(7)	C(2b)–Br(1b)	1.911(9)
O(1b)–Cu–O(2a ⁱ)	89.4(3)	O(1b)–C(1b)–O(2b)	126.7(6)
O(1b)–Cu–O(2b ⁱ)	167.1(3)	O(1b)–C(1b)–C(2b)	116.5(7)
O(1a)–C(1a)–O(2a)	127.1(8)	O(2b)–C(1b)–C(2b)	116.7(7)
O(1a)–C(1a)–C(2a)	115.9(7)	C(1b)–C(2b)–Br(1b)	104.4(6)
O(2a)–C(1a)–C(2a)	117.0(7)	C(1b)–C(2b)–C(3b)	118(1)
Br(1a)–C(2a)–C(3a)	111(1)	Br(1b)–C(2b)–C(3b)	108(1)

Symmetry code: (i) 2 – *x*, 1 – *y*, 1 – *z*.

The EPR spectrum was measured using either a digitized Bruker ER 200 E SRC with an internal marked for determining the frequency.

Results and discussion

The crystal structure of the entitled compound consists of a centrosymmetric binuclear complex where the two Cu atoms are coordinated by eight oxygen atoms of four bridging 2-bromopropionates. Two nitrogen atoms of two caffeine ligands are located at the apex of a bicapped square prism. The structure of the binuclear complex is shown in Fig. 1. Each Cu(II) is five coordinated by four carboxylic O atoms in an equatorial plane (maximum deviation of the mean plane passing through the four oxygen atoms: 0.001(5) Å) and axially by one N atom from the molecule of caffeine, to give a tetragonal pyramid. The displacement of the Cu atom from the equatorial plane toward the apical atom N(3) [0.221(1) Å] is comparable with that found for other Cu(II) carboxylates.^{1a,22} The distance between the two basal planes of the studied compound as defined by Yablokov *et al.*²³ is 2.247(1) Å.

Both 2-bromopropionato ligands show a disorder by permutation of their –Br and –CH₃ substituents leading to an inversion of configuration at the C(2) position. The disorders were resolved by refining the population parameter of the major bromine substituent and fixing the sum with




Fig. 1. Ortep view of the crystal structure of $[\text{Cu}_2(2\text{-bromopropionato})_4(\text{caffeine})_2]$ showing the atom numbering of the asymmetric unit (black bonds) and the part of the binuclear complex generated by a centre of inversion (white bonds). Ellipsoids are represented with 50% probability level. Only the major disordered 2-bromopropionates are represented.

the minor to unity. These population parameters are also applied to the involved methyl C(3) and hydrogen atoms. The refinement of such a model (see Fig. 2) shows final disorder ratios of 0.907(4)/0.093(4) and 0.650(4)/0.350(4) for 2-bromopropionates **a**, and **b** respectively. The bond lengths and angles of 2-bromopropionate groups do not show significant deviations to those found in the structure of compound **9**. It should be noted that a refinement in the space group $P1$, led to similar distribution of the population parameters and a Flack parameter²⁵ $x = 0.48(7)$ confirming that the structure is centrosymmetric.

The caffeine ligand of $[\text{Cu}_2(2\text{-bromopropionato})_4(\text{caffeine})_2]$ complex is slightly deviated from planarity (maximum deviation = 0.052 Å for C(3)) and is involved in intermolecular $\pi \cdots \pi$ stacking interactions²⁶ through a centre of inversion (mean interplane distance = 3.56(3) Å) connecting the complex units into a infinite chain along *c* direction. The dihedral angle between the 6- and 5-membered rings of the caffeine moiety is 2.1(4)°. The methyl groups of caffeine molecule are slightly deviated from the planes of five- and six-membered rings. The torsion angles




Fig. 2. Perspective view of the crystal structure of $[\text{Cu}_2(2\text{-bromopropionato})_4(\text{caffeine})_2]$ (caffeine moieties are omitted for clarity) showing the disorder of the bromopropionates substituents. Bromine atoms are represented by spheres. The population parameters of the major (black) and minor (white) moieties are reported in parenthesis and the relative configuration at the C(2) centers²⁴ of the major part are mentioned.

C(8)–N(4)–C(1)–C(2) and C(7)–N(2)–C(4)–N(3) are 9(1)° and 2.4(9)°, respectively.

The $\text{Cu}_2(\text{COO})_4$ cage of studied compound is geometrically very close to the same cages of binuclear complexes **3–5**. The coordination of atom Cu for these complexes is square pyramidal. But the geometry around each atom Cu of the 2-bromopropionato complex **9** is distorted tetragonal bipyramid with coordination number 4 + 2. In Fig. 3 is shown the correlation between the Cu–O and O–C bond lengths of the independent Cu–O–C moieties. The data for binuclear copper complexes were collected by Cambridge Structural Database (CSD),²⁷ Version 2.3.8. The function (2) for non-linear regression of the correlation in Fig. 3 was derived using the logarithmic




Fig. 3. Dependence of between the Cu–O and C–O bond lengths (in Å) of the independent Cu–O–C moieties in the $\text{Cu}_2(\text{COO})_4$ cages with the scatterogram of compounds from CSD with crystallographic R -factors less than 0.06 (\circ). For chemically equivalent bonds of the compounds **3–5**, $[\text{Cu}_2(\text{CH}_3\text{CHBrCOO})_4(\text{caffeine})_2]$ and the compound **2**(*) $R = 0.17$ and $wR = 0.18$, the mean values are shown. The coefficient of linear correlation is -0.314 .

formula^{4a} for O–C bond valence and the formula based on orbital metal–ligand interaction of Cu–O bond:⁴ⁱ

$$r_{\text{O-C}} = R^* - b \ln \left(V^* - \frac{\alpha_1}{r_{\text{Cu-O}}} - \frac{\alpha_2}{r_{\text{Cu-O}}^2} - \frac{\alpha_3}{r_{\text{Cu-O}}^3} - \frac{\alpha_4}{r_{\text{Cu-O}}^4} - \frac{\alpha_5}{r_{\text{Cu-O}}^5} \right) \quad (2)$$

$r_{\text{O-C}}$ and $r_{\text{Cu-O}}$ are O–C and Cu–O bond lengths. Variable V^* is the valence of the atom O and R^* is the bond valence parameter^{4c} of the C–O bond and b is commonly accepted constant equal to 0.37 Å^4 . α_i are constants of the bond valence formula for the Cu–O bond. The fitted parameters are $R^* = 1.51(3) \text{ Å}$ and $V^* = 2.4(2)$. This correlation implies very weak influence of Cu–O bond on O–C bond lengths of carboxylato group.

The lengths of shorter and longer Cu–O bonds within the chelate tetrametallocycle

containing the chelated carboxylato group in this work are assigned: r_S and r_L respectively. Figure 4 shows the correlation r_S vs. r_L for hexacoordinated copper(II) carboxylates. The data were collected from Cambridge Structural Database (CSD),²⁷ Version 2.3.8. Only the structures solved from the single crystal data with crystallographic R -factor less than 0.05 were involved. The structures with at least one disordered atom were excluded. A continuous range of states occur between the monodentate and bidentate coordination of carboxylate group to one Cu atom which can be explained as the manifestation of coordination sphere plasticity.^{2a} The function (3) which express the correlation shown in Fig. 4 was derived from the BVS model. For the shorter Cu–O bond, the bond valence function based on the orbital metal–ligand interaction was used. The best fit was achieved using the formula for the bond valence of longer Cu–O bond $s_L = (R_o/r_L)^N$ in the range of $r_L = 2.6\text{--}3.07 \text{ Å}$. Within this range




Fig. 4. Dependence between r_L and r_S values (in Å) with the scatterogram from CSD data for hexacoordinated copper(II) carboxylates (open circles). Spearman rank-correlation coefficient²⁹ is $\rho = -0.336$. (■) **6**, (●) **9**, (▽) **10a**, (△) **10b**, (▼) **11**, (▲) **12**.

is the bond valence function s_L greater than bond valence function based on orbital interaction.⁴ⁱ R_o and N are constants.^{4a} The bond valence function s_L can be derived with a purely ionic approach.^{4e,g} On the other hand, there is a close correlation between the bond valence (bond strengths) s_L and the metal-oxygen covalence.^{4h}

$$r_L = R_o \left[\frac{\alpha_1}{r_S} + \frac{\alpha_2}{r_S^2} + \frac{\alpha_3}{r_S^3} + \frac{\alpha_4}{r_S^4} + \frac{\alpha_5}{r_S^5} + 2 - V_o^* \right]^{-1/N} \quad (3)$$

Variable V_o^* is the contribution of remaining copper-ligand bonds to the bond valence sum around the central atom.²⁸ The standard statistical procedure for the test of correlation significance and goodness of fit was used.²⁹ The fitted value of V_o^* is 1.493(2). The curve of function (3) approaches the minimum of short Cu–O bond length asymptotically at 1.90(5) Å (Fig. 4). Such interatomic distance occurs in complex **1** [1.903(2) Å], and consequently the pentafluoropropionate ligand of this complex is monodentate. In the range

where r_L is greater than 3.1 Å, r_S is almost insensitive towards further increase of the value of r_S . The carboxylato groups of the studied compounds with $r_L > 3.1$ Å are monodentate. The points of compound **9** reveal the greatest deviation from the fitted curve which can be explained by the relative strong hydrogen bonds of uncoordinated oxygen atoms to the methanol hydrogen atoms of 2-pyridylmethanol. The structure of compound **10** contains two kinds of independent complex units mutually bonded by the hydrogen bonds. The 2,2-chloropropionate anions of the unit $[\text{Cu}(\text{CH}_3\text{CCl}_2\text{COO})_2(\text{dena})_2(\text{H}_2\text{O})]$ (**10a**) are monodentate. Cu(II) atom of $[\text{Cu}(\text{CH}_3\text{CCl}_2\text{COO})_2(\text{dena})_2(\text{H}_2\text{O})_2]$ (**10b**) is coordinated by the monodentate and bidentate 2,2-chloropropionate anions. Similarly as in complex **9** the uncoordinated oxygen atoms of monodentate 2,2-chloropropionate anion are bonded *via* hydrogen bonds to the water molecule of the inner copper(II) coordination sphere. The pentafluoro- and 2-chloropropionate anions of complex **6** are monodentate with very weak Cu···O interactions.




Fig. 5. The dependences of Cu...Cu separation (full line) and the deviations of the copper(II) cation from the basal O₄ plane in CuO₄N chromophore (dashed line) on Cu-N bond length with the scatter plots of the data in Table 3.

The room temperature magnetic moment for the [Cu₂(2-bromopropionato)₄(caffeine)₂] complex of $\mu_{\text{eff}} = 1.48$ B.M. is lower than d^9 spin-only moment, $\mu_{\text{eff}} = 1.73$ B.M., and is similar to those observed for other binuclear Cu(II) complexes of the type [Cu₂(RCOO)₄L₂]³⁰ with antiferromagnetic super-exchange interaction between the copper(II) atoms. The magnetic moment of the complex at 80 K decreases to the value of 0.42 B.M., as a consequence of depopulation of the excited triplet ($S = 1$) state.³¹

The solid electronic spectrum of the complex shows a broad absorption band (band I) in the visible region at 13500 cm⁻¹, which could be assigned to a $d_{x^2-y^2} \rightarrow d_{x^2-y^2}$ transition. Moreover, the spectrum displays a shoulder at 26 000 cm⁻¹ (band II). Band II has been assigned to a charge-transfer LMCT absorption and is believed to be indicative of a binuclear complex.³²

The EPR spectrum is characteristic of a triplet state which can be characterized by the singlet-triplet separation $|J|$. The spectrum of the complex may be interpreted in terms of the spin Hamiltonian

$$\hat{H} = \beta B g \bar{S} + D \left[S_y^2 - \frac{1}{3} S(S+1) \right] + E [S_x^2 - S_z^2] \quad (4)$$

where D is the zero-field splitting parameter, x , y and z are the principal axes of the coordinate

system fixed with respect to the Cu...Cu contact, and the other symbols have their usual meaning.³³ The spin parameters are $g_{\perp} = 2.045$, $g_{\parallel} = 2.325$, $|D| = 0.332$ cm⁻¹ and $E = 0.005$ cm⁻¹. The energy separation ($-2J$) between the triplet and singlet states of 310 cm⁻¹ is similar to those found in the series of binuclear copper(II) carboxylates.^{22a, 30a}

In Fig. 5 are shown the Cu...Cu distances plotted against the Cu-N distances of [Cu₂X₄(caffeine)₂] complexes listed in Table 3. The Cu-O bond length of these compounds reveals only small variation. The mean value of Cu-O bond lengths of compounds in Table 3 is 1.97 ± 0.04 Å. If Cu-O distances are kept constant the expression (5) holds which was derived from the BVS model. The best fit of correlation shown in Fig. 5 was achieved using the exponential formula for the bond valence of Cu-N, Cu-O bonds^{4c} and formula $s_{\text{Cu} \cdots \text{Cu}} = (R^*/r_{\text{Cu} \cdots \text{Cu}})^{N^*}$ for the bond valence of Cu...Cu contact. $r_{\text{Cu} \cdots \text{Cu}}$ is internuclear Cu...Cu distance and R^* , N^* are constants.

$$r_{\text{Cu} \cdots \text{Cu}} = R^* \left[2 - \bar{s}_{\text{Cu-O}} - \exp \left(\frac{R_{\text{Cu-N}} - r_{\text{Cu-N}}}{b} \right) \right]^{-1/N^*} \quad (5)$$

Table 3. Selected structural parameters of $[\text{Cu}_2\text{X}_4(\text{caffeine})_2]$ complexes

X	Cu–N (Å)	Cu···Cu (Å)	Cu out of plane deviation (Å)	Φ (°)	R^a	Mark	Ref
Trichloroacetato	2.045(2)	3.0622(7)	0.415(1)	19.11(9)	0.035	A	3b
Trichloroacetato ^b	2.121(6)	2.852(2)	0.315(1)	12.7(2)	0.068	B	15a
Benzoylformato ^b	2.161(4)	2.745(1)	0.259(1)	4.5(2)	0.067	C	15b
Benzoato ^c	2.222(3)	2.6470(7)	0.204(1)	81.6(1)	0.053	D	15c
Naproxenato	2.226(6)	2.649(2)	0.217(1)	79.8(2)	0.071	E	15d
Monochloroacetato	2.23(1)	2.713(2)	0.237(1)	81.7(3)	0.067	F	15e
2-bromopropionato	2.231(6)	2.694(1)	0.221(1)	79.9(2)	0.038	G	This work
2-iodobenzoato	2.238(3)	2.704(6)	0.236(1)	81.6(1)	0.039	H	15f,g

^a $R = \sum ||F_o| - |F_c|| / \sum |F_o|$.

^b Dibenzene solvate.

^c Caffeine solvate.

$\bar{r}_{\text{Cu–O}}$ is the mean value of Cu–O bond valences for the compounds listed in Table 3 and $r_{\text{Cu–N}}$ is internuclear Cu–N distance. $R_{\text{Cu–N}}$ is bond valence parameter of Cu–N bond.^{4c} The fitted values of R^* and N^* are 4.1(1) and 0.71(6). In Fig. 5 is the graph of the function (5) plotted by the full line. The out-of-plane deviation of atom Cu for the binuclear complex unit with the center of symmetry between two Cu atoms in cage is equal to $(r_{\text{Cu} \cdots \text{Cu}} - \text{basal plane-basal plane distance})/2$. The mean value of basal plane-basal plane distance of $2.22 \pm 0.03 \text{ Å}$ ^{1a} was used for calculations (dashed line in Fig. 5). The strong correlation between Cu···Cu distance and out-of-plane deviation of atom Cu has been reported.^{22b} The correlations in Fig 5, fitted by described BVS model suggest predominantly electrostatic nature of Cu···Cu interaction which is consistent with negligibly weak Cu···Cu spin-exchange compared to the spin superexchange through the bridging carboxylates.³⁴ The coordination copper(II) geometry of these complexes is square-pyramid deformed toward the trigonal bipyramid. Such deformation can be expressed by the angle between Cu–N bond line and basal least-squares plane (Φ), Cu out-of plane deviation and Cu–N bond length (Table 3). For the complexes **D–H** is such deformation almost equal. But for the complexes **A–C** deformation of copper(II) coordination geometry increase with elongation of Cu···Cu separation. The dependences shown in Fig. 5 may be caused by the integral

effect of antiferromagnetic spin-exchange interaction between the two copper(II) atoms through the four carboxylate ions,^{3b} electrostatic Cu···Cu interaction within the binuclear cage, and crystal packing.

Conclusions

Halogenopropionate anions can act in copper(II) complexes as bridging ligands, chelate ligands or monodentate ligand. Within a four-membered metallocycles with chelated carboxylic ligands the interatomic distances Cu–O (r_S and r_L) are unambiguously mutually dependent. The dependence r_S vs. r_L is expressed by the equation (3). Increase of the shorter interatomic Cu–O distance (r_S) leads to a strengthening of the longer Cu–O bond, and consequently, to a shortening of the longer Cu–O distance (r_L) and *vice versa*. At the minimal shorter interatomic distance Cu–O (r_S) being 1.92(3) Å long no further increase of the longer Cu–O distance (r_L) takes place. There is a continual transition between the bidentate and monodentate bonding mode of the carboxylato group. Within the $\text{Cu}_2(\text{COO})_4$ cage of binuclear Cu(II) complex the bond lengths C–O slightly depend on the interatomic distance Cu–O of the same Cu–O–C moiety. This dependence ($r_{\text{Cu–O}}$ vs. $r_{\text{C–O}}$) is expressed by the equation (2). For the complexes $[\text{Cu}_2\text{X}_4(\text{caffeine})_2]$ listed in Table 3 the dependence of Cu–N distance vs. Cu···Cu separation is expressed by the

equation (5). If the bond lengths Cu–O are kept constant any elongation of the Cu···Cu separation leads to the shortening of Cu–N distance and *vice versa*. The novel 2-bromopropionato copper(II) complex with caffeine has been prepared and characterized. The crystal structure of $[\text{Cu}_2(2\text{-bromopropionato})_4(\text{caffeine})_2]$ consists of binuclear units connected by the stacking $\pi \cdots \pi$ interactions between adjacent molecules of caffeine in 1-D networks. –Br and –CH₃ substituents of 2-bromopropionato ligands are disordered in major and minor positions. The asymmetric 2-carbon atoms exhibit the chirality of major and invert chirality of minor atoms. The molecular structure and nonbonding character of Cu···Cu contacts within each unit are consistent with the magnetic data and EPR spectra of the substance.

Acknowledgements

This work was sponsored by the grants 1/0056/03 and 1/3572/06 of the Slovak Grant Agency for Science (VEGA). The authors gratefully acknowledge the Professor Alan F. Williams for his helpful discussions.

References

- (a) Melník, M. *Coord. Chem. Rev.* **1982**, *42*, 259. (b) Catterick, J.; Thornton, P. *Adv. Inorg. Chem. Radiochem.* **1977**, *20*, 291. (c) Melník, M. *Coord. Chem. Rev.* **1981**, *36*, 1.
- (a) Gažo, J.; Bersuker, I.B.; Garaj, J.; Kabešová, M.; Kohout, J.; Langfelderová, H.; Melník, M.; Serátor, M.; Valach, F. *Coord. Chem. Rev.* **1976**, *19*, 253. (b) Gažo, J.; Boča, R.; Jóna, E.; Kabešová, M.; Macášková, L.; Šima, J.; Pelikán, P.; Valach, F. *Coord. Chem. Rev.* **1982**, *43*, 87.
- (a) Koh, Y.B.; Christoph, G.G. *Inorg. Chem.* **1979**, *18*, 1122. (b) Uekusa, H.; Ohba, S.; Tokii, T.; Muto, Y.; Kato, M.; Husebye, S.; Steward, O.W.; Chang, Shih-Chi; Rose, J.P.; Pletcher, J.F.; Suzuki, I. *Acta Crystallogr. B* **1992**, *48*, 650.
- (a) Brown, I.D. *Structure and Bonding in Crystals*. Vol. 2; M. O'Keeffe, A. Navrotsky, Eds.; Academic Press: New York, 1981; pp. 1–30. (b) O'Keeffe, M. *Struct. Bonding (Berlin)*. **1989**, *71*, 162. (c) Breese, N.E.; O'Keeffe, M. *Acta Crystallogr. B* **1991**, *47*, 192. (d) Brown, I.D.; Alternatt, D. *Acta Crystallogr. B* **1985**, *41*, 244. (e) Urusov, U.S. *Acta Crystallogr. B* **1995**, *51*, 641. (f) Gopinathan, M.S.; Jug, K. *Theoret. Chim. Acta (Berlin)* **1983**, *63*, 497. (g) Naskar, J.P.; Hati, S.; Datta, D. *Acta Crystallogr. B* **1997**, *53*, 885. (h) Brown, I.D.; Shannon, R.D. *Acta Crystallogr. A* **1973**, *29*, 266. (i) Valach, F. *Polyhedron* **1999**, *18*, 699. (j) Brown, I.D. *The Chemical Bond in Inorganic Chemistry: The Bond-Valence Model*; Oxford University Press: Oxford, 2002.
- Jiang, Z.-H.; Sun, B.-W.; Liao, D.-Z.; Wang, G.-L.; Dahan, F.; Tuchagues, J.P. *Inorg. Chim. Acta* **1998**, *279*, 69.
- Zhang, L.; Li, S.-Q.; Sun, B.-W.; Liao, D.-Z.; Jiang, Z.-H.; Yan, S.-P.; Wang, G.-L.; Yao, X.-K.; Wang, H.-G. *Polyhedron* **1999**, *18*, 781.
- Sharrock, P.; Melník, M.; Belangier-Gariepy, F.; Beachamp, L. *Can. J. Chem.* **1985**, *63*, 2564.
- Koreň, B.; Sivý, P.; Valach, F.; Melník, M.; Ječný, J. *Acta Crystallogr. C* **1988**, *44*, 646.
- Smolander, K.; Melník, M. *Acta Chem. Scand. A* **1984**, *38*, 619.
- Smolander, K. *Acta Chem. Scand. A* **1983**, *37*, 5.
- Moncol, J.; Koman, M.; Melník, M.; Černáková, M.; Glowiak, T. *Polyhedron* **2000**, *19*, 2573.
- Moncol, J.; Mudra, M.; Lönnecke, P.; Koman, M.; Melník, M. *J. Coord. Chem.* **2004**, *57*, 1065.
- Moncol, J.; Svorec, J.; Valko, M.; Melník, M.; Mazur, M.; Koman, M. *Advances in Coordination, Bioinorganic, and Inorganic Chemistry*; M. Melník, J. Šima, M. Tatarko, Eds.; Slovak Technical University Press: Bratislava, 2005; pp. 240–245.
- Cahn, R.S.; Ingold, C.K. *J. Chem. Soc. (London)* **1951**, 612.
- (a) Horie, H.; Husebye, S.; Kato, M.; Meyers, E.A.; Muto, Y.; Suzuki, I.; Tokii, T.; Zingaro, R.A. *Acta Chem. Scand. A* **1986**, *40*, 579. (b) Harada, A.; Tsuchimoto, M.; Ohba, S.; Iwasawa, K.; Tokii, T. *Acta Crystallogr. B* **1997**, *53*, 654. (c) Kawata, T.; Uekusa, H.; Ohbe, S.; Furukawa, T.; Tokii, T.; Muto, Y.; Kato, M. *Acta Crystallogr. B* **1992**, *48*, 253. (d) Koman, M.; Melník, M.; Moncol, J.; Glowiak, T. *Inorg. Chem. Comm.* **2000**, *3*, 489. (e) Koreň, B.; Valach, F.; Sivý, P.; Melník, M. *Acta Crystallogr. C* **1985**, *41*, 1160. (f) Valach, F.; Tokarčík, M.; Maris, T.; Watkin, D.J.; Prout, C.K. *J. Organomet. Chem.* **2001**, *622*, 166. (g) Valach, F.; Koreň, B.; Tokarčík, M.; Melník, M. *Chem. Papers* **1998**, *52*, 140. (h) Melník, M.; Koman, M.; Glowiak, T. *Polyhedron* **1998**, *17*, 1767. (i) Cingi, M.B.; Villa, A.C.; Manfredotti, A.G.; Gaustini, C. *Cryst. Struct. Commun.* **1972**, *1*, 363. (j) Bandoli, G.; Biagini, M.C.; Clemente, D.A.; Rizzardi, G. *Inorg. Chim. Acta* **1976**, *20*, 71.
- Melník, M.; Näsänen, R. *Suomen Kemistilehti B* **1969**, *42*, 298.
- Stoe & Cie, X-RED v. 1.14; Darmstadt, Germany, 1999.
- Main, P.; Fiske, S.J.; Hull, S.E.; Lessinger, L.; Germain, G.; Declercq, J.-P.; Woolfson, M.M. *MULTAN-87, A System of Computer Programs for the Automatic Solution of Crystal Structures from X-Ray Diffraction Data*. University of York: England, and Louvain-la-Neuve: Belgium, 1987.
- Hall, S.R.; Flack, H.D.; Stewart, J.M., Eds. *XTAL3.2 User's Manual*. University of Western Australia and Maryland 1992.
- Farugia, L.J. *J. Appl. Cryst.* **1997**, *30*, 565.
- Earnshaw, A. *Introduction to Magnetochemistry*. Academic Press: London, 1968.
- (a) Melník, M. *Coord. Chem. Rev.* **1982**, *47*, 293. (b) Sundberg, M.R.; Uggla, R.; Melník, M. *Polyhedron* **1996**, *15*, 1157.
- Yablokov, J.V.; Mosina, M.N.; Simonov, Yu.A.; Milkova, L.N.; Ablov, A.V. *Zh. Strukt. Khim.* **1978**, *19*, 42.
- Cahn, R.S.; Ingold, Ch.; Prelog, V. *Angew. Chem. Intern. Edit.* **1966**, *5*, 385.
- Flack, H.D. *Acta Crystallogr. A* **1983**, *39*, 876.
- Janiak, C. *J. Chem. Soc., Dalton Trans.* **2000**, 3885.
- (a) Allan, F.H. *Acta Crystallogr. B* **2002**, *58*, 380. (b) Allan, F.H.; Motherwell, W.D.S. *Acta Crystallogr. B* **2002**, *58*, 407.

28. Valach, F.; Tokarčík, M.; Maris, M.; Watkin, D.J.; Prout, C.K. *Z. Kristallogr.* **2000**, *215*, 56.
29. Havlicek, L.L.; Crain, R.D.; *Practical Statistics for the Physical Sciences*; American Chemical Society: Washington, DC 1988; pp. 109–110.
30. (a) Melník, M. *Coord. Chem. Rev.* **1982**, *42*, 259. (b) Mikloš, D.; Segla, P.; Palicová, M.; Kopcová, M.; Melník, M.; Valko, M.; Glowiak, T.; Korabík, M.; Mrozinski, J. *Polyhedron* **2001**, *20*, 1867.
31. Kahn, O. *Molecular Magnetism*, VCH publishers: Weinheim, 1993.
32. Melník, M.; Potočník, I.; Macášková, L.; Mikloš, D. *Polyhedron* **1996**, *15*, 2159.
33. Abragam, A.; Bleaney, B. *EPR of transition metal complexes*, Clarendon Press: Oxford, 1970.
34. Yamanaka, M.; Uekusa, H.; Ohba, S.; Saito, Y.; Iwata, S.; Kato, M.; Tokii, T.; Muto, Y.; Steward, O.W. *Acta Crystallogr. B* **1992**, *47*, 650.