

The Role of *Aeromonas* in Diarrhea: a Review

A. von Graevenitz

Abstract

The evidence for an enteropathogenic role of *Aeromonas* spp. is still controversial. This review examines various parameters related to a causative role of *Aeromonas* and concludes that infraspecific subsets of strains with a particular array of enterotoxin genes are potential enteropathogens. The consequences for stool bacteriology are discussed.

Infection 2007; 35: 59–64
DOI 10.1007/s15010-007-6243-4

Introduction

The genus *Aeromonas* [1] consists of facultatively anaerobic Gram-negative rods that are predominantly motile by a single polar flagellum and produce oxidase, catalase, nitrate reductase, and an array of exoenzymes. Most species are mesophilic. Members occur in water sources, soil, and foodstuffs. They can be agents of disease in poikilothermic animals (e.g., fish, reptiles) and in man.

The following infections in man have become known [2]:

- (a) wound infections or cellulites related to water or soil exposure;
- (b) septicemia, mainly associated with malignancies (particularly hematological) or hepatobiliary disease, rarely with diabetes mellitus or no immune defect;
- (c) localized extraintestinal infections such as meningitis, peritonitis, otitis media, endocarditis, osteomyelitis, etc.

The first case of a human *Aeromonas* infection reported was one of myositis, observed in 1954 in a woman from Jamaica [3]; and the first series of infections was published from the author's laboratory in 1968 [4]. Since 1961, when the first strain was isolated from human feces [5], aeromonads have been incriminated as agents of diarrhea but the evidence for enteropathogenicity has never been unequivocal [6] because various yardsticks used to solve the question have proven insufficient, including Koch's original postulates and their expansion into a "unified concept" by Evans [7]. For the clinician, the clinical microbiology laboratory, and public health authorities, however, a solution would be of considerable importance: a positive answer would mean

to search for aeromonads in practically every stool specimen by means of selective media (which are different from those used for other bacterial enteropathogens) [8], and to decide whether to treat patients with fecal aeromonads and whether to regularly check drinking water for the presence of these organisms. In fact, the US Environmental Protection Agency has already included *A. hydrophila* in the Contaminant Candidate List of organisms which require future regulation under the Safe Drinking Water Act [9].

The purpose of this review is to examine and interpret the data that have accumulated over approximately 40 years and to make some, albeit preliminary, recommendations for the clinical laboratory. Right now, most laboratories have their own rules regarding the isolation of aeromonads from stools. *Aeromonas*-associated diarrhea (A.-a.d.) will be defined as loose stools associated with the isolation of aeromonads in the stool.

Diarrhoic Patients vs Controls

Of the 17 phenospecies of the genus *Aeromonas* [1, 2], only four have been isolated from human feces with significant frequencies: *A. hydrophila*, *A. caviae*, *A. veronii* biovar. *sobria* (formerly "*A. sobria*"), and *A. trota*; i.e., hybridization groups (HG) 1,4,8, and 14. The latter is a "newcomer" whose frequency was probably underestimated in earlier studies since it is susceptible to ampicillin, an agent used in many selective media for aeromonads [8]. *A. jandaei* (HG 9) and *A. schubertii* (HG 12) have been rare fecal isolates [10]. If these species are agents of diarrhea, one should expect that they are more frequently present in patients with A.-a.d. than in controls [7]. Appropriate studies would have used such controls, selective media, speciation of the

A. von Graevenitz

Institute of Medical Microbiology, University of Zurich,
8028 Zurich, Switzerland; e-mail: avg@immv.unizh.ch

Presented at the summer school workshop, "Rethinking Koch's Postulates", Oct. 14–16, 2005, Berlin-Schwanenwerder.

Dedicated to Professor F.-H. Caselitz, who first described an *Aeromonas* infection.

Received: August 31, 2006 • Revision accepted: January 8, 2007

isolates, and would have eliminated samples containing aeromonads together with other enteropathogens.

While not all studies have included enteropathogenic *Escherichia coli*, viruses, and parasites, results of those available have been contradictory. Not only did the ranking of aeromonads differ from first [11–13] to fifth [14] among enteropathogenic bacteria, but there were also differences in seasonal distribution and predominating species. Some studies observed isolation peaks in warmer seasons [12, 13, 15], others saw no seasonal preference [16]. *A. caviae* was the predominating species in most studies, followed by *A. hydrophila* and *A. veronii* biovar. *sobria* [6, 12, 13, 16, 17]. *A. hydrophila* predominated in Brazil [18] and Thailand [19]; in Bangladesh, it was initially *A. caviae* [20] but later *A. trota* [21] (it should be kept in mind that later studies used more advanced methods of identification and differentiation). In Finnish residents traveling in Morocco, *A. veronii* biovar. *sobria* came in first [22]. While some studies did find significant differences in the frequency of aeromonads isolated from diarrhoic patients including travelers [22] as opposed to controls [12, 14–16], others did not [17, 23, 24]. One study found differences in children – more prone to diarrhea anyway – but not in adults [25]. In Thailand, the difference was significant for Peace Corps volunteers but not for indigenous people [19]. The average isolation rate from diarrhoic stools was 0.8 to 7.4%; in healthy controls, it was 0.4 to 2.1% [26]. Individual studies, however, vary widely in this regard, e.g., a study of infants in Peru found 52.4% in diarrhoic patients vs 8.7% in controls [12]. Of note, if the concomitant enteropathogens were removed from the statistics and isolates were speciated, the significant difference between patients and controls would disappear in this study. The frequency of copathogens was also high in India and Bangladesh (ci. 40%; 10,14) and in Chile (50%, 23). Control groups showed frequencies of 0 to 2.1% in Western Europe and the USA [15, 17, 25] and higher ones in warmer climates (4.8% in Bangladesh, 5.2% in Chile) [14, 23]; in Peru, they ranged from 0 to 21.4% [12]. The carrier state was transient [16], sometimes involving multiple phenotypes [10, 13, 27]. Ranking of individual species was generally equal in patients and controls [12, 17]. Finally, a quantitative study showed considerable variations in the number of colony-forming units (CFU) of aeromonads during different sampling periods, and CFU overlapped between patients (3.5 to 9.9 log₁₀ CFU/gm stool) and controls (5.7 to 6.0 log₁₀ CFU/gm) [11]. In another study, the amount of growth on a selective (Monsur agar) plate was not associated with the presentation of diarrhea [6].

Symptomatology

Symptoms of A.-a.d. were quite variable. Stool consistency varied from loose to watery to bloody; and diarrhea was either self-limited, lasting up to one week, or took a prolonged course of up to two weeks, or became “chronic” with more than one month duration [11, 13, 15]. Nausea,

abdominal cramps, fever, and vomiting were observed only in part of the patients [11, 13]. In one study, duration was longer and fever more prominent with *A. caviae* than with *A. hydrophila* or *A. veronii* biovar. *sobria* [13]. While in the majority of patients the small intestine seemed to be affected only, up to one-third showed colitis by endoscopy [11, 28, 29]. A few cases of hemolytic-uremic syndrome (HUS) following A.-a.d. [2, 30, 31] but no cases of reactive arthritis have been reported.

Similar to some cases of bacterial gastroenteritis, low stomach acidity, liver and gastrointestinal disease as well as recent therapy with antimicrobials ineffective against aeromonads have been reported as associated factors [11, 13, 32]. AIDS patients, however, did not show a higher frequency of A.-a.d. than non-AIDS patients [33].

Cessation of diarrhea following therapy with antimicrobials effective against aeromonads has been reported [11] but its significance for a causation of diarrhea remains questionable in view of the frequently self-limited course of A.-a.d. More significant would be the disappearance of diarrhea or of colonic lesions together with newly negative stool cultures, as has been reported for single cases [11, 28, 29, 34]. This could, of course, also be due to a self-limited disease or to a transient state of colonization with aeromonad(s) [16, 27] in the presence of other enteropathogens. Unfortunately, follow-up typing of carrier isolates has never been done.

Common-Source Outbreaks and Transmission of Fecal Isolates

Although outbreaks are well known for enteropathogenic bacteria, the evidence for outbreaks with aeromonads is almost non-existent in spite of their ubiquitous occurrence in water sources like drinking water [9] and in foods such as seafood, meat, and vegetables [35].

The first outbreaks of A.-a.d. were reported from two day care centers where 6 of 25 and 5 of 24 diarrhoic children, respectively, yielded fecal *Aeromonas* strains [36]. However, the PFGE patterns differed in all except two strains. Of 17 diarrhoic patients from Wisconsin whose stools contained aeromonads, 14 had their drinking water examined for these organisms; only one source was positive. Additional isolates from private wells were unrelated by PFGE to patient isolates [9]. A similar result was obtained in another study using ribotyping [37]. Conversely, during a 68-week observation period in Melbourne in which 795 stool samples from diarrhoic patients were examined, none yielded aeromonads but half of the drinking water samples of the patients, drawn weekly, yielded the organisms [38]. In another study from Wisconsin, A.-a.d. was more often associated with consumption of aeromonad-free drinking water than with consumption of untreated water [39]. Seemingly in contradiction was a study associating A.-a.d. with drinking untreated water but that water had not been examined for aeromonads [32].

Two tropical outbreaks have been reported. In one [40], 28 of 69 patients admitted during one month to a Benghazi, Libya, hospital had “*A. sobria*” isolated from their stools. A source could not be found. The biochemical characteristics suggested *A. hydrophila* but the identification was questionable because the strains produced hydrogen sulfide. In the other report [41], six children in a hematology-oncology unit in Chandigarh, India, developed acute A.-a.d. with “*A. sobria*” during a four-week period. All isolates had similar biotypes and antibiograms. Since neither study had used controls or molecular typing they cannot be used as proof for the presence of an outbreak.

That molecular methods are crucial in this connection was also shown in cases where transmission might have occurred from food, the environment or from patient(s) to patient(s). In one suggestive case of transmission from egg salad, no typing had been done [42]. In a further study mentioned previously [31], several isolates of “*A. sobria*” were cultured from an aquarium with which an infant had contact. The infant developed A.-a.d. and HUS. While all *A. sobria* isolates produced aerolysin (see below) and the diarrhea disappeared with the disappearance of *A. sobria* from the stool, the isolates from the aquarium and from the stool were not identical in their nucleotide sequences (in view of the multiplicity of environmental types it may, however, have been possible that not enough colonies had been checked). In this connection, it is of interest that studies on *Aeromonas* strains from patients' environments have revealed that identical ribotypes may be found in epidemiologically related, but also in unrelated strains [43]. In one instance, *A. caviae* was isolated from three children, from their mother, and from well water used by them. Ribotyping showed identical types in the children but unique patterns in the mother and in the well water [44]. Cultures of *Aeromonas* strains in India obtained during a two-year period have shown multiple types, virulence genes, and antibiograms [10].

Only two studies have found identical ribotypes to suggest transmission, one from food (shrimp) to a patient with A.-a.d. [45], and another one from a child to her foster parents [44]. The outbreak of food poisoning with diarrhea after consumption of Swedish “landgang”, however, from which 6 to 7 log₁₀ CFU/gm of enterotoxin-producing *A. hydrophila* was isolated, lacked stool cultures to prove the causative agent [46].

Experimental Infections

Surprising for an allegedly enteropathogenic organism, oral challenge of 57 human volunteers with 5 different *A. hydrophila* strains producing cytotoxin and cholera toxin-reactive factor (presumably Act) as well as “hemolysin” (see below), using 4 to 10 log₁₀ CFU, led only to mild/moderate diarrhea in two participants who received 7 and 9 log₁₀ CFU, respectively. 55 remained unaffected, although most became colonized [47]. Likewise, oral challenge of protein-malnourished mice [48] or of mice pretreated with strep-

tomycin [49] or clindamycin [50] led to colonization only. In the highly artificial RITARD (removable intestinal tie adult rabbit diarrhea) model [51] using 12 isolates of three *Aeromonas* species from A.-a.d. patients and from carriers diarrhea, bacteremia, and death were strain-dependent. Diarrhea was seen in 23 of 37 rabbits who died and in 11 of 27 who survived ($p = 0.15$). Bacteremia was found in 36 of the moribund ones and in 2 of the survivors. Colonization was detected in all animals but, as in studies mentioned previously [16, 27], lasted only up to 10 days. The ileum was the most heavily colonized part of the intestine.

Host Response

Similar to other diarrheas in which the causative agent is found in the intestine only, serum assays for antibody responses in A.-a.d. have shown insufficient sensitivity and specificity. 7 of 10 sera from patients with A.-a.d. showed no agglutinating antibodies; 5 of these 7 had titers of 1:16 to 1:64 by passive hemagglutination, but that was similar to controls [32]. An ELISA for IgM and IgG responses to the lipopolysaccharides (LPS) of homologous strains had a sensitivity of 30% and a specificity (vs healthy blood donors) of 74% [52]. A cytotoxin-neutralizing assay had a 46% sensitivity and a 94% specificity; only patients with severe acute diarrhea over 60 years of age reacted positively [52]. In a case of HUS following A.-a.d., these neutralizing antibodies showed rising titers [30]. SIgA responses to fecal extracts of 13 patients with A.-a.d. showed fourfold or more titer increases in 11 by dot-blot and in 8 by Western blot but were limited to *A. hydrophila* and “*A. sobria*”; two patients with *A. caviae* did not react [53]. If extracellular products similar to hemolytic toxins were used, specific SIgA responses were seen in 11 of 13 extracts from patients with A.-a.d. [54]. Neither study had examined carriers or blood donors.

Virulence Factors: Occurrence

Aeromonads may produce a variety of biologically active extracellular substances similar to virulence factors known in other enteropathogenic bacteria. They include fimbriae, flagella, outer membrane proteins, an S layer, lipopolysaccharides (endotoxin), capsules, proteases, glycerophospholipid:cholesterol acyl transferase (GCAT), and siderophores (amonabactin, enterobactin) [26, 55]. Their relationship to enteropathogenicity has not been elucidated except that type IV pili have been reported to be associated with gastroenteritis [56].

Toxins of particular interest for gastrointestinal infections, i.e., enterotoxins, are also present in *aeromonads* [26, 55] and were found to be chromosomally encoded [57]. All cause fluid secretion in rabbit ileal loops. They include:

- the cytotoxic, heat-labile (56 °C, 20 min) type II-secreted enterotoxin Act which is also hemolytic and

- destroys the intestinal epithelium via cytokines and activation of the arachidonic acid metabolism;
- two (the heat-labile *Alt* and the heat-stable *Ast*) cytotoxic enterotoxins which cause an increase in intracellular cAMP and prostaglandins and elongate Chinese hamster ovary (CHO) cells;
- various hemolysins, among them *AerA*, *HlyA*, *Ahh1*, *Asa1* [59, 60], and the “*A. sobria*” hemolysin [61] which increases intracellular cAMP. It should be emphasized that hemolysis on blood agar plates used in the clinical laboratories is not a reliable indicator of hemolysin production since it depends on other factors such as the animal source of the erythrocytes and, in experimental models, the number of passages in ileal loops [51, 62, 63]. Generally, erythrocytes are hemolyzed on sheep blood agar plates by *A. hydrophila* and *A. veronii* biovar. *sobria* but rarely by *A. caviae*.

Here again, studies have found the cytotoxin more frequently in *Aeromonas* isolates from diarrhoic patients than from carriers [64] while in Brazil, the heat-stable cytotoxic toxin, the cytotoxin and adhesins were not more frequent in diarrhoic patients than in controls [18]. Others found them more frequently in isolates of *A. hydrophila* than in those of *A. caviae* and *A. veronii* biovar. *sobria* but not more often in diarrhoic than in carrier or environmental strains [23]. The presence of various hemolysins did not show any relationship to the occurrence of diarrhea in one study [6].

Fecal, extraintestinal, and environmental strains of *A. hydrophila*, *A. caviae*, and *A. veronii* biovar. *sobria* were recently found to have a type III secretion system [65].

Virulence Factors: Genetic Analysis

In a study of Bangladeshi children [21], DNA probes were used for identification of the genes for the enterotoxins. *Act* was never found alone, and only in the species *A. hydrophila* together with the other two genes. Diarrhoic children yielded significantly higher numbers of isolates carrying *alt* plus *ast* than carriers and environmental samples but yielded lower numbers of isolates than environmental strains that carried *ast* only. For isolates with *alt* alone and for different *Aeromonas* species, significant differences in the frequency of these genes between the three types of samples could not be detected. The presence of *alt* plus *ast* was associated with watery stools whereas *alt* alone was associated with loose stools. It was concluded that *alt* plus *ast* may synergistically lead to more severe diarrhea, whereas *act* was found associated with bloody diarrhea [21]. In the Indian study [10], only diarrhoic isolates were investigated and yielded a similar distribution as in [21] for *alt* between the species but not for *act* and *ast*. Transposon mutants of an A.-a.d. isolate of *A. hydrophila* which affected the transcription of *act* resulted in reduced virulence by 2 log₁₀ in mice when injected intraperitoneally. Culture filtrates of isogenic mutants with a truncated *act* caused no damage to

the microvilli, did not show hemolytic activity any more, and showed reduced mouse virulence by 3 log₁₀. Reintegration of the native *act* resulted in complete restoration of its former activity [67]. Knockout mutants caused by marker-exchange mutagenesis showed reductions in fluid secretions in ileal loops by 64, 48, and 43% in mutants affecting *act*, *alt*, and *ast*, respectively [66].

Double knockout mutants showed reductions of 36, 62, and 73% in *alt* plus *ast*, *act* plus *ast*, and *act* plus *alt*, respectively. A triple knockout mutant failed to elicit fluid secretion. The greatest damage, thus, seemed to come from *act*, followed by *alt* and *ast* [67].

Genes for hemolysis were found in a majority of *A. hydrophila* and *A. veronii* biovar. *sobria*, but not of *A. caviae*, from diarrhoic patients [6, 60]. The absence of *hly* and *aer* has led to a 20-fold reduction in the LD 50 in a suckling mouse model [61]. Deletion of the gene encoding the *Aeromonas* OMP which is a structural component of the type III secretion system resulted in a significant reduction in inflammatory cytokines and chemokines in the sera of infected mice [68].

Conclusions

At this time, it seems as if only certain infraspecific subsets of aeromonads provided with certain genes for enterotoxicity are significantly associated with diarrhea, the best evidence arising from genetic analysis. Falkow's proposed “molecular Koch Postulates” [69] would almost have been fulfilled, depending on the interpretation of the first postulate. Unfortunately, this will be of little help to the clinical microbiologist and to public health personnel. The microbiologist may, however, base his judgment upon the obvious rarity of person-to-person transmission, the lack of proven outbreaks and of experimental pathogenicity for man, and the frequently self-limited character of A.-a.d. At the present time, he may, therefore, decide to limit the search for aeromonads in stools to those from patients with chronic diarrhea for which no other cause can be found. The epidemiologist will, however, insist on isolation and species identification, if not typing. The fact that recovery of aeromonads from water did not correlate with recovery from stool suggests caution in recommending public health measures to control aeromonads in water.

References

1. Martin-Carnahan A, Joseph SW: Genus I *Aeromonas* Stanier 1943. In: Brenner DJ, Krieg NR, Staley JT (eds): *Bergey's Manual of systematic bacteriology* (2nd edn). Springer, Berlin 2005, 2: 557–578.
2. Figueras MJ: Clinical relevance of *Aeromonas*. *Rev Med Microbiol* 2005; 16: 145–153.
3. Hill KR, Caselitz FH, Moody LM: A case of acute metastatic myositis caused by a new organism of the family Pseudomonadaceae. *West Indian Med J* 1954; 3: 9–11.

4. von Graevenitz A, Mensch AH: The genus *Aeromonas* in human bacteriology. Report of 30 cases and review of the literature. *New Engl J Med* 1968; 278: 245–249.
5. Lautrop H: *Aeromonas hydrophila* isolated from human feces and its possible pathological significance. *APMIS* 1961; 51/5 144: 299–301.
6. Chu YW, Wong CH, Tsang GKL, Kwok MSW, Wong RKO, Lo JYC, Kam KM: Lack of association between presentation of diarrheal symptoms and faecal isolation of *Aeromonas* spp. amongst outpatients in Hong Kong. *J Med Microbiol* 2006; 55: 349–351.
7. Evans AE: Causation and disease: the Henle-Koch postulates revisited. *Yale J Biol Med* 1976; 49: 175–195.
8. Moyer NP, Geiss HK, Marinescu M, Rigby A, Robinson J, Altwegg M: Media and methods for isolation of aeromonads from fecal specimens. A multilaboratory study. *Experientia* 1991; 47: 409–412.
9. Borchardt MA, Stemper ME, Standridge JH: *Aeromonas* isolates from human diarrheic stool and groundwater compared by pulsed-field gel electrophoresis. *Emerg Infect Dis* 2003; 9: 224–228.
10. Sinha S, Shimada T, Ramamurthy T, Bhattacharya SK, Yamasaki S, Takeda Y, Balakrishna Nair G: Prevalence, serotype distribution, antibiotic susceptibility and genetic profiles of mesophilic *Aeromonas* species isolated from hospitalized diarrhoeal cases in Kolkata, India. *J Med Microbiol* 2004; 53: 527–534.
11. George WL, Nakata MM, Thompson J, White M: *Aeromonas*-related diarrhea in adults. *Arch Intern Med* 1985; 145: 2207–2211.
12. Pazzaglia G, Sack RB, Salazar E, Yi A, Chea E, Leon-Barua R, Guerrero CE, Palomino J: High frequency of coinfecting enteropathogens in *Aeromonas*-associated diarrhea of hospitalized Peruvian infants. *J Clin Microbiol* 1991; 29: 1151–1156.
13. Moyer NP: Clinical significance of *Aeromonas* species isolated from patients with diarrhea. *J Clin Microbiol* 1987; 25: 2044–2048.
14. Albert MJ, Faruque ASG, Faruque SM, Sack RB, Malahanabis D: Case-control study of enteropathogens associated with childhood diarrhea in Dhaka, Bangladesh. *J Clin Microbiol* 1999; 37: 3458–3464.
15. Agger WA, McCormick JD, Gurwith MJ: Clinical and microbiological features of *Aeromonas hydrophila*-associated diarrhea. *J Clin Microbiol* 1985; 21: 909–913.
16. Challapalli M, Tess BR, Cunningham DG, Chopra AK, Houston CW: *Aeromonas*-associated diarrhea in children. *Pediatr Infect Dis J* 1988; 7: 693–698.
17. Figura N, Marri L, Verdiani S, Ceccherini C, Barberi A: Prevalence, species differentiation, and toxigenicity of *Aeromonas* strains in cases of childhood gastroenteritis and in controls. *J Clin Microbiol* 1986; 23: 595–599.
18. Bauab TM, Levy CE, Rodrigues J, Falcao DF: Niche-specific association of *Aeromonas* ribotypes from human and environmental origin. *Microbiol Immunol* 2003; 47: 7–16.
19. Pitarangsi C, Echeverria P, Whitmire R, Tirapat C, Formal S, Dammin GJ, Tingtalapong M: Enteropathogenicity of *Aeromonas hydrophila* and *Plesiomonas shigelloides*: prevalence among individuals with and without diarrhea in Thailand. *Infect Immun* 1982; 35: 666–673.
20. Kühn I, Albert MJ, Ansaruzzaman M, Bhuiyan MA, Alabi SA, Islam MS, Neogi PKB, Huys G, Janssen P, Kersters K, Möllby R: Characterization of *Aeromonas* spp. isolated from humans with diarrhea, from healthy controls, and from surface water in Bangladesh. *J Clin Microbiol* 1997; 35: 369–373.
21. Albert MJ, Ansaruzzaman M, Talukder KA, Chopra AK, Kühn I, Rahman M, Faruque ASG, Islam MS, Sack RB, Mollby R: Prevalence of enterotoxin genes in *Aeromonas* spp. isolated from children with diarrhea, healthy controls, and the environment. *J Clin Microbiol* 2000; 38: 3785–3790.
22. Hänninen ML, Salmi S, Mattila L, Taipalinen R, Siitonen A: Association of *Aeromonas* spp. with travelers' diarrhoea in Finland. *J Med Microbiol* 1995; 42: 26–31.
23. Figueroa G, Galeno H, Soto V, Troncoso M, Hinrichsen V, Yudelevich A: Enteropathogenicity of *Aeromonas* species isolated from infants: a cohort study. *J Infect* 1988; 17: 205–213.
24. Svenungsson B, Lagergren K, Ekwall E, Evengard B, Hedlund KO, Karnell A, Lofdahl S, Svensson L, Weintraub A: Enteropathogens in adult patients with diarrhea and healthy control subjects: a 1-year prospective study in a Swedish clinic for infectious diseases. *Clin Infect Dis* 2000; 30: 770–778.
25. Geiss HK, Fogel W, Sonntag HG: Häufigkeit von *Aeromonas* species im Stuhl von Gesunden und Durchfallkranken. *Immun Infekt* 1988; 16: 115–117.
26. Thornley JP, Shaw JG, Gryllos IA, Eley A: Virulence properties of clinically significant *Aeromonas* species: evidence for pathogenicity. *Rev Med Microbiol* 1997; 8: 61–72.
27. Pazzaglia G, Escalante JR, Sack RB, Rocca C, Benavides V: Transient intestinal colonization with multiple phenotypes of *Aeromonas* species during the first week of life. *J Clin Microbiol* 1990; 28: 1842–1846.
28. Deutsch SF, Wedzina W: *Aeromonas sobria*-associated left-sided segmental colitis. *Am J Gastroenterol* 1997; 92: 2104–2106.
29. Farraye FA, Peppercorn MA, Ciano PS, Kavesh WN: Segmental colitis associated with *Aeromonas hydrophila*. *Am J Gastroenterol* 1989; 84: 436–438.
30. Bogdanovic R, Cobeljic M, Markovic V, Nikolic V, Ogujanovic M, Sarjanovic L, Makic D: Hemolytic-uremic syndrome associated with *Aeromonas hydrophila* enterocolitis. *Pediatr Nephrol* 1991; 5: 293–295.
31. Filler G, Ehrlich JHH, Strauch E, Beutin L: Acute renal failure in an infant associated with cytotoxic *Aeromonas sobria* isolated from patient's stool and from aquarium water as suspected source of infection. *J Clin Microbiol* 2000; 38: 469–470.
32. Holmberg SD, Schell WL, Fanning GR, Wachsmuth IK, Hickman-Brenner FW, Blake PA, Brenner DJ, Farmer JJ: *Aeromonas* intestinal infections in the United States. *Ann Intern Med* 1986; 105: 683–689.
33. Suthienkul O, Aiumlaor P, Siripanichgon K, Eampokalap B, Likhanon-sakul S, Utrachakij F, Rakue Y: Bacterial causes of AIDS-associated diarrhea in Thailand. *Southeast Asian J Trop Med Public Health* 2001; 32: 158–170.
34. Nathwani D, Laing RBS, Harvey G, Smith CC: Treatment of symptomatic enteric *Aeromonas hydrophila* infection with ciprofloxacin. *Scand J Infect Dis* 1991; 23: 653–654.
35. Palumbo S: The *Aeromonas hydrophila* group in food. In: Austin B, Altwegg M, Gosling PJ, Joseph SW (eds): *The Genus Aeromonas*. Wiley, New York, pp 1996; 287–310.
36. de la Morena M, Van R, Singh K, Brian M, Murray BE, Pickering LK: Diarrhea associated with *Aeromonas* species in children in day care centers. *J Infect Dis* 1993; 168: 215–218.
37. Moyer NP, Martinetti Lucchini G, Holcomb LA, Hall NH, Altwegg M: Application of ribotyping for differentiating aeromonads isolated from clinical and environmental sources. *Appl Env Microbiol* 1992; 58: 1940–1944.
38. Hellard ME, Sinclair MI, Forbes AB, Fairley CK: A randomized, blinded, controlled trial investigating the gastrointestinal health effects of drinking water quality. *Environ Health Perspect* 2001; 109: 773–778.
39. Agger WA, Callister SM: Intestinal infections with *Aeromonas*. *Ann Intern Med* 1987; 106: 479.
40. Taher AA, Rao BN, Alganay KG, el-Arabi MB: An outbreak of acute gastroenteritis due to *Aeromonas sobria* in Benghazi, Libyan Arab Jamahiriya. *East Mediterr Health J* 2000; 6: 497–499.

41. Taneja N, Khurana S, Trehan A, Marwaha RK, Sharma M: An outbreak of hospital-acquired diarrhea due to *Aeromonas sobria*. *Indian Pediatr* 2004; 41: 912–916.
42. Bottone E: Correlation between known exposure to contaminated food or surface water and development of *Aeromonas hydrophila* and *Plesiomonas shigelloides* diarrhea. *Med Microbiol Lett* 1993; 2: 217–225.
43. Demarta A, Tonolla M, Caminada AP, Beretta M, Peduzzi R: Epidemiological relationships between *Aeromonas* strains isolated from symptomatic children and household environments as determined by ribotyping. *Eur J Epidemiol* 2000; 16: 447–453.
44. Moyer NP, Martinetti G, Lüthy-Hottenstein J, Altwegg M: Value of rRNA gene restriction patterns of *Aeromonas* spp. for epidemiological investigations. *Curr Microbiol* 1992; 24: 15–21.
45. Altwegg M, Martinetti Lucchini G, Lüthy-Hottenstein J, Rohrbach M: *Aeromonas*-associated gastroenteritis after consumption of contaminated shrimp. *Eur J Clin Microbiol* 1991; 10: 4.
46. Krovacek K, Dumontet S, Eriksson E, Baloda SB: Isolation and virulence profiles of *Aeromonas hydrophila* implicated in an outbreak of food poisoning in Sweden. *Microbiol Immunol* 1995; 39: 655–661.
47. Morgan DP, Johnson PC, DuPont HL, Satterwhite TK, Wood LV: Lack of correlation between known virulence properties of *Aeromonas hydrophila* and enteropathogenicity for humans. *Infect Immun* 1985; 50: 62–65.
48. Pazzaglia G, Winoto I, Jennings G: Oral challenge with *Aeromonas* in protein-malnourished mice. *J Diarrhoeal Dis Res* 1994; 12: 108–112.
49. Sanderson K, Ghazali FM, Kirov SM: Colonization of streptomycin-treated mice by *Aeromonas* species. *J Diarrhoeal Dis Res* 1996; 14: 27–32.
50. Kelleher A, Kirov S: *Rattus norvegicus*: not a model for *Aeromonas*-associated gastroenteritis in man. *FEMS Immunol Med Microbiol* 2000; 28: 313–318.
51. Pazzaglia G, Sack RB, Bourgeois AL, Froehlich J, Eckstein J: Diarrhea and intestinal invasiveness of *Aeromonas* strains in the removable intestinal tie model. *Infect Immun* 1990; 58: 1924–1931.
52. Kuijper EJ, van Alphen V, Peeters MF, Brenner DJ: Human serum antibody response to the presence of *Aeromonas* spp. in the intestinal tract. *J Clin Microbiol* 1990; 28: 584–590.
53. Jiang ZD, Nelson AC, Mathewson JJ, Ericsson CD, duPont H: Intestinal secretory immune response to infection with *Aeromonas* species and *Plesiomonas shigelloides* among students from the United States in Mexico. *J Infect Dis* 1991; 164: 979–982.
54. Crivelli C, Demarta A, Peduzzi R: Intestinal secretory immunoglobulin (sIgA) response to *Aeromonas* exoproteins in patients with naturally acquired *Aeromonas* diarrhea. *FEMS Immunol Med Microbiol* 2001; 30: 31–35.
55. Chopra AK, Houston CW: Enterotoxins in *Aeromonas*-associated gastroenteritis. *Microbes Infect* 1999; 1: 1129–1137.
56. Barnett TC, Kirov SM, Strom MS, Sanderson K: *Aeromonas* spp. possess at least two distinct type IV pilus families. *Microb Pathog* 1997; 23: 241–247.
57. Brown RL, Sanderson K, Kirov SM: Plasmids and *Aeromonas* virulence. *FEMS Immunol Med Microbiol* 1997; 17: 217–223.
58. Vadivelu J, Puthuchery SD, Navaratnam P: Exotoxin profiles of clinical isolates of *Aeromonas hydrophila*. *J Med Microbiol* 1991; 34: 363–367.
59. Tanoue N, Takahashi A, Okamoto K, Fujii Y, Taketani Y, Harada N, Nakano M, Nakaya Y: A pore-forming toxin produced by *Aeromonas sobria* activates cAMP-dependent Cl-secretory pathways to cause diarrhea. *FEMS Microbiol Lett* 2005; 242: 195–201.
60. Wang G, Clark CG, Liu C, Pucknell C, Munro CK, Kruk TMAC, Caldeira R, Woodward DL, Rodgers FG: Detection and characterization of the hemolysin genes in *Aeromonas hydrophila* and *Aeromonas sobria* by multiplex PCR. *J Clin Microbiol* 2003; 41: 1048–1054.
61. Wong CYF, Heuzenroeder MW, Flower RLP: Inactivation of two hemolytic toxin genes in *Aeromonas hydrophila* attenuates virulence in a suckling mouse model. *Microbiology* 1998; 144: 291–29.
62. Joseph SM, Ali A: *Aeromonas aerolysin* – when systematics and genetics collide. *Med Microbiol Lett* 1993; 2: 314–321.
63. Singh DV, Sanyal SC: Production of hemolysis and its correlation with enterotoxicity in *Aeromonas* spp. *J Med Microbiol* 1992; 37: 262–267.
64. Kuijper EJ, Bol P, Peeters MF, Steigerwalt AG, Zanen HC, Brenner DJ: Clinical and epidemiological aspects of members of *Aeromonas* DNA hybridization groups isolated from human feces. *J Clin Microbiol* 1989; 27: 1531–1537.
65. Chacon MR, Soler L, Groisman EA, Guarro J, Figueras MJ: Type III secretion system genes in clinical *Aeromonas* isolates. *J Clin Microbiol* 2004; 42: 1285–1287.
66. Sha J, Kozlova EV, Chopra AK: Role of various enterotoxins in *Aeromonas hydrophila*-induced gastroenteritis: generation of enterotoxin-gene deficient mutants, evaluation of their enterotoxic activity. *Infect Immun* 2002; 70: 1924–1935.
67. Xu X-J, Ferguson MR, Popov VL, Houston CW, Peterson JW, Chopra AK: Role of a cytotoxic enterotoxin in *Aeromonas*-mediated infections: development of transposon and isogenic mutants. *Infect Immun* 1998; 66: 3501–3509.
68. Fadl AA, Galindo CL, Sha J, Erova TE, Houston CW, Olano JP, Chopra AK: Deletion of the genes encoding type III secretion system and cytotoxic enterotoxin alters host responses to *Aeromonas hydrophila* infection. *Microb Pathog* 2006; 40: 198–210.
69. Falkow S: Molecular Koch's postulates applied to bacterial pathogenicity – a personal recollection 15 years later. *Nature Rev Microbiol* 2004; 2: 67–72.