

Center-stable manifold of the ground state in the energy space for the critical wave equation

Joachim Krieger · Kenji Nakanishi ·
Wilhelm Schlag

Received: 16 May 2013 / Revised: 16 May 2013 / Published online: 3 July 2014
© Springer-Verlag Berlin Heidelberg 2014

Abstract We construct a center-stable manifold of the ground state solitons in the energy space for the critical wave equation without imposing any symmetry, as the dynamical threshold between scattering and blow-up, and also as a collection of solutions which stay close to the ground states. Up to energy slightly above the ground state, this completes the 9-set classification of the global dynamics in our previous paper (DCDS 33:6, 2013). We can also extend the manifold to arbitrary energy size by adding large radiation. The manifold contains all the solutions scattering to the ground state solitons, and also some of those blowing up in finite time by concentration of the ground states.

Keywords Nonlinear wave equation · Sobolev critical exponent · Hamiltonian · Blowup · Scattering · Ground state · Stability · Invariant manifold

Mathematics Subject Classification (2010) 35L70 · 35B40

Support by the Swiss National Science foundation for the first author is acknowledged.
Support of the National Science Foundation DMS-0617854, DMS-1160817 for the third author is gratefully acknowledged.

J. Krieger (✉)
Bâtiment des Mathématiques, EPFL, Station 8, 1015 Lausanne, Switzerland
e-mail: joachim.krieger@epfl.ch

K. Nakanishi
Department of Mathematics, Kyoto University, Kyoto 606-8502, Japan

W. Schlag
Department of Mathematics, The University of Chicago, 5734 South University Avenue,
Chicago, IL 60615, USA

Contents

1	Introduction	2
1.1	Strichartz norms and strong solutions	7
1.2	Symmetry, solitons and linearization	8
1.3	Coordinates around the ground states	9
2	Center-stable manifold around the ground states	13
2.1	Igniting the unstable mode	15
2.2	Construction of the manifold	22
2.3	Smoothness of the manifold	24
3	Extension of the manifold and the 9-set dynamics	27
3.1	Extension by the backward flow	27
3.2	Space-time extension and restriction	28
3.3	Extension by the Lorentz transform	31
4	Detaching the radiation	34
5	Center-stable manifold with large radiation	41
6	One-pass theorem with large radiation	43
	Appendix A: Concentration blowup in the interior of blowup region	47
	Appendix B: Table of notation	49
	References	50

1 Introduction

We study global dynamics of the critical wave equation (CW)

$$\begin{aligned} \ddot{u} - \Delta u &= f'(u) := |u|^{2^*-2}u, \quad 2^* := \frac{2d}{d-2}, \\ u(t, x) &: I \times \mathbb{R}^d \rightarrow \mathbb{R}, \quad I \subset \mathbb{R}, \quad d = 3 \text{ or } 5, \end{aligned} \quad (1.1)$$

in the energy space¹

$$\vec{u}(t) := \begin{pmatrix} u(t) \\ \dot{u}(t) \end{pmatrix} \in \mathcal{H} := \dot{H}^1(\mathbb{R}^d) \times L^2(\mathbb{R}^d). \quad (1.2)$$

Henceforth, the arrow on a function $\vec{u}(t)$ indicates the vector $\vec{u}(t) = (u(t), \dot{u}(t))$ given by a scalar function $u(t)$. We do not distinguish column and row vectors. The above equation CW is in the Hamiltonian form

$$\vec{u}_t = J E'(\vec{u}), \quad J := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (1.3)$$

and so the energy or the Hamiltonian

$$E(\vec{u}(t)) := \int_{\mathbb{R}^d} \frac{|\dot{u}|^2 + |\nabla u|^2}{2} - f(u) dx, \quad f(u) := \frac{|u|^{2^*}}{2^*}, \quad (1.4)$$

¹ The exclusion of $d = 4$ is by the same reason as in [14], namely to preclude type-II blow-up in the scattering region by [6]. The argument in this paper or [14] is not sensitive to the dimension.

is conserved. Another important conserved quantity is the total momentum

$$P(\vec{u}(t)) := \int_{\mathbb{R}^d} \dot{u} \nabla u dx. \quad (1.5)$$

The Nehari functional

$$K(u) := \int_{\mathbb{R}^d} |\nabla u|^2 - |u|^{2^*} dx \quad (1.6)$$

plays a crucial role in the variational argument. CW is invariant under translation, the Lorentz transform, and the scaling

$$u(t, x) \mapsto u_\lambda(t, x) := \lambda^{d/2-1} u(\lambda t, \lambda x), \quad (1.7)$$

which also preserves the energy $E(u_\lambda) = E(u)$, making (CW) special and critical. It also gives rise to the ground state solutions in the explicit form

$$W_\lambda(x) := \lambda^{d/2-1} W(\lambda x), \quad W(x) := \left[1 + \frac{|x|^2}{d(d-2)} \right]^{1-d/2} \in \dot{H}^1(\mathbb{R}^d), \quad (1.8)$$

$$\forall \lambda > 0, \quad -\Delta W_\lambda + f'(W_\lambda) = 0,$$

which has the minimal energy among all the stationary solutions. The scale and translation invariance of CW generates a family of ground states as a smooth manifold in \mathcal{H} with dimension $1 + d$:

$$W_{\lambda,c}(x) := \lambda^{d/2-1} W(\lambda(x-c)) \implies -\Delta W_{\lambda,c} + f'(W_{\lambda,c}) = 0, \quad (1.9)$$

$$S_{\text{static}}(W) := \{\vec{W}_{\lambda,c} \in \mathcal{H} \mid \lambda > 0, c \in \mathbb{R}^d\}.$$

Then the Lorentz invariance generates a family of solitons on a smooth manifold in \mathcal{H} with dimension $1 + 2d$:

$$u(t, x) = W_{\lambda,c,p}(t, x) := W_{\lambda,c+pt}(x + (\langle p \rangle - 1)p|p|^{-2}p \cdot x) \implies (CW).$$

$$S_{\text{soliton}}(W) := \{\vec{W}_{\lambda,c,p}(0) \in \mathcal{H} \mid \lambda > 0, c \in \mathbb{R}^d, p \in \mathbb{R}^d\}. \quad (1.10)$$

Other types of solutions are the scattering (to 0) solutions with the property

$$\exists \varphi \in \mathcal{H}, \quad \|\vec{u}(t) - U(t)\varphi\|_{\mathcal{H}} \rightarrow 0 \quad (t \rightarrow \infty) \quad (1.11)$$

where $U(t)$ denotes the free propagator, defined as the Fourier multiplier

$$U(t) := \begin{pmatrix} \cos(t|\nabla|) & |\nabla|^{-1} \sin(t|\nabla|) \\ -|\nabla| \sin(t|\nabla|) & \cos(t|\nabla|) \end{pmatrix}, \quad |\nabla| := \sqrt{-\Delta}, \quad (1.12)$$

the norm blow-up (called type-I blow-up in [8])

$$\limsup_{t \nearrow t^*} \|\vec{u}(t)\|_{\mathcal{H}} = \infty, \quad (1.13)$$

and the more subtle type-II blow-up, for which $\|\vec{u}(t)\|_{\mathcal{H}}$ is bounded but $\vec{u}(t)$ fails to be strongly continuous in \mathcal{H} beyond some $t < \infty$.

In [14], the authors gave a partial classification of dynamics of (CW) in the region

$$E(\vec{u}) < \sqrt{(E(W) + \varepsilon^2)^2 + |P(\vec{u})|^2} \quad (1.14)$$

for a small $\varepsilon > 0$, which is, by the Lorentz invariance, reduced to the region

$$E(\vec{u}) < E(W) + \varepsilon^2. \quad (1.15)$$

It was proved that if $u \in C([0, T_+); \mathcal{H})$ is a strong solution up to the maximal existence time $T_+ \in (0, \infty]$, which does not stay close to the ground state solitons near $t = T_+$, then u either blows up away from the ground state solitons, or it scatters (to 0) as $t \rightarrow \infty$. We have the same for $t < 0$, and moreover, the 2×2 combinations of scattering and blow-up in $t > 0$ and in $t < 0$ respectively are realized by initial-data sets in \mathcal{H} which have non-empty interior. The key ingredient for proof is the existence of a small neighborhood of the ground states such that any solution exiting from it can never come back again, called the one-pass theorem.

A missing piece in the above result of [14] is the global dynamics around the ground states, compared with the corresponding results for the subcritical Klein-Gordon equation [21] and for the Schrödinger equation in the radial symmetry [20], where we have 3×3 complete classification of (1.14) including the scattering to the ground states on a center-stable manifold of codimension 1.

On the other hand, there have been many papers [1, 3, 4, 9, 10, 16, 17] for (CW) constructing various types of solutions around the ground states, including center-stable manifolds in some stronger topology than the energy space, on which the solutions scatter to the ground states, type-II blow-up at prescribed power law rate or with eternal oscillations between such rates, type-II blow-up at time infinity. The latter phenomena clearly distinguish (CW) from the dynamics of the subcritical equation.

In this paper, we construct a smooth center-stable manifold of codimension 1 in the energy space, which embraces all the solutions scattering to, or staying close to the ground state solitons. Indeed the last property is the defining characterization of the manifold. Plugging it into the result in [14], we complete the 3×3 classification for (CW) in the region (1.14), which is now described.

Denote \mathcal{H} -distance to the ground states by

$$\text{dist}_W(\varphi) := \inf\{\|\vec{u}(t) - \psi\|_{\mathcal{H}} \mid \psi \text{ or } -\psi \in S_{\text{static}}(W)\}, \quad (1.16)$$

and the time inversion for any initial data $\varphi \in \mathcal{H}$ and any initial data set $A \subset \mathcal{H}$ by

$$\varphi^\dagger := (\varphi_1, -\varphi_2), \quad A^\dagger := \{\varphi^\dagger \mid \varphi \in A\}. \quad (1.17)$$

Theorem 1.1 *There exist positive constants $\varepsilon < \delta < 1 < C$, and an unbounded connected C^1 manifold $\mathcal{M} \subset \mathcal{H}$ with codimension 1 satisfying the following. $S_{\text{static}}(W) \subset \mathcal{M}$ is tangent to the center-stable subspace of the linearized equation at each point of $S_{\text{static}}(W)$. \mathcal{M} is invariant by the flow, translation, rotation and the \mathcal{H} -invariant scaling. Let u be any solution with $E(\vec{u}(0)) < E(W) + \varepsilon^2$, and let $T \in (0, \infty]$ be its maximal existence time. Then we have only one of the following (1)–(3).*

- (1) $\vec{u}(0) \notin \pm\mathcal{M}$, $T = \infty$ and $\lim_{t \rightarrow \infty} \|\vec{u}(t) - \vec{v}(t)\|_{\mathcal{H}} = 0$ for some free solution v .
- (2) $\vec{u}(0) \notin \pm\mathcal{M}$, $T < \infty$ and $\varliminf_{t \rightarrow T} \text{dist}_W(\vec{u}(t)) > \delta > C\varepsilon$.
- (3) $\vec{u}(0) \in \pm\mathcal{M}$ and $\varliminf_{t \rightarrow T} \text{dist}_W(\vec{u}(t)) \leq C\sqrt{E(\vec{u}) - E(W)}$.

Let A_1, A_2, A_3 be the corresponding sets of the initial data $\vec{u}(0)$. Then $A_1 \cap A_1^\dagger$ is a non-empty open set. $A_1 \cap A_2^\dagger$, $A_2 \cap A_1^\dagger$ and $A_2 \cap A_2^\dagger$ have non-empty interior. A_1 and A_2 have non-empty interior in $\pm\mathcal{M}^\dagger$ in the relative topology. $\mathcal{M} \cap \mathcal{M}^\dagger \subset \mathcal{H}$ is a connected C^1 manifold with codimension 2. $\mathcal{M} \cap -\mathcal{M} = \emptyset = \mathcal{M}^\dagger \cap -\mathcal{M}$.

The above case (3) also contains blow-up solutions, but they are distinguished from (2) by the asymptotic distance from the ground states. By the characterization of type-II blow-up by Duyckaerts, Kenig and Merle [6, Theorem 1], in case (3) with $T < \infty$, there are a smooth $(\lambda, c) : [0, T) \rightarrow (0, \infty) \times \mathbb{R}^d$ and $\varphi \in \mathcal{H}$ such that

$$\lambda(t) \rightarrow \infty, \quad \|\vec{u}(t) - \vec{W}_{\lambda(t), c(t)} - \vec{\varphi}\|_{\mathcal{H}} \rightarrow 0 \quad (1.18)$$

as $t \nearrow T$. The gap between $C\varepsilon$ and δ is actually huge in the proof. The above theorem except for the existence of \mathcal{M} was essentially proved in [14].

Next we can exploit the Lorentz transform to include all the ground state solitons.

Theorem 1.2 *There exist a small constant $\varepsilon > 0$, a connected C^1 manifold $\mathcal{M}_L \subset \mathcal{H}$ with codimension 1, and two open sets $O_1, O_2 \subset \mathcal{H}$ satisfying the following. $\mathcal{M} \cup S_{\text{soliton}}(W) \subset \mathcal{M}_L$. $S_{\text{soliton}}(W) \subset O_1 \subset \overline{O_1} \subset O_2$. \mathcal{M}_L is invariant by the flow, translation, \mathcal{H} -invariant scaling and the Lorentz transform. Let u be any solution with $E(\vec{u}(0)) < \sqrt{|E(W) + \varepsilon^2|^2 + |P(\vec{u}(0))|^2}$ and let $T \in (0, \infty]$ be its maximal existence time. Then we have only one of the following (1)–(3).*

- (1) $\vec{u}(0) \notin \pm\mathcal{M}_L$, $T = \infty$ and $\lim_{t \rightarrow \infty} \|\vec{u}(t) - \vec{v}(t)\|_{\mathcal{H}} = 0$ for some free solution v .
- (2) $\vec{u}(0) \notin \pm\mathcal{M}_L$, $T < \infty$ and $\vec{u}(t) \notin \pm\overline{O_2}$ for all t near T .
- (3) $\vec{u}(0) \in \pm\mathcal{M}_L$ and $\vec{u}(t) \in \pm O_1$ for all t near T .

The initial data sets and \mathcal{M}_L enjoy the same properties as in the previous theorem.

The manifolds \mathcal{M} and \mathcal{M}_L are center-stable manifolds of $S_{\text{static}}(W)$ and $S_{\text{soliton}}(W)$ respectively, but they contain solutions blowing up in finite time. The invariance by the flow should be understood that the solutions starting on the manifold stay there as long as they exist, and similarly for the Lorentz transform. Again by [6], in case (3) with $T < \infty$, we have a smooth $(\lambda, c), p \in \mathbb{R}^d$ and $\varphi \in \mathcal{H}$ such that, as $t \nearrow T$,

$$\lambda(t) \rightarrow \infty, \quad \|\vec{u}(t) - \vec{W}_{\lambda(t), c(t), p} - \varphi\|_{\mathcal{H}} \rightarrow 0. \quad (1.19)$$

Our argument to construct the center-stable manifold is somehow similar to the numerical bisection in [2, 24], where the center-stable manifold was searched for as the threshold between scattering and blowup. Indeed, our proof does not touch the delicate dynamics of those solutions on the manifold, but relies on the behavior of those off the manifold. In particular, we do not need any dispersive estimate on the linearized operator as in [1, 16, 20, 21], which makes our proof much simpler. In this respect, it is similar to [22] in the subcritical case. On the other hand, the criticality or the concentration phenomenon forces us to work in the space-time rescaled according to the solution itself. For that part we employ the same argument as in the previous paper [14].

The next question is if we can remove the energy restriction (1.14). Concerning it, Duyckaerts, Kenig and Merle [8] recently established an outstanding result of asymptotic soliton resolution for $d = 3$: Every solution with radial symmetry is either type-I blow-up, or decomposes into a sum of ground states with time-dependent scaling and a free solution

$$\exists N \geq 0, \exists \lambda_j(t), \exists \varphi \in \mathcal{H}, \lim_{t \nearrow T} \left\| \vec{u}(t) - \sum_{j=1}^N \vec{W}_{\lambda_j(t), 0} - U(t)\varphi \right\|_{\mathcal{H}} = 0, \quad (1.20)$$

where T is the maximal existence time of u . Given this expansion, one might expect that the dividing manifold of dynamics could be extended as the collection of all such solutions with $N > 0$. However, it is very hard to prove such a statement even if we know the above asymptotics, because of the instability of the ground state. Moreover, one can easily observe that the above naive guess is not correct when $T < \infty$ and the energy is larger, as one can construct such blow-up solutions in the deep interior of blow-up solutions, by using finite speed of propagation (see Appendix A).

Instead of pursuing that approach, we extend our center-stable manifold globally by adding large radiation, thereby including at least all solutions (1.20) with $N = 1$ and $T = \infty$, as well as some of them with $T < \infty$. A simple procedure is proposed to reduce the analysis to the previous case $E < E(W) + \varepsilon^2$ by detaching large radiation, which relies on the asymptotic Huygens principle, valid for all $d \in \mathbb{N}$ and without radial symmetry. The extended manifold splits the energy space into the scattering and blow-up regions locally around itself, although the entire dynamical picture is still far beyond our analysis.

Theorem 1.3 *There exist a connected C^1 manifold $\mathcal{M}_D \subset \mathcal{H}$ with codimension 1 and two open sets $O_3, O_4 \subset \mathcal{H}$ satisfying the following. $\mathcal{M}_L \subset \mathcal{M}_D$. $S_{\text{soliton}}(W) \subset O_3 \subset \bar{O}_3 \subset O_4$. \mathcal{M}_D is invariant by the flow, translation, \mathcal{H} -invariant scaling and the Lorentz transform. Let u be any solution with $\vec{u}(0) \in O_3$. Then we have only one of the following (1)–(3).*

- (1) $\vec{u}(0) \notin \pm \mathcal{M}_D$, $T = \infty$ and $\lim_{t \rightarrow \infty} \|\vec{u}(t) - \vec{v}(t)\|_{\mathcal{H}} = 0$ for some free solution v .
- (2) $\vec{u}(0) \notin \pm \mathcal{M}_D$, $T < \infty$ and $\vec{u}(t) \notin \pm \bar{O}_4$ for t near T .
- (3) $\vec{u}(0) \in \pm \mathcal{M}_D$ and $\vec{u}(t) \in \pm O_3$ for all t near T .

If $u \in C([0, \infty); \mathcal{H})$ is a solution satisfying

$$\lim_{t \rightarrow \infty} \|\vec{u}(t) - \vec{W}_{\lambda(t), c(t), p(t)} - \vec{v}(t)\|_{\mathcal{H}} = 0, \quad (1.21)$$

for some $(\lambda, c, p) : [0, \infty) \rightarrow (0, \infty) \times \mathbb{R}^{1+2d}$ and a free solution v , then $\vec{u}(0) \in \mathcal{M}_D$.

More detailed statements are given in the main body of paper.

This paper is organized as follows. In the rest of this section, we introduce some notation and coordinates, together with a few basic facts and estimates, mostly overlapping with the previous paper [14].

In Part I starting with Sect. 2, we deal with the solutions with energy slightly above the ground state. The center-stable manifold is constructed as a threshold between scattering and blowup, which completes the 9-set classification of dynamics, in a form similar to the subcritical case [21]. The main new ingredient is the ignition Lemma 2.2, which roughly says that for any solution staying close to the ground states, any arbitrarily small perturbation in the unstable direction eventually leads to the ejection from a small neighborhood as in the ejection lemma of [14]. These are extended by the Lorentz transform in the end of Sect. 3.

In Part II starting with Sect. 4, we extend the results in Part I to large energy by adding out-going radiation. The extended manifold contains all the solutions scattering to the ground state solitons, while it is still a dynamical threshold between the scattering and the blowup. The main ingredient is the detaching Lemma 4.4, which allows one to detach out-going radiation energy from a solution to produce another solution with smaller energy but the same behavior. We also extend the one-pass theorem of [14] by allowing out-going large radiation.

1.1 Strichartz norms and strong solutions

For any $I \subset \mathbb{R}$, we use the Strichartz norms for the wave equation with the following exponents

$$\begin{aligned} \text{St}_s &:= L_t^{q_s}(I; \dot{B}_{q_s, 2}^{1/2}(\mathbb{R}^d)), \quad \text{St}_m := L_{t,x}^{q_m}(I \times \mathbb{R}^d), \quad \text{St}_s^* := L_t^{q'_s}(I; \dot{B}_{q'_s, 2}^{1/2}(\mathbb{R}^d)), \\ \text{St}_p &:= L_t^{(d+2)/(d-2)}(I; L^{2(d+2)/(d-2)}(\mathbb{R}^d)) \quad (d \leq 6), \\ \text{St}(I) &:= \text{St}_s(I) \cap \text{St}_m \cap \text{St}_p(I), \quad q_s := \frac{2(d+1)}{d-1}, \quad q_m := \frac{2(d+1)}{d-2}, \end{aligned} \quad (1.22)$$

where $q' := q/(q-1)$ denotes the Hölder conjugate. Slightly abusing the notation, we often apply these norms to the first component of vector functions such as

$$\|(v_1, v_2)\|_{\text{St}(I)} := \|v_1\|_{\text{St}(I)}. \quad (1.23)$$

The small data theory using the Strichartz estimate implies that there is a small $\varepsilon_S > 0$ such that for any $T > 0$ and $\varphi \in \mathcal{H}$ satisfying

$$\vec{v}(t) := U(t)\varphi \implies \|v\|_{\text{St}_m(0,T)} \leq \varepsilon_S, \quad (1.24)$$

there is a unique solution u of (CW) on $[0, T)$ satisfying

$$\begin{aligned} \vec{u}(0) = \varphi, \quad \|\vec{u} - \vec{v}\|_{L^\infty \mathcal{H}(0,T)} + \|u - v\|_{\text{St}(0,T)} &\lesssim \|f'(u)\|_{\text{St}_s^*(0,T)} \\ &\lesssim \|u\|_{\text{St}_m(0,T)}^{2^*-2} \|u\|_{\text{St}_s(0,T)} \ll \|u\|_{\text{St}(0,T)} \sim \|v\|_{\text{St}(0,T)}, \end{aligned} \quad (1.25)$$

which is scattering to 0 if $T = \infty$. The uniqueness holds in

$$\{\vec{u} \in C([0, T]; \mathcal{H}) \mid \forall S \in (0, T), u \in \text{St}(0, S)\}, \quad (1.26)$$

and a solution u in this space can be extended beyond $T < \infty$ if and only if $\|u\|_{\text{St}_m(0, T)} < \infty$. For any interval $I \subset \mathbb{R}$, we denote by

$$\text{Solution}(I) \quad (1.27)$$

the set of all solutions u of (CW) on I such that $\vec{u} \in C(J; \mathcal{H}) \cap \text{St}(J)$ for any compact $J \subset I$, and that u can not be extended beyond any open boundary of I . For example, $\vec{u} \in \text{Solution}([0, T])$ with $T < \infty$ means that u is a solution in (1.26) with $\|u\|_{\text{St}(0, T)} = \infty$, so that it can be extended to $t < 0$ but not to $t > T$, whereas $\vec{u} \in \text{Solution}([0, T])$ means that u is a solution in $C([0, T]; \mathcal{H}) \cap \text{St}(0, T)$, which can be extended both to $t < 0$ and to $t > T$. Hence, if I is an open interval, then I is the maximal life for any $u \in \text{Solution}(I)$.

1.2 Symmetry, solitons and linearization

The groups of space translation and scaling are denoted by

$$\begin{aligned} T^c \varphi(x) &:= \varphi(x - c) \quad (c \in \mathbb{R}^d), \\ S^\sigma(\varphi_1, \varphi_2)(x) &:= (e^{\sigma(d/2-1)}\varphi_1(e^\sigma x), e^{\sigma d/2}\varphi_2(e^\sigma x)) \quad (\sigma \in \mathbb{R}), \end{aligned} \quad (1.28)$$

and for any $a \in \mathbb{R}$, $(S_a^\sigma \varphi)(x) := e^{\sigma(d/2+a)}\varphi(e^\sigma x)$. Their generators are $T' = -\nabla$, $S' = S'_{-1} \otimes S'_0$, and $S'_a = r\partial_r + d/2 + a$. For any $\vec{u} \in \text{Solution}(I)$ and $(\sigma, c) \in \mathbb{R}^{1+d}$,

$$T^c S^\sigma \vec{u}(e^\sigma t) \in \text{Solution}(e^{-\sigma} I). \quad (1.29)$$

The linearization around the ground state W is written by the operator

$$L_+ := -\Delta - f''(W), \quad f''(W) = (2^* - 1)(1 + |x|^2/(d(d-2)))^{-2}, \quad (1.30)$$

as well as the nonlinear term

$$N(v) := f'(W + v) - f'(W) - f''(W)v = O(v^2). \quad (1.31)$$

The matrix version of the linearization is given by $J\mathcal{L}$ with

$$\mathcal{L} := \begin{pmatrix} L_+ & 0 \\ 0 & 1 \end{pmatrix}, \quad J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (1.32)$$

The generator of each invariant transform of (CW) gives rise to generalized null vectors, namely with $A = S'_{-1}$ or $T' = -\nabla$,

$$J\mathcal{L} \begin{pmatrix} AW \\ 0 \end{pmatrix} = 0, \quad J\mathcal{L} \begin{pmatrix} 0 \\ AW \end{pmatrix} = \begin{pmatrix} AW \\ 0 \end{pmatrix}. \quad (1.33)$$

It is well known that for the ground state W , there is no other generalized null vector. Note however that $S'_{-1}W$ is not an eigenfunction but a threshold resonance, i.e. $S'_{-1}W \notin L^2$ for $d \leq 4$. Besides $L_+^{-1}(0) = \{\nabla W\}$ and the absolutely continuous spectrum $[0, \infty)$, L_+ has only one negative eigenvalue and the ground state,

$$L_+\rho = -k^2\rho, \quad 0 < \rho \in H^2(\mathbb{R}^d), \quad k > 0, \quad \|\rho\|_2 = 1, \quad (1.34)$$

for which the orthogonal subspace and projection are denoted by

$$\mathcal{H}_\perp := \{\varphi \in \mathcal{H} \mid \langle \varphi | \rho \rangle = 0 \in \mathbb{R}^2\}, \quad P_\perp := 1 - \rho \langle \rho |. \quad (1.35)$$

Henceforth, the L^2 inner product is denoted by

$$\langle \varphi | \psi \rangle := \operatorname{Re} \int_{\mathbb{R}^d} \varphi(x) \overline{\psi(x)} dx \in \mathbb{R}, \quad (1.36)$$

and for vector functions

$$\begin{aligned} \langle (\varphi_1, \varphi_2) | (\psi_1, \psi_2) \rangle &:= \langle \varphi_1 | \psi_1 \rangle + \langle \varphi_2 | \psi_2 \rangle \in \mathbb{R}, \\ \langle (\varphi_1, \varphi_2) | \psi \rangle &:= (\langle \varphi_1 | \psi \rangle, \langle \varphi_2 | \psi \rangle) =: \langle \psi | (\varphi_1, \varphi_2) \rangle \in \mathbb{R}^2, \end{aligned} \quad (1.37)$$

which may be applied to column vectors as well as higher dimensional vectors. Throughout the paper, a pair with a comma (\cdot, \cdot) denotes a vector, but never an inner product.

1.3 Coordinates around the ground states

We recall from [14] our dynamical coordinates for the solution \vec{u} around the ground states $S_{\text{static}}(W)$:

$$\vec{u}(t) = T^{c(t)} S^{\sigma(t)} (\vec{W} + v(t)), \quad v(t) = \lambda(t)\rho + \gamma(t), \quad \begin{cases} \lambda(t) = \langle v(t) | \rho \rangle, \\ \gamma(t) = P_\perp v(t), \end{cases} \quad (1.38)$$

where $v(t) = (v_1(t), v_2(t)) \in \mathcal{H}$ does not generally satisfy $v_2(t) = \dot{v}_1(t)$ because of the modulation $(\sigma(t), c(t))$. The unstable and stable modes are denoted by λ_\pm :

$$\lambda\rho = \lambda_+g_+ + \lambda_-g_-, \quad \lambda_\pm := \sqrt{\frac{k}{2}}\lambda_1 \pm \sqrt{\frac{1}{2k}}\lambda_2, \quad g_\pm := \frac{1}{\sqrt{2k}}(1, \pm k)\rho, \quad (1.39)$$

for which we introduce linear functionals $\Lambda_{\pm} : \mathcal{H} \rightarrow \mathbb{R}$ by

$$\Lambda_{\pm}\varphi = \frac{1}{\sqrt{2k}} \langle \varphi | (k, \pm 1) \rho \rangle = \langle J\varphi | \mp g_{\mp} \rangle, \tag{1.40}$$

so that we have

$$v = \Lambda_+(v)g_+ + \Lambda_-(v)g_- + P_{\perp}v. \tag{1.41}$$

If $\vec{u}(t)$ is close to $S_{\text{tatic}}(W)$, then we can uniquely choose $(\sigma(t), c(t))$ such that the orthogonality condition holds²

$$\mathbb{R}^{1+d} \ni (\alpha(t), \mu(t)) := \langle v_1(t) | (S'_0, T') \rho \rangle = 0 \tag{1.42}$$

by the implicit function theorem. Note that it is not preserved by the linearized equation, since neither $S'_0\rho$ nor $T'\rho$ is an eigenfunction of L_+ . The linearized energy norm E is defined on the entire \mathcal{H} by

$$\begin{aligned} \|v\|_E^2 &:= k^2\lambda_1^2 + \lambda_2^2 + \langle \mathcal{L}\gamma | \gamma \rangle + |\alpha|^2 + |\mu|^2 \sim \|v\|_{\mathcal{H}}^2 \\ &= |k\langle v_1 | \rho \rangle|^2 + |\langle v_2 | \rho \rangle|^2 + \langle \mathcal{L}P_{\perp}v | v \rangle + |\langle \gamma_1 | S'_0\rho \rangle|^2 + |\langle \gamma_1 | T'\rho \rangle|^2 \\ &= k|\lambda_+|^2 + k|\lambda_-|^2 + \langle \mathcal{L}P_{\perp}v | v \rangle + |\langle v_1 | S'_0\rho \rangle|^2 + |\langle v_1 | T'\rho \rangle|^2. \end{aligned} \tag{1.43}$$

See [14, Lemma2.1] for a proof of the equivalence to \mathcal{H} .

Note that all the above are static operations in \mathcal{H} defined around $S_{\text{tatic}}(W)$. More precisely, we define the bi-continuous affine maps $\Phi_{\sigma,c} : \mathbb{R}^2 \times \mathcal{H}_{\perp} \rightarrow \mathcal{H}$ and $\Psi_{\sigma,c} : \mathcal{H} \rightarrow \mathcal{H}$ for each $(\sigma, c) \in \mathbb{R}^{1+d}$ by

$$\Phi_{\sigma,c}(\lambda, \gamma) = \Psi_{\sigma,c}(\lambda\rho + \gamma), \quad \Psi_{\sigma,c}(v) = T^c S^{\vec{\sigma}}(\vec{W} + v). \tag{1.44}$$

For any $\delta > 0$, define open neighborhoods of 0 and $S_{\text{tatic}}(W)$ in \mathcal{H} , by

$$\mathcal{B}_{\delta} := \{v \in \mathcal{H} \mid \|v\|_{\mathcal{H}} < \delta\}, \quad \mathcal{N}_{\delta} := \bigcup_{(\sigma,c) \in \mathbb{R}^{1+d}} \Psi_{\sigma,c}(\mathcal{B}_{\delta}) \subset \mathcal{H}. \tag{1.45}$$

Then $\{(\Psi_{\sigma,c}(\mathcal{B}_{\delta}), \Psi_{\sigma,c}^{-1})\}_{(\sigma,c) \in \mathbb{R}^{1+d}}$ is an atlas for the open set $\mathcal{N}_{\delta} \subset \mathcal{H}$. Here is a precise statement on the orthogonality (1.42).

Lemma 1.4 *There exist $\delta_{\Phi} \in (0, 1)$ and a smooth map $(\vec{\sigma}, \vec{c}) : \mathcal{N}_{\delta_{\Phi}} \rightarrow \mathbb{R}^{1+d}$ such that for any $(\sigma, c) \in \mathbb{R}^{1+d}$ and $\varphi \in \Psi_{\sigma,c}(\mathcal{B}_{\delta_{\Phi}})$, we have $\langle \Psi_{\sigma,c}^{-1}(\varphi)_1 | (S'_0, T') \rho \rangle = 0$ if and only if $(\sigma, c) = (\vec{\sigma}(\varphi), \vec{c}(\varphi))$, and moreover*

$$|\vec{\sigma}(\varphi) - \sigma| + e^{\sigma} |\vec{c}(\varphi) - c| \lesssim \|P_{\perp} \Psi_{\sigma,c}^{-1}(\varphi)_1\|_{\dot{H}^1}. \tag{1.46}$$

² The sign of μ is switched from [14] for better notational symmetry.

Proof For any $\psi = \lambda\rho + \gamma \in \mathcal{B}_\delta$ and $(\sigma, c) \in \mathbb{R}^{1+d}$, define $(\alpha, \mu) : \mathbb{R}^{1+d} \rightarrow \mathbb{R}^{1+d}$ by

$$\begin{aligned} (\alpha, \mu)(s, y) &:= \langle \Psi_{s,y}^{-1}(\Psi_{\sigma,c}(\psi))_1 | (S'_0, T')\rho \rangle \\ &= \langle S_{-1}^{\sigma-s} T^{e^\sigma(c-y)}(W + \psi_1) - W | (S'_0, T')\rho \rangle, \end{aligned} \tag{1.47}$$

where we used the identity $T^c S_a^\sigma = S_a^\sigma T^{e^\sigma c}$. Hence we have

$$\begin{aligned} |(\alpha, \mu)(\sigma, c)| &= |\langle \gamma_1 | (S'_0, T')\rho \rangle| \lesssim \|\gamma_1\|_{\dot{H}^1} \lesssim \delta, \\ \partial_s(\alpha, \mu) &= \langle W + \psi_1 | T^{e^\sigma(y-c)} S_1^{s-\sigma} S'_1(S'_0, T')\rho \rangle \\ &= (-b_W, 0) + O(\delta + |s - \sigma| + e^\sigma|y - c|), \end{aligned} \tag{1.48}$$

where $b_W := \langle S'_{-1} W | S'_0 \rho \rangle = k^{-2}(2^* - 1)(2^* - 2)\langle W^{2^*-3}(S'_{-1} W)^2 | \rho \rangle > 0$ (see [14, (2.26)] for the identity), and

$$\begin{aligned} e^{-\sigma} \partial_y(\alpha, \mu) &= \langle W + \psi_1 | T' T^{e^\sigma(y-c)} S_1^{s-\sigma} (S'_0, T')\rho \rangle \\ &= -(0, a_W I) + O(\delta + |s - \sigma| + e^\sigma|y - c|), \end{aligned} \tag{1.49}$$

where $a_W := \langle -\Delta W | \rho \rangle / d = \langle f'(W) | \rho \rangle / d > 0$ and I denotes the identity matrix acting on \mathbb{R}^d . Then the implicit function theorem implies that there is a unique $(s, y) \in \mathbb{R}^{1+d}$ such that

$$(\alpha, \mu)(s, y) = 0, \quad |s - \sigma| + e^\sigma|y - c| \lesssim \|\gamma_1\|_{\dot{H}^1} \lesssim \delta, \tag{1.50}$$

provided that $\delta > 0$ is small enough. Since (α, μ) is obviously smooth in ψ , the implicit function is also smooth in $\Psi_{\sigma,c}(\mathcal{B}_\delta)$. For the uniqueness on \mathcal{N}_δ , suppose that $\Psi_{\sigma,c}(\psi) \in \Psi_{s,y}(\mathcal{B}_\delta)$ for some $(s, y) \in \mathbb{R}^{1+d}$, then

$$\delta \gtrsim \|\Psi_{\sigma,c}(0) - \Psi_{s,y}(0)\|_{\mathcal{H}} = \|(T^c S^\sigma - T^y S^s)\vec{W}\|_{\mathcal{H}} \sim |s - \sigma| + e^\sigma|y - c|. \tag{1.51}$$

Hence the uniqueness on \mathcal{N}_δ follows from the implicit function theorem. \square

For brevity, we define $\mathcal{T}_\varphi : \mathcal{H} \rightarrow \mathcal{H}$ for $\varphi \in \mathcal{N}_{\delta_\Phi}$, and $\tilde{\lambda} : \mathcal{N}_{\delta_\Phi} \rightarrow \mathbb{R}^2$ by

$$\mathcal{T}_\varphi := T^{\tilde{c}(\varphi)} S^{\tilde{\sigma}(\varphi)}, \quad \tilde{\lambda}(\varphi) := \langle \mathcal{T}_\varphi^{-1}(\varphi) - \vec{W} | \rho \rangle, \tag{1.52}$$

and similarly $\tilde{\lambda}_\pm : \mathcal{N}_{\delta_\Phi} \rightarrow \mathbb{R}$.

Remark 1.1 $\Phi_{\sigma,c}$ and $\Psi_{\sigma,c}$ are not smooth in (σ, c) for the γ component, since the derivative in (σ, c) induces $S'\gamma$ and $T'\gamma$, which are not generally in \mathcal{H} . Indeed $\Phi_{\sigma,c}$ is continuous for (σ, c) at each fixed point on \mathcal{H} , but not uniformly on any ball in \mathcal{H} . In [22] this was remedied by introducing a topology (“mobile distance”) in which translations are also Lipschitz continuous. Instead of that, we will fix (σ, c) with respect to perturbation of the initial data, even though we modulate it in time.

Next we change the time variable from t to τ by

$$\tau(0) = 0, \quad \frac{d\tau}{dt} = e^{\sigma t}. \tag{1.53}$$

Then we get the equation of v as an evolution in τ :

$$\begin{aligned} \partial_\tau v &= J\mathcal{L}v + \underline{N}(v_1) - Z(\vec{W} + v), \\ \underline{N}(\varphi) &:= (0, N(\varphi)), \quad Z = (Z_1, Z_2) := \sigma_\tau \mathcal{S}' + e^\sigma c_\tau \cdot \mathcal{T}', \end{aligned} \tag{1.54}$$

and differentiating the orthogonality condition (1.42) yields

$$\begin{aligned} 0 &= \partial_\tau \alpha = \langle \partial_\tau v_1 | S'_0 \rho \rangle = \langle \gamma_2 | S'_0 \rho \rangle + e^\sigma c_\tau \langle v_1 | \mathcal{T}' S'_0 \rho \rangle - \sigma_\tau [b_W - \langle v_1 | S'_1 S'_0 \rho \rangle], \\ 0 &= \partial_\tau \mu = \langle \partial_\tau v_1 | \mathcal{T}' \rho \rangle = \langle \gamma_2 | \mathcal{T}' \rho \rangle - e^\sigma c_\tau [a_W I - \langle v_1 | \nabla^2 \rho \rangle] + \sigma_\tau \langle v_1 | S'_1 \mathcal{T}' \rho \rangle. \end{aligned}$$

Hence, as long as v_1 is small,

$$\begin{aligned} \begin{pmatrix} \sigma_\tau \\ e^\sigma c_\tau \end{pmatrix} &= \begin{pmatrix} b_W - \langle v_1 | S'_1 S'_0 \rho \rangle & -\langle v_1 | \mathcal{T}' S'_0 \rho \rangle \\ -\langle v_1 | S'_1 \mathcal{T}' \rho \rangle & a_W I - \langle v_1 | \nabla^2 \rho \rangle \end{pmatrix}^{-1} \begin{pmatrix} \langle \gamma_2 | S'_0 \rho \rangle \\ \langle \gamma_2 | \mathcal{T}' \rho \rangle \end{pmatrix} \\ &= (1 + O(\|v_1\|_{\dot{H}^1})) \begin{pmatrix} b_W^{-1} \langle \gamma_2 | S'_0 \rho \rangle \\ a_W^{-1} \langle \gamma_2 | \mathcal{T}' \rho \rangle \end{pmatrix}. \end{aligned} \tag{1.55}$$

This is linear in v (or γ), because the orthogonality (1.42) is not preserved by the linearized equation, a notable difference from the standard modulation analysis in the subcritical case. In the original time t , it yields

$$(e^{-\sigma} \sigma_t, c_t) = (1 + O(\|v_1\|_{\dot{H}^1})) \langle \gamma_2 | (S'_0 \rho / b_W, \mathcal{T}' \rho / a_W) \rangle. \tag{1.56}$$

For the eigenmode we have

$$\begin{aligned} \partial_\tau \lambda &= \begin{pmatrix} 0 & 1 \\ k^2 & 0 \end{pmatrix} \lambda - \langle Zv | \rho \rangle + \langle \underline{N}(v_1) | \rho \rangle \\ &= \begin{pmatrix} \lambda_2 + \sigma_\tau (\alpha + \lambda_1) + e^\sigma c_\tau \mu \\ k^2 \lambda_1 + \langle N(v_1) - Z_2 \gamma_2 | \rho \rangle \end{pmatrix} = \begin{pmatrix} \lambda_2 + \sigma_\tau \lambda_1 \\ k^2 \lambda_1 + \langle N(v_1) - Z_2 \gamma_2 | \rho \rangle \end{pmatrix}, \end{aligned} \tag{1.57}$$

where we used $\alpha = 0 = \mu$ only in the last step, since we will consider the case $(\alpha, \mu) \neq 0$ as well. In the unstable/stable modes, the equation reads

$$\partial_\tau \lambda_\pm = \pm k \lambda_\pm + \sqrt{\frac{k}{2}} [\sigma_\tau (\alpha + \lambda_1) + e^\sigma c_\tau \mu] \pm \sqrt{\frac{1}{2k}} \langle N(v_1) - Z_2 \gamma_2 | \rho \rangle. \tag{1.58}$$

We also recall the distance function $d_W : \mathcal{H} \rightarrow [0, \infty)$ defined in [14], which satisfies $d_W(\varphi) \sim \text{dist}_W(\varphi)$, and, for some constant $\delta_E > 0$, if $d_W(\varphi) \leq \delta_E$ then $\pm\varphi \in \mathcal{N}_{\delta_\Phi}$ and

$$d_W(\varphi)^2 = E(\varphi) - E(W) + k^2 \tilde{\lambda}_1(\pm\varphi)^2 \tag{1.59}$$

for either sign \pm .

Part I: Slightly above the ground state energy

In the first part of paper, we study the global dynamics in the region $E(u) < E(W) + \varepsilon^2$, and its Lorentz extension, completing the picture in [14] with a center-stable manifold and the dynamics around it.

2 Center-stable manifold around the ground states

First we construct a center-stable manifold around the ground states $S_{\text{tatic}}(W)$. This will be later extended in three ways:

- (1) By the backward flow, to the region $E < E(W) + \varepsilon^2$,
- (2) By the Lorentz transform, to the region $E < \sqrt{E(W)^2 + \varepsilon^4 + |P(\bar{u})|^2}$
- (3) By adding large radiation, which may have arbitrarily large energy.

In order to define the manifold as a graph of $(\lambda_-, \gamma) \mapsto \lambda_+$, we define

$$\begin{aligned} \mathcal{B}_\delta^+ &:= \{\lambda_+ \in \mathbb{R} \mid \|\lambda_+ g_+ \|_{\mathcal{H}} < \delta\}, \\ \mathcal{B}'_\delta &:= \{\lambda_- g_- + \gamma \mid \lambda_- \in \mathbb{R}, \gamma \in \mathcal{H}_\perp, \|\lambda_- g_- + \gamma \|_{\mathcal{H}} < \delta\}, \end{aligned} \quad (2.1)$$

then $\mathcal{B}_{\delta/C} \subset \mathcal{B}_\delta^+ g_+ \oplus \mathcal{B}'_\delta \subset \mathcal{B}_{C\delta}$ for some constant $C > 1$. The corresponding neighborhood of $S_{\text{tatic}}(W)$ is denoted by

$$\mathcal{N}_{\delta_1, \delta_2} := \{\Psi_{\sigma, c}(\lambda_+ g_+ + \varphi) \mid (\sigma, c) \in \mathbb{R}^{1+d}, \lambda_+ \in \mathcal{B}_{\delta_1}^+, \varphi \in \mathcal{B}'_{\delta_2}\}. \quad (2.2)$$

Theorem 2.1 *There exist constants $\delta_m, \delta_X > 0$ satisfying $\delta_m \ll \delta_X \ll \delta_\Phi$ and $\delta_m \ll \varepsilon_S$, and a unique C^1 function $m_+ : \mathcal{B}'_{\delta_m} \rightarrow \mathcal{B}_{\delta_m}^+$ with the following property. Let $\lambda_+ \in \mathcal{B}_{\delta_m}^+$, $\varphi \in \mathcal{B}'_{\delta_m}$, $(\sigma, c) \in \mathbb{R}^{1+d}$, $T > 0$ and $\bar{u} \in S_{\text{solution}}([0, T])$ with $\bar{u}(0) = \Psi_{\sigma, c}(\lambda_+ g_+ + \varphi)$. Then we have the trichotomy:*

- (1) *If $\lambda_+ > m_+(\varphi)$, then u blows up away from the ground states. More precisely*

$$T < \infty, \quad \liminf_{t \nearrow T} d_W(\bar{u}(t)) > \delta_X \quad (2.3)$$

or $\liminf_{t \nearrow T} \text{dist}_{L^{2^}}(u(t), S_{\text{tatic}}(W)_1) \gg \delta_m$.*

- (2) *If $\lambda_+ = m_+(\varphi)$, then u is trapped by the ground states. More precisely, $d_W(\bar{u}(t))$ is decreasing until it reaches*

$$d_W(\bar{u}(t))^2 \leq 2(E(u) - E(W)) \leq 2d_W(\bar{u}(0))^2 \lesssim \|\varphi\|_{\mathcal{H}}^2, \quad (2.4)$$

and stays there for the rest of $t < T$.

- (3) *If $\lambda_+ < m_+(\varphi)$, then u scatters to 0. More precisely,*

$$T = \infty, \quad \exists \varphi_\infty \in \mathcal{H}, \quad \lim_{t \rightarrow \infty} \|\bar{u}(t) - U(t)\varphi_\infty\| = 0. \quad (2.5)$$

Moreover, in the cases (1) and (3), there exists $T_X \in (0, T)$ such that

$$0 < t < T_X \implies d_W(\tilde{u}(t)) < \delta_X, \quad T_X < t < T \implies d_W(\tilde{u}(t)) > \delta_X, \tag{2.6}$$

$$\tilde{\lambda}_+(\tilde{u}(T_X)) \sim \tilde{\lambda}_1(\tilde{u}(T_X)) \sim -K(u(T_X)) \sim \pm \delta_X,$$

with the sign + for (1) and - for (3). In addition, $m_+(0) = m'_+(0) = 0$ and

$$|m_+(\varphi^1) - m_+(\varphi^2)| \lesssim (\|\varphi^1\|_{\mathcal{H}} + \|\varphi^2\|_{\mathcal{H}})^{1/6} \|\varphi^1 - \varphi^2\|_{\mathcal{H}}. \tag{2.7}$$

Obviously, the three asymptotics in (1)–(3) are distinctive. From the preceding results around the ground states, we know that the case (2) contains type-II blowup and global solutions scattering to the ground states. Type-I blowup is contained in the case (1), but it may also contain type-II blowup. (2.6) comes from the one-pass theorem proved in [14].

Thus we obtain a manifold of codimension 1 in \mathcal{H} :

$$\mathcal{M}_0 := \{\Psi_{\sigma,c}(m_+(\varphi)g_+ + \varphi) \mid \varphi \in \mathcal{B}'_{\delta_m}, (\sigma, c) \in \mathbb{R}^{1+d}\}, \tag{2.8}$$

which contains $S_{\text{tatic}}(W)$ and is invariant by the forward flow within $\mathcal{N}_{\delta_m, \delta_m}$. It is also invariant by \mathcal{T} and \mathcal{S} by definition.

Then (2.7) implies that it is tangent to the center-stable subspace of the linearized evolution at each point on $S_{\text{tatic}}(W) = \{\Psi_{\sigma,c}(0)\}_{\sigma,c} \subset \mathcal{M}_0$, and that \mathcal{M}_0 is transverse to its time inversion \mathcal{M}_0^\dagger , since $\varphi \mapsto \varphi^\dagger$ exchanges λ_+ and λ_- . More explicitly

$$\mathcal{M}_0 \cap \mathcal{M}_0^\dagger = \bigcup_{(\sigma,c)} \Phi_{\sigma,c}\{(\lambda, \gamma) \mid (\lambda_\pm, \gamma) \in \mathcal{B}'_{\delta_m}, \lambda_\pm = m_+(\lambda_\mp g_+ + (\gamma_1, \pm \gamma_2))\} \tag{2.9}$$

is a local center manifold of codimension 2, on which every solution u satisfies

$$\delta_m \gtrsim d_W(\tilde{u}(t)) \gg |\tilde{\lambda}(\tilde{u}(t))| \tag{2.10}$$

all over its life, though it is not necessarily global. Obviously, \mathcal{M}_0^\dagger is relatively closed in \mathcal{M}_0 , splitting it into two non-empty, relatively open sets where $\lambda_- > m_+(\lambda_+ g_+ + \gamma^\dagger)$ or $\lambda_- < m_+(\lambda_+ g_+ + \gamma^\dagger)$. The solutions starting from the first set blow up away from the ground states in $t < 0$, while those starting from the second set scatters to 0 as $t \rightarrow -\infty$.

A C^1 functional $M_+ : \mathcal{N}_{\delta_m, \delta_m} \rightarrow \mathbb{R}$ is defined such that $\mathcal{M}_0 = M_+^{-1}(0)$, by putting

$$M_+(\varphi) = \lambda_+ - m_+(\lambda_- g_- + \gamma), \quad \varphi = \Phi_{\bar{\sigma}(\varphi), \bar{c}(\varphi)}(\lambda, \gamma). \tag{2.11}$$

It is clearly non-degenerate in the direction $\mathcal{T}_\varphi g_+$ by (2.7). Moreover, $M_+ > 0$, $M_+ = 0$ and $M_+ < 0$ respectively give the trichotomy (1)–(3).

The proof of the above theorem goes as follows. First we observe that if $\|\varphi\|_{\mathcal{H}} \ll |\lambda_+| \ll 1$ then we can apply the ejection lemma and the one-pass theorem from [14], and obtain (1) for $\lambda_+ \gg \|\varphi\|_{\mathcal{H}}$ and (3) for $-\lambda_+ \gg \|\varphi\|_{\mathcal{H}}$. Moreover, the ejection

lemma implies that every solution ejected at $t = T_X$ from a small neighborhood of $S_{\text{static}}(W)$ is categorized either in (1) with $\tilde{\lambda}_+(\tilde{u}(T_X)) > 0$, or in (3) with $\tilde{\lambda}_+(\tilde{u}(T_X)) < 0$. So each set of such initial data is open in \mathcal{H} . Hence there is at least one λ_+ in between, for which the solution is never ejected, i.e. the case (2). The uniqueness of such λ_+ follows also from the instability of W , or the exponential growth of the unstable component λ_+ for the difference of two solutions. The next section is devoted to its estimate, which is essentially the only ingredient in addition to [14].

As in [22], we abbreviate the differences by the following notation:

$$\langle X^\triangleright \rangle := X^1 - X^0, \quad \langle F(X^\triangleright) \rangle := F(X^1) - F(X^0), \quad (2.12)$$

for any symbol X and any function F .

2.1 Igniting the unstable mode

In this subsection, we prove the following: For any solution trapped by the ground states, an arbitrarily small perturbation leads to the ejection from the small neighborhood unless the perturbation is almost zero in the unstable direction. More precisely,

Lemma 2.2 (Ignition lemma) *There exist constants $0 < \iota_I < 1 < C_I < \infty$ with the following property. Let $T > 0 < \iota \leq \iota_I$, $\tilde{u}^0 \in S_{\text{solution}}([0, T])$, $(s, y) \in \mathbb{R}^{1+d}$ and $\varphi \in \mathcal{H}$ satisfy $\tilde{u}^0(0) \in \Psi_{s,y}(\mathcal{B}_\iota)$,*

$$\|d_W(\tilde{u}^0)\|_{L_t^\infty(0,T)} < \iota^3, \quad C_I \iota \|\varphi\|_{\mathcal{H}} < |\Lambda_+ \varphi|, \quad \|\varphi\|_{\mathcal{H}} < \iota^6. \quad (2.13)$$

Then there exist $t_I \in (0, T)$, $\lambda_+ \in \mathbb{R}$, and $\tilde{u}^1 \in S_{\text{solution}}([0, t_I])$ such that

$$\begin{aligned} \tilde{u}^1(0) &= \tilde{u}^0(0) + \mathcal{T}^y \mathcal{S}^s \varphi, \\ \|\tilde{u}^0 - \tilde{u}^1\|_{L^\infty(0,t_I;\mathcal{H})} &= \|\tilde{u}^0(t_I) - \tilde{u}^1(t_I)\|_{\mathcal{H}} = \iota^3 \sim \lambda_+ \text{sign}(\Lambda_+ \varphi), \\ \|\tilde{u}^0(t_I) - \tilde{u}^1(t_I) - \mathcal{F}_{\tilde{u}^0(t_I)} \lambda_+ g_+\|_{\mathcal{H}} &\lesssim \iota^4. \end{aligned} \quad (2.14)$$

In particular, we have

$$d_W(\tilde{u}^0(t_I)) + d_W(\tilde{u}^1(t_I)) \sim \iota^3. \quad (2.15)$$

This lemma is proved by exponential growth in the unstable direction of the difference $u^0 - u^1$ in the rescaled coordinate for u^0 . It may take very long depending on the initial size of the perturbation, but in the rescaled time τ , where the solution u^0 is (forward) global in both the scattering and the blow-up cases. The difference is estimated mainly by the energy argument, rather than dispersive estimates. The nonlinearity is too strong to be controlled solely by Sobolev, for which we employ Strichartz norms which are uniform on unit intervals of τ .

Hence the main idea is similar to [22], but we do not use the mobile distance, but instead the same modulation parameters $(\sigma(t), c(t))$ for both u^0 and u^1 , in order to avoid destroying the energy structure for the difference. This is indeed much simpler,

whereas the former idea seems hard to apply in the critical setting because of the change of time variable.

The main difference from the ejection lemma in [14] is that there is no bound on the time for the unstable mode to grow to some amount, and we estimate the difference of two solutions rather than the difference from the ground state. In particular, the equation for the difference naturally contains linear terms coming from the nonlinearity, which prevents us from a crude Duhamel argument as in [14].

Before starting the proof, we see that the Strichartz norms can be uniformly bounded on unit time intervals in the rescaled variables:

Lemma 2.3 (Locally uniform perturbation in St_τ) *There is a constant $\eta_l \in (0, 1]$ with the following property. Let $T > 0 < \delta \leq \eta_l$, $\vec{u} \in \text{S}_{\text{olution}}([0, T])$,*

$$\begin{aligned} \vec{u}(t) &= \Psi_{\sigma(t), c(t)} v(\tau(t)), \quad \frac{d\tau}{dt}(t) = e^{\sigma(t)}, \quad \tau(0) = 0, \\ \|v(0)\|_{\mathcal{H}} + |e^{-\sigma(t)} \sigma'(t)| + |c'(t)| &\leq \delta, \end{aligned} \tag{2.16}$$

for $0 < t < T$. Then we have $\tau(T) > \eta_l$ and

$$\|v\|_{\text{St} \cap L^\infty \mathcal{H}(0, \eta_l)} \lesssim \delta. \tag{2.17}$$

Moreover, if $\text{S}_{\text{olution}}([0, T^1]) \ni \vec{u}^1(t) = \Psi_{\sigma(t), c(t)} v^1(\tau(t))$ satisfies $\|\vec{u}^1(0) - \vec{u}(0)\|_{\mathcal{H}} < \eta_l$, then we have $\tau(T^1) > \eta_l$ and

$$\|v^1 - v\|_{\text{St} \cap L^\infty \mathcal{H}(0, \eta_l)} \lesssim \|\vec{u}^1(0) - \vec{u}(0)\|_{\mathcal{H}}. \tag{2.18}$$

Proof We obtain from the inequality on (σ', c') that

$$|e^{\sigma(t)} - e^{\sigma(0)}| < e^{\sigma(t)}/4 < e^{\sigma(0)}/2, \quad |\sigma(t) - \sigma(0)| + e^{\sigma(t)}|c(t) - c(0)| < \delta \tag{2.19}$$

for $0 < t < e^{-\sigma(0)}/4$. Let $\vec{W}^0 := \mathcal{T}^{c(0)} \mathcal{S}^{\sigma(0)} \vec{W}$ and $\vec{w}(t) := \vec{u}(t) - \vec{W}^0$. Then we have

$$\|W^0\|_{\text{St}(0, e^{-\sigma(0)\eta})} = \|W\|_{\text{St}(0, \eta)} \lesssim \eta^{1/q_m} \tag{2.20}$$

for $0 < \eta < 1$, and w solves on $(0, T)$

$$(\partial_t^2 - \Delta)w = f'(W^0 + w) - f'(W^0). \tag{2.21}$$

Hence by Strichartz we have for small $\eta > 0$

$$\begin{aligned} \|\vec{w} - \vec{w}_F\|_{\text{St} \cap L^\infty \mathcal{H}(0, e^{-\sigma(0)\eta})} &\ll \|w\|_{\text{St}(0, e^{-\sigma(0)\eta})} \\ &\sim \|w_F\|_{\text{St}(0, e^{-\sigma(0)\eta})} \lesssim \|\vec{w}(0)\|_{\mathcal{H}} = \|v(0)\|_{\mathcal{H}}, \end{aligned} \tag{2.22}$$

where $\vec{w}_F := U(t)\vec{w}(0) = U(t)v(0)$. In particular there is some $\eta_l < 1/8$ such that the above estimates hold for $\eta \leq 2\eta_l$, and so $\tau(T) > \tau(2e^{-\sigma(0)\eta_l}) > \eta_l$. Under the scaling property

$$1/p + d/q - s = d/2 - 1, \tag{2.23}$$

we have, for any $T' \in (0, T)$,

$$\begin{aligned} \|v_1\|_{L^p_{\tau}(0, \tau(T'); \dot{B}^s_{q,2})} &= \int_0^{T'} \|v_1(\tau(t))\|_{\dot{B}^s_{q,2}}^p e^{\sigma(t)} dt \\ &= \int_0^{T'} \|\mathcal{T}^{c(t)} \mathcal{S}^{\sigma(t)} v_1(\tau(t))\|_{\dot{B}^s_{q,2}}^p dt \\ &= \|u_1 - \mathcal{T}^{c(t)} \mathcal{S}^{\sigma(t)} W\|_{L^p_{t'}(0, T'; \dot{B}^s_{q,2})}^p, \end{aligned} \tag{2.24}$$

and similarly for v_2 and in $L^\infty \mathcal{H}$. Hence putting $T' = \tau^{-1}(\eta_l) < 2e^{-\sigma(0)} \eta_l$,

$$\|v\|_{\text{St} \cap L^\infty \mathcal{H}(0, \eta_l)} \leq \|\vec{w}\|_{\text{St} \cap L^\infty \mathcal{H}(0, T')} + \|(\mathcal{T}^c \mathcal{S}^\sigma - \mathcal{T}^{c(0)} \mathcal{S}^{\sigma(0)}) \vec{W}\|_{\text{St} \cap L^\infty \mathcal{H}(0, T')}, \tag{2.25}$$

and the last norm is bounded by

$$\begin{aligned} &\|(\mathcal{T}^{c-c(0)} - I) \mathcal{S}^\sigma \vec{W}\|_{\text{St} \cap L^\infty \mathcal{H}(0, T')} + \|(\mathcal{S}^{\sigma-\sigma(0)} - I) \mathcal{S}^{\sigma(0)} \vec{W}\|_{\text{St} \cap L^\infty \mathcal{H}(0, T')} \\ &\lesssim \|e^\sigma |c - c(0)| + |\sigma - \sigma(0)|\|_{L^\infty(0, T')} \lesssim \delta, \end{aligned} \tag{2.26}$$

which concludes the estimate on v .

For the difference, the same change of variable as in (2.24) yields for $T' \in (0, T)$

$$\|v^1 - v\|_{\text{St} \cap L^\infty \mathcal{H}(0, \tau(T'))} = \|\vec{u}^1 - \vec{u}\|_{\text{St} \cap L^\infty \mathcal{H}(0, T')}. \tag{2.27}$$

Since $\|u\|_{\text{St}(0, \tau^{-1}(\eta_l))} + \|\vec{u}(0) - \vec{u}^1(0)\|_{\mathcal{H}} \ll 1$, we have $T^1 > \tau^{-1}(\eta_l)$ and

$$\|\vec{u}^1 - \vec{u}\|_{\text{St} \cap L^\infty \mathcal{H}(0, \tau^{-1}(\eta_l))} \lesssim \|\vec{u}^1(0) - \vec{u}(0)\|_{\mathcal{H}}, \tag{2.28}$$

which leads to the estimate on $v^1 - v$. □

Proof of Lemma 2.2 Let $(\sigma, c) : [0, T] \rightarrow \mathbb{R}^{1+d}$ be the modulation for u^0 defined by

$$(\sigma(t), c(t)) = (\tilde{\sigma}(\vec{u}^0(t)), \tilde{c}(\vec{u}^0(t))), \tag{2.29}$$

and let $\tau : [0, T] \rightarrow (0, \infty)$ be the rescaled time variable for u^0 defined by

$$\frac{d\tau}{dt} = e^{\sigma(t)}, \quad \tau(0) = 0. \tag{2.30}$$

Let $\vec{u}^1 \in \text{S}_{\text{olution}}([0, T^1])$ be the solution with the initial data

$$\vec{u}^1(0) = \vec{u}^0(0) + \mathcal{T}^y \mathcal{S}^s \varphi. \tag{2.31}$$

We use the same coordinates for u^0 and u^1 by putting for $j = 0, 1$ and at each t

$$\bar{u}^j = \Psi_{\sigma,c}(v^j) = \Phi_{\sigma,c}(\lambda^j, \gamma^j). \tag{2.32}$$

For the initial perturbation, we have, using (1.46),

$$\begin{aligned} |\langle \lambda^\triangleright(0) - \langle \varphi | \rho \rangle| &= | \langle (\mathcal{S}^{s-\sigma(0)} \mathcal{T}^{e^s(y-c(0))} - I) \varphi | \rho \rangle | \\ &\lesssim (|s - \sigma(0)| + e^s |y - c(0)|) \|\varphi\|_{\mathcal{H}} \\ &\lesssim \|(\Psi_{s,y})^{-1}(\bar{u}^0(0))\|_{\mathcal{H}} \|\varphi\|_{\mathcal{H}} \lesssim \iota \| \langle v^\triangleright(0) \|_{\mathcal{H}}, \end{aligned} \tag{2.33}$$

which, together with (2.13), implies

$$C_I \iota \| \langle v^\triangleright(0) \|_{\mathcal{H}} < 2 | \langle \lambda^\triangleright_+(0) |, \tag{2.34}$$

if C_I is large and ι_I is small enough.

Since the modulation (σ, c) was chosen for u^0 , we have

$$\|v^0\|_{L^\infty(0,T;\mathcal{H})} \sim \sup_{0 < t < T} d_W(\bar{u}^0) =: \delta < \iota^3. \tag{2.35}$$

For the Strichartz norms, Lemma 2.3 implies

$$\|v^0\|_{\text{St}(\tau_0 < \tau < \tau_0 + \eta)} \lesssim \delta, \quad \| \langle v^\triangleright \|_{\text{St}(\tau_0 < \tau < \tau_0 + \eta)} \lesssim \| \langle v^\triangleright(\tau_0) \|_{\mathcal{H}}, \tag{2.36}$$

as long as $\| \langle v^\triangleright \|_{\mathcal{H}}$ remains small, while (1.55) implies

$$|\tau_0 - \tau_1| \lesssim \delta^{-1} \implies |\sigma(\tau_0) - \sigma(\tau_1)| + |e^{\sigma(\tau_0)}[c(\tau_0) - c(\tau_1)]| \lesssim \delta |\tau_0 - \tau_1|. \tag{2.37}$$

In particular, we have $\tau(t) \nearrow \infty$ as $t \nearrow T$, since otherwise $|\sigma|$ is bounded as $t \nearrow T$, and so, if $T < \infty$ then $\|u^0\|_{\text{St}(0,T)} < \infty$ contradicting the blowup at T , and if $T = \infty$ then the boundedness of $\dot{\tau} = e^\sigma$ implies that $\tau \rightarrow \infty$.

In the following, we regard all the dynamical variables as functions of τ rather than t , unless explicitly specified. We have the equations for the difference

$$\begin{aligned} \partial_\tau \langle \lambda^\triangleright_1 &= \langle \lambda^\triangleright_2 + \sigma_\tau (\alpha^1 + \langle \lambda^\triangleright_1) + e^\sigma c_\tau \cdot \mu^1, \\ \partial_\tau \langle \lambda^\triangleright_2 &= k^2 \langle \lambda^\triangleright_1 + \langle N(v^\triangleright_1) - Z_2 \gamma^\triangleright_2 | \rho \rangle, \\ \partial_\tau (\alpha^1, \mu^1) &= \langle \gamma^\triangleright_2 | (S'_0, T') \rho \rangle - \langle Z_1 \langle v^\triangleright_1 | (S'_0, T') \rho \rangle, \\ \partial_\tau \langle \gamma^\triangleright &= J \mathcal{L} \langle \gamma^\triangleright + P_\perp [\langle N(v^\triangleright_1) - Z \langle v^\triangleright], \end{aligned} \tag{2.38}$$

Also remember that $(\alpha^0, \mu^0) = \langle v^0 | (S'_0, T') \rho \rangle = 0$ and so

$$(\alpha^1, \mu^1) = \langle \alpha^\triangleright, \mu^\triangleright \rangle = \langle \gamma^\triangleright_1 | (S'_0, T') \rho \rangle. \tag{2.39}$$

Hence the third equation follows from the fourth one in (2.38). By the assumption,

$$\delta < \iota^3, \quad \|\langle v^\triangleright(0) \rangle_{\mathcal{H}}\| < \iota^6. \quad (2.40)$$

Suppose that for some $\tau_0 > 0$ we have

$$\|\langle v^\triangleright \|_{L^\infty(0, \tau_0; \mathcal{H})} < \iota^3, \quad \iota^2 |\langle \lambda_-^\triangleright(\tau_0) \rangle| + \iota v(\tau_0) < |\langle \lambda_+^\triangleright(\tau_0) \rangle|, \quad (2.41)$$

where we put

$$v(\tau) := \sqrt{\iota^2 |(\alpha^1, \mu^1)|^2 + \langle \mathcal{L}\gamma | \gamma \rangle}. \quad (2.42)$$

Choosing $C_I > 1$ large enough, we have (2.41) at $\tau_0 = 0$. We will prove that the second condition of (2.41) is preserved until the first one is broken.

The linearized energy in (1.43) implies that at each time

$$\|\langle \gamma^\triangleright \|_{\mathcal{H}} \gtrsim v \gtrsim \iota \|\langle \gamma_1^\triangleright \|_{\dot{H}^1} + \|\langle \gamma_2^\triangleright \|_{L^2} + \iota |(\alpha^1, \mu^1)|. \quad (2.43)$$

Lemma 2.3 implies that u^1 exists at least for $\tau < \tau_0 + \eta_l$ and

$$\|v^0\|_{\text{St} \cap L^\infty \mathcal{H}(\tau_0, \tau_0 + \eta_l)} \lesssim \delta, \quad \|\langle v^\triangleright \|_{\text{St} \cap L^\infty \mathcal{H}(\tau_0, \tau_0 + \eta_l)} \lesssim \|\langle v^\triangleright(\tau_0) \rangle_{\mathcal{H}} < \iota^3. \quad (2.44)$$

Using (1.55) as well, we derive from the difference equations

$$\begin{aligned} |(\partial_\tau \mp k) \langle \lambda_\pm^\triangleright | &\lesssim \delta \|\langle v^\triangleright \|_{\mathcal{H}} + \delta |(\alpha^1, \mu^1)| + \|\langle v^\triangleright \|_{\mathcal{H}}^2 \lesssim \iota^3 |\langle \lambda^\triangleright | + \iota^2 v, \\ |\partial_\tau (\alpha^1, \mu^1)| &\lesssim \|\langle \gamma_2^\triangleright \|_{L^2} + \delta \|\langle v^\triangleright \|_{\mathcal{H}}. \end{aligned} \quad (2.45)$$

To control $\langle \gamma^\triangleright$, we use the linearized energy identity

$$\begin{aligned} \partial_\tau \langle \mathcal{L} \langle \gamma^\triangleright | \langle \gamma^\triangleright \rangle / 2 &= \langle \langle N(v_1^\triangleright) - Z \langle v^\triangleright | \mathcal{L} \langle \gamma^\triangleright \rangle \rangle \\ &= \langle \langle N(v_1^\triangleright) | \langle \gamma_2^\triangleright \rangle \rangle + e^\sigma c_\tau [\langle L_+ \nabla \rho | \langle \gamma_1^\triangleright \rangle \rangle \langle \lambda_1^\triangleright \rangle + \langle \nabla \mathcal{W} | \langle \gamma_1^\triangleright \rangle^2 / 2 \rangle + \langle \nabla \rho | \langle \gamma_2^\triangleright \rangle \rangle \langle \lambda_2^\triangleright \rangle] \\ &\quad - \sigma_\tau [\langle L_+ S'_{-1} \rho | \langle \gamma_1^\triangleright \rangle \rangle \langle \lambda_1^\triangleright \rangle + \langle S'_{2-d/2} \mathcal{W} | \langle \gamma_1^\triangleright \rangle^2 / 2 \rangle + \langle S'_0 \rho | \langle \gamma_2^\triangleright \rangle \rangle \langle \lambda_2^\triangleright \rangle], \end{aligned} \quad (2.46)$$

where $\mathcal{W} := f''(W)$. The terms on the right except for the first one are simply bounded by $\delta \|\langle \gamma^\triangleright \|_{\mathcal{H}} \|\langle v^\triangleright \|_{\mathcal{H}}$. The nonlinear term can be bounded only via τ -integral. For any interval $I = (\tau_0, \tau_1) \subset [0, \infty)$ with $|I| < \eta_l$, we have,

$$\int_I | \langle \langle N(v_1^\triangleright) | \langle \gamma_2^\triangleright \rangle \rangle | d\tau \lesssim \|\langle \gamma_2^\triangleright \|_{L^\infty_\tau(I; L^2_x)} \|\langle N(v_1^\triangleright) \|_{L^1_\tau(I; L^2_x)}. \quad (2.47)$$

Since

$$\langle N(v_1^\triangleright) \rangle = \int_0^1 [f''(W + v_1^\theta) - f''(W)] \langle v_1^\triangleright d\theta, \quad v^\theta := (1 - \theta)v^0 + \theta v^1, \quad (2.48)$$

we have³

$$\begin{aligned} \|\langle N(v_1^\triangleright) \rangle\|_{L^1 L^2(I)} &\lesssim \sup_{0 < \theta < 1} \| [f''(W + v_1^\theta) - f''(W)] \langle v_1^\triangleright \rangle \|_{L^1 L^2_x(I)} \\ &\lesssim (\|v_1^0\|_{\text{St}_p(I)} + \|\langle v_1^\triangleright \rangle\|_{\text{St}_p(I)}) \|\langle v_1^\triangleright \rangle\|_{\text{St}_p(I)} \\ &\lesssim (\delta + \|\langle v^\triangleright(\tau_0) \rangle\|_{\mathcal{H}}) \|\langle v^\triangleright(\tau_0) \rangle\|_{\mathcal{H}}, \end{aligned} \tag{2.49}$$

where we used Lemma 2.3 in the last step. We thus obtain

$$|[(\mathcal{L}\langle \gamma^\triangleright | \langle \gamma^\triangleright \rangle)]_{\tau_0}^{\tau_1}]| \lesssim \iota^3 \|\langle \gamma^\triangleright \rangle\|_{L^\infty(\mathcal{H}_\tau)} \|\langle v^\triangleright(\tau_0) \rangle\|_{\mathcal{H}}. \tag{2.50}$$

Combining this with the estimate on (α^1, μ^1) yields

$$\begin{aligned} [v^2]_{\tau_0}^{\tau_1} &\lesssim \iota \nu [\|\langle \gamma^\triangleright \rangle\|_{L^2_x} + \delta \|\langle v^\triangleright(\tau_0) \rangle\|_{\mathcal{H}}] + \iota^3 \|\langle \gamma^\triangleright \rangle\|_{L^\infty \mathcal{H}} \|\langle v^\triangleright(\tau_0) \rangle\|_{\mathcal{H}} \\ &\lesssim \iota \|\nu\|_{L^\infty_\tau}^2 + \iota^2 \|\nu\|_{L^\infty_\tau} \|\langle v^\triangleright(\tau_0) \rangle\|_{\mathcal{H}} \lesssim \iota \|\nu\|_{L^\infty_\tau}^2 + \iota^3 \|\langle \lambda^\triangleright \rangle\|_{L^\infty_\tau}^2. \end{aligned} \tag{2.51}$$

Thus we obtain for $\tau_0 < \tau < \tau_0 + \eta_l$,

$$\begin{aligned} v &\leq (1 + C\iota)v(\tau_0) + C\iota^{3/2} \|\langle \lambda^\triangleright \rangle\|_{L^\infty_\tau}, \\ |(\partial_\tau \mp k)\langle \lambda^\triangleright_\pm \rangle| &\lesssim \iota^3 \|\langle \lambda^\triangleright \rangle\| + \iota^2 v. \end{aligned} \tag{2.52}$$

Suppose that $\|\langle \lambda^\triangleright \rangle\|_{L^\infty_\tau} \leq M \|\langle \lambda^\triangleright(\tau_0) \rangle\|$. Then using that $\iota v(\tau_0) < \|\langle \lambda^\triangleright_+(\tau_0) \rangle\|$,

$$|(\partial_\tau \mp k)\langle \lambda^\triangleright_\pm \rangle| \lesssim [\iota^3 M + \iota(1 + \iota^{3/2} M)] \|\langle \lambda^\triangleright(\tau_0) \rangle\| \lesssim \iota \|\langle \lambda^\triangleright(\tau_0) \rangle\|, \tag{2.53}$$

provided that $\iota^{3/2} M \ll 1$. Hence by continuity in τ , we deduce that

$$|\langle \lambda^\triangleright \rangle| \lesssim \|\langle \lambda^\triangleright(\tau_0) \rangle\|, \tag{2.54}$$

for $\tau_0 < \tau < \tau_0 + \eta_l$, and plugging this into (2.52),

$$\begin{aligned} v &\leq (1 + C\iota)v(\tau_0) + C\iota^{3/2} \|\langle \lambda^\triangleright(\tau_0) \rangle\|, \\ |\langle \lambda^\triangleright_\pm \rangle - e^{\pm k(\tau - \tau_0)} \langle \lambda^\triangleright_\pm(\tau_0) \rangle| &\leq C\iota \|\langle \lambda^\triangleright_+(\tau_0) \rangle\|. \end{aligned} \tag{2.55}$$

In particular, if $0 < \iota \ll 1$ then

$$\begin{aligned} |[\langle \lambda^\triangleright_+ \rangle]_{\tau_0}^{\tau_0 + \eta_l}| &> (e^{(k - C\iota)\eta_l} - 1) \|\langle \lambda^\triangleright_+(\tau_0) \rangle\| \\ &\gg \iota \|\langle \lambda^\triangleright_+(\tau_0) \rangle\| \gtrsim [\iota^2 \|\langle \lambda^\triangleright_-(\tau) \rangle\| + \iota \nu]_{\tau_0}^{\tau_0 + \eta_l}. \end{aligned} \tag{2.56}$$

³ Here for simplicity we use the exponents available only for $d \leq 6$, but it is clear that we only need Hölder continuity of f'' , i.e. $2^* > 2$, and so it can be easily modified for all dimensions $d \geq 3$.

Hence the last condition of (2.41) is transferred to $\tau = \tau_0 + \eta_l$. Therefore by iteration, (2.41) and the above estimates hold with $\tau_0 = n\eta_l$ for integers $0 \leq n < N$, where either $N = \infty$ or $2 \leq N < \infty$ and $\|v\|_{L^\infty_\tau(N\eta_l, (N+1)\eta_l; \mathcal{H})} \geq \iota^3$.

We can improve the above estimates as follows. First from the second estimate for λ_+ in (2.55), we have for all $0 \leq n < N$,

$$e^{(k-C\iota)\eta_l} < \langle \lambda_+^\triangleright((n+1)\eta_l) \rangle / \langle \lambda_+^\triangleright(n\eta_l) \rangle < e^{(k+C\iota)\eta_l}, \tag{2.57}$$

which precludes the case $N = \infty$. Plugging this exponential growth in the same estimate for λ_- , we obtain for $0 \leq n \leq N$

$$|\langle \lambda_-^\triangleright(n\eta_l) \rangle| \leq e^{-(k-C\iota)n\eta_l} |\langle \lambda_-^\triangleright(0) \rangle| + C\iota |\langle \lambda_+^\triangleright(n\eta_l) \rangle|. \tag{2.58}$$

Using those two in the first estimate of (2.55), we obtain for $0 \leq n \leq N$

$$v(n\eta_l) \leq e^{C\iota n\eta_l} v(0) + C\iota^{3/2} [|\langle \lambda_-^\triangleright(0) \rangle| + |\langle \lambda_+^\triangleright(n\eta_l) \rangle|]. \tag{2.59}$$

Iterating once again, we obtain continuous versions for $0 < \tau < N\eta_l$

$$\begin{aligned} e^{(k-C\iota)\tau} &\lesssim \langle \lambda_+^\triangleright(\tau) \rangle / \langle \lambda_+^\triangleright(0) \rangle \lesssim e^{(k+C\iota)\tau}, \\ |\langle \lambda_-^\triangleright(\tau) \rangle| &\lesssim e^{-(k-C\iota)\tau} |\langle \lambda_-^\triangleright(0) \rangle| + \iota |\langle \lambda_+^\triangleright(\tau) \rangle|, \\ \|v\|_{L^\infty(0, \tau)} &\lesssim e^{C\iota\tau} v(0) + \iota^{3/2} [|\langle \lambda_-^\triangleright(0) \rangle| + |\langle \lambda_+^\triangleright(\tau) \rangle|]. \end{aligned} \tag{2.60}$$

In particular, we have

$$\begin{aligned} \|\langle v^\triangleright \|_{L^\infty(0, \tau; \mathcal{H})} &\lesssim \|\langle \lambda^\triangleright \|_{L^\infty(0, \tau)} + \iota^{-1} \|v\|_{L^\infty(0, \tau)} \\ &\lesssim |\langle \lambda_+^\triangleright(\tau) \rangle| + \iota^{-1} e^{C\iota\tau} \|\langle v^\triangleright(0) \rangle\|_{\mathcal{H}}. \end{aligned} \tag{2.61}$$

Since $\|\langle v^\triangleright(0) \rangle\|_{\mathcal{H}} < \iota^6$, the last term is bounded by ι^4 for $e^{C\iota\tau} < \iota^{-1}$, while

$$|\langle \lambda_+^\triangleright(\tau) \rangle| \gtrsim \left[e^{C\iota\tau} \right]^{(k-C\iota)/C\iota} \iota^2 \|\langle v^\triangleright(0) \rangle\|_{\mathcal{H}} \gg \iota^{-10} e^{C\iota\tau} \|\langle v^\triangleright(0) \rangle\|_{\mathcal{H}}, \tag{2.62}$$

for $e^{C\iota\tau} \geq \iota^{-1}$, choosing $\iota_I \ll k$. Hence for some $\tau_I \in (N\eta_l, (N+1)\eta_l)$ we have

$$\begin{aligned} \iota^3 &= \|\langle v^\triangleright(\tau_I) \rangle\|_{\mathcal{H}} = \|\langle v^\triangleright \|_{L^\infty(0, \tau_I; \mathcal{H})} \sim |\langle \lambda_+^\triangleright(\tau_I) \rangle| \sim |\langle \lambda_+^\triangleright(\tau_I/2) \rangle|, \\ \iota^4 &\gtrsim \|\langle \lambda_-^\triangleright \|_{L^\infty(0, \tau_I)} + \|v/\iota\|_{L^\infty(0, \tau_I)} \\ &\gtrsim \|\langle \lambda_-^\triangleright \|_{L^\infty(0, \tau_I)} + \|\langle \gamma^\triangleright \|_{L^\infty(0, \tau_I; \mathcal{H})} + \|(\alpha^1, \mu^1)\|_{L^\infty(0, \tau_I)}. \end{aligned} \tag{2.63}$$

$\tau_I < \infty$ means that $t_I := \tau^{-1}(\tau_I) \in (0, T)$ and $\vec{u}^1 \in \text{Solution}(\{0, t_I\})$. For the last statement of the lemma, suppose $d_W(\vec{u}^0(t_I)) \ll \iota^3$. Then $\|\gamma^1(t_I)\|_{\mathcal{H}} \lesssim \iota^4 + d_W(\vec{u}^0(t_I)) \ll \iota^3$ and via (1.46) we have $|\langle \tilde{\sigma}(\vec{u}^\triangleright(t_I)) \rangle| + e^{\sigma(t_I)} |\langle \tilde{c}(\vec{u}^\triangleright(t_I)) \rangle| \lesssim \|\gamma^1(t_I)\|_{\mathcal{H}} \ll \iota^3$, and so

$$d_W(\vec{u}^1(t_I)) \gtrsim \|\vec{v}^1(t_I)\|_{\mathcal{H}} - C \|\mathcal{F}_{\vec{u}^0(t_I)} \vec{W} - \mathcal{F}_{\vec{u}^1(t_I)} \vec{W}\|_{\mathcal{H}} \gtrsim \iota^3. \tag{2.64}$$

Therefore $d_W(\vec{u}^0(t_I)) + d_W(\vec{u}^1(t_I)) \sim \iota^3$ in any case (the upper bound is obvious). □

2.2 Construction of the manifold

Now we are ready to prove Theorem 2.1. Let $0 < \delta' \ll \delta_+ \ll \delta_\Phi, \lambda_+ \in \mathcal{B}_{\delta_+}^+, \varphi \in \mathcal{B}'_{\delta'}$ and $\vec{u} \in \text{Solution}([0, T])$ with $\vec{u}(0) = \lambda_+ g_+ + \varphi$. Choosing δ_+ small enough ensures that

$$E(u) < E(W) + \varepsilon_*^2, \quad \delta_+ \ll \delta_*, \tag{2.65}$$

where $\delta_* \gg \varepsilon_* > 0$ are the small constants in the one-pass theorem [14, Theorem 5.1]. For each fixed φ , we divide the set $\mathcal{B}_{\delta_+}^+$ for λ_+ according to the behavior of \vec{u} . Let A_\pm be the totality of $\lambda_+ \in \mathcal{B}_{\delta_+}^+$ for which there exists $t_0 \in [0, T)$ such that

$$\begin{aligned} \delta_*^2 > d_W(\vec{u}(t_0))^2 > 2(E(u) - E(W)), \\ \partial_t d_W(\vec{u}(t_0)) > 0, \quad \pm \tilde{\lambda}_+(\vec{u}(t_0)) > 0. \end{aligned} \tag{2.66}$$

Then the ejection lemma [14, Lemma 3.2] followed by the one-pass theorem [14, Theorem 5.1] implies the following. If $\lambda_+ \in A_\pm$ then the solution \vec{u} is exponentially ejected out of the small neighborhood $d_W < d_W(\vec{u}(t_0))$ and never comes back again. Moreover, if $\lambda_+ \in A_+$ then u blows up in $t > t_0$, and if $\lambda_+ \in A_-$ then u scatters to 0. By the local wellposedness of (CW) in \mathcal{H} , both A_\pm are open. To see that both are non-empty, consider the case $\delta_+ \sim |\lambda_+| \gg \|\varphi\|_{\mathcal{H}}$. Let

$$\vec{u}(t) = \mathcal{F}_{\vec{u}(t)}(\vec{W} + \tilde{\lambda}(t)\rho + \gamma(t)), \quad \gamma(t) \perp \rho. \tag{2.67}$$

Then (1.46) implies that

$$|\tilde{\lambda}_+(0) - \lambda_+| + |\tilde{\lambda}_-(0)| \lesssim \|\varphi\|_{\mathcal{H}} \ll |\lambda_+|, \tag{2.68}$$

and, by definition of d_W , we have at $t = 0$

$$\begin{aligned} d_W(\vec{u})^2 &= E(u) - E(W) + k^2 |\tilde{\lambda}_1|^2, \\ e^{-\tilde{\sigma}} \partial_t d_W(\vec{u})^2 &= \frac{k^2}{2} (|\tilde{\lambda}_+|^2 - |\tilde{\lambda}_-|^2) + O(\|\gamma\|_{\mathcal{H}} |\tilde{\lambda}_1|^2), \\ E(u) - E(W) &= -k \tilde{\lambda}_+ \tilde{\lambda}_- + \frac{1}{2} \langle \mathcal{L}\gamma | \gamma \rangle - o(|\tilde{\lambda}|^2 + \|\gamma\|_{\mathcal{H}}^2). \end{aligned} \tag{2.69}$$

Hence we deduce

$$|E(u) - E(W)| \ll |\lambda_+|^2 \sim d_W(\vec{u}(0))^2 \sim e^{-\tilde{\sigma}} \partial_t d_W(\vec{u}(0))^2 > 0, \tag{2.70}$$

and thus $\lambda_+ \in A_{\text{sign } \lambda_+}$ for $|\lambda_+| \sim \delta_+$. In particular, both A_\pm are non-empty, which implies that the remainder

$$A_0 := \mathcal{B}_{\delta_+}^+ \setminus (A_+ \cup A_-) \tag{2.71}$$

is also non-empty. Every solution u for $\lambda_+ \in A_0$ must violate (2.66) for each $t \in [0, T)$, to avoid the ejection. Hence at each $t \in (0, T)$, one of the following holds

- (1) $\partial_t d_W(\tilde{u}(t)) \leq 0$
- (2) $d_W(\tilde{u}(t))^2 \leq 2(E(u) - E(W)) \leq 2d_W(\tilde{u}(0))^2$
- (3) $d_W(\tilde{u}(t)) \geq \delta_*$,

where the last condition in (2.66) is not considered, since it is implied by the others, due to the ejection lemma. Since $d_W(\tilde{u}(0)) \lesssim \delta_+ \ll \delta_*$, either (1) or (2) holds for small $t > 0$. Since the ejection lemma can be applied with $\partial_t d_W(\tilde{u}) = 0$ as well, we deduce that $d_W(\tilde{u}(t))$ is strictly decreasing in $t > 0$ until (2) is satisfied, where \tilde{u} spends its remaining life (hence never reaching (3)).

Choosing $\delta_+ \ll \iota_I^6$ small enough, we can ensure that

$$d_W(\tilde{u}(0))^2 + |\lambda_+| \ll \iota_I^6 \tag{2.72}$$

for all $(\lambda_+, \varphi) \in \mathcal{B}_{\delta_+}^+ \times \mathcal{B}'_{\delta'}$. If there are more than one $\lambda_+ \in A_0$ for the same φ , say $\lambda_+^0 \neq \lambda_+^1$, then we can apply Lemma 2.2 to the corresponding solutions with the initial data $\tilde{u}^j(0) = \tilde{W} + \lambda_{+}^j g_+ + \varphi$ and with $\iota \in (0, \iota_I]$ satisfying

$$d_W(\tilde{u}^j(0))^2 + |\lambda_{+}^j| \ll \iota^6. \tag{2.73}$$

Then its conclusion together with (2) leads to a contradiction

$$\iota^3 \sim d_W(\tilde{u}^0(\iota_I)) + d_W(\tilde{u}^1(\iota_I)) \lesssim d_W(\tilde{u}^0(0)) + d_W(\tilde{u}^1(0)) \ll \iota^3. \tag{2.74}$$

Thus we can define the functional m_+ by putting $A_0 = \{m_+(\varphi)\}$, and then $A_+ = (m_+(\varphi), \delta_+)$, $A_- = (-\delta_+, m_+(\varphi))$. The same reasoning as above implies that we can never apply Lemma 2.2 for different $\varphi^0, \varphi^1 \in \mathcal{B}'_{\delta'}$, with $\tilde{u}^j(0) = \tilde{W} + m_+(\varphi^j)g_+ + \varphi^j$ and $d_W^2(\tilde{u}^j(0)) + \|\varphi^j\|_{\mathcal{H}} \ll \iota^6$. Therefore

$$\max_{j=0,1} \left[|m_+(\varphi^j)| + \|\varphi^j\|_{\mathcal{H}} \right] \ll \iota^6 < \iota_I^6 \implies |\langle m_+(\varphi^\triangleright) | \rangle| \lesssim \iota \|\varphi^\triangleright\|_{\mathcal{H}}, \tag{2.75}$$

which implies the Lipschitz continuity (2.7). Since $m_+(0) = 0$ is obvious, it also implies that $|m_+(\varphi)| = o(\|\varphi\|_{\mathcal{H}})$. In particular, we may restrict both the domain and the range of m_+ to have the same radius δ_m as in the statement of the theorem, though there is no merit for that besides reducing the number of parameters.

The trichotomic dynamics readily follows from the ejection lemma and the one-pass theorem in [14]. The estimate on the L^{2^*} distance in (1) is derived from the bound $K(u(t)) \leq -\kappa(\delta_*)$ in the variational region [14, Lemma 4.1] as follows. Choose $\delta_m \ll \kappa(\delta_*)$. Let u be a solution in the case (1) and let t be after the ejection, namely $d_W(\tilde{u}(t)) > \delta_*$. Let $u(t) = \psi + \varphi$ with $\psi = \mathcal{T}^c S_{-1}^\sigma W$ for some $(\sigma, c) \in \mathbb{R}^{1+d}$. Then

$$\begin{aligned} \kappa(\delta_*) \leq -K(u(t)) &= -K(\psi) - \|\nabla\varphi\|_{L^2}^2 + 2\langle \Delta\psi | \varphi \rangle + \|\psi + \varphi\|_{L^{2^*}} - \|\psi\|_{L^{2^*}} \\ &= -\|\nabla\varphi\|_{L^2}^2 - 2\langle \psi^{2^*-1} | \varphi \rangle + \|\psi + \varphi\|_{L^{2^*}} - \|\psi\|_{L^{2^*}}, \end{aligned} \tag{2.76}$$

which implies $\kappa(\delta_*) \lesssim \|\varphi\|_{L^{2^*}}$ and so $\text{dist}_{L^{2^*}}(u(t), S_{\text{tatic}}(W)_1) \gtrsim \kappa(\delta_*) \gg \delta_m$. Thus it only remains to prove that m_+ is C^1 .

2.3 Smoothness of the manifold

Next we prove that the center-stable manifold obtained above is at least C^1 in \mathcal{H} . Let $\tilde{u}^0 \in \text{Solution}([0, T])$ be a solution on the manifold with the modulation and the rescaled variables

$$(\sigma, c) = (\tilde{\sigma}, \tilde{c})(\tilde{u}^0(t)), \quad \frac{d\tau}{dt} = e^{\sigma(t)}, \quad \tau(0) = 0. \tag{2.77}$$

We consider solutions \tilde{u}^1 close to \tilde{u}^0 , in the form

$$\begin{aligned} \tilde{u}^1(t) &= \tilde{u}^0(t) + \mathcal{F}_{\tilde{u}^0(t)} h \check{v}(t, h), \quad \check{v}(t, h) = \check{\lambda}(t, h)\rho + \check{\gamma}(t, h), \quad \check{\gamma} \perp \rho, \\ \lambda_-^1(0)g_- + \gamma^1(0) &:= \varphi^0 + h\varphi', \quad (h \rightarrow 0), \end{aligned} \tag{2.78}$$

for each $\varphi^0 \in \mathcal{B}'_{\delta_m}$ and $\varphi' \in \mathcal{B}'_{\infty}$. Let $\tilde{u}^j = \Phi_{\sigma, c}(\lambda^j, \gamma^j)$, then $(\check{\lambda}, \check{\gamma}) = \langle \lambda^{\triangleright}, \gamma^{\triangleright} \rangle / h$. To put both \tilde{u}^j on the manifold, we need

$$\lambda_+^0(0) = m_+(\varphi^0), \quad \lambda_+^1(0) = m_+(\varphi^0 + h\varphi'). \tag{2.79}$$

Let $(\check{\alpha}, \check{\mu}) := \langle \check{\gamma}_1 | (S'_0, T')\rho \rangle$. From the equation (2.38) of $\langle \tilde{u}^{\triangleright} \rangle$, we obtain

$$\begin{aligned} \partial_{\tau} \check{\lambda}_1 &= \check{\lambda}_2 + \sigma_{\tau}(\check{\alpha} + \check{\lambda}_1) + e^{\sigma} c_{\tau} \check{\mu}, \\ \partial_{\tau} \check{\lambda}_2 &= k^2 \check{\lambda}_1 + \langle \triangleleft N(v_1^{\triangleright}) / h - Z_2 \check{\gamma}_2 | \rho \rangle, \\ \partial_{\tau} (\check{\alpha}, \check{\mu}) &= \langle \check{\gamma}_2 - Z_1 \check{v}_1 | (S'_0, T')\rho \rangle, \\ \partial_{\tau} \check{\gamma} &= J\mathcal{L}\check{\gamma} + P_{\perp}[\langle \triangleleft N(v_1^{\triangleright}) / h - Z\check{v} \rangle]. \end{aligned} \tag{2.80}$$

By Lemma 2.3, \check{v} is bounded in $L^{\infty}\mathcal{H} \cap \text{St}$ as $h \rightarrow 0$ locally uniformly on $0 < \tau < \infty$. Moreover, the small Lipschitz property of m_+ implies that

$$|\check{\lambda}_+(0, h)| \ll \|\varphi'\|_{\mathcal{H}}. \tag{2.81}$$

Hence there is a sequence $h \rightarrow 0$ along which $\check{\lambda}_+(0, h)$ converges to some $\lambda_+^{\infty} \in \mathbb{R}$, and the limit of $\check{v} = \check{\lambda}\rho + \check{\gamma} \rightarrow v' = \lambda'\rho + \gamma'$ satisfies the linearized equation

$$\begin{aligned} \partial_{\tau} \lambda'_1 &= \lambda'_2 + \sigma_{\tau} \lambda'_1 + \langle \gamma'_1 | Z_1 \rho \rangle, \\ \partial_{\tau} \lambda'_2 &= k^2 \lambda'_1 + \langle N'(v_1^0) v'_1 - Z_2 \gamma'_2 | \rho \rangle, \\ \partial_{\tau} \gamma' &= J\mathcal{L}\gamma' + P_{\perp}[N'(v_1^0) v'_1 - Z v'], \end{aligned} \tag{2.82}$$

where $N'(v_1^0) := f''(W + v_1^0) - f''(W)$, with the initial data $\lambda'(0)\rho + \gamma'(0) = \lambda_+^{\infty}g_+ + \varphi'$. Regarding (λ', γ') as the unknown variables, this system is almost the same

as (2.38) for $\langle v^p \rangle$, except for $N'(v_1^0)v_1'$, which is the leading term of the nonlinearity in the latter system. In particular, we can show, by the same argument⁴ as for Lemma 2.2, that unless $|\lambda'_+(0)|$ is much smaller than $|\lambda'_-(0)| + \|\gamma'(0)\|_{\mathcal{H}}$, we have that $\lambda'_+(\tau)$ grows exponentially and so becomes dominant over the other components. More precisely,

Lemma 2.4 *There exists a constant $1 < C_D < \infty$ with the following property. Let $\bar{u}^0 \in \text{Solution}([0, T])$ satisfy⁵ $\|d_W(\bar{u}^0)\|_{L^\infty(0, T)} < \iota^3 \leq \iota_I^3$. Let*

$$(\sigma, c) = (\bar{\sigma}, \bar{c})(\bar{u}), \quad v^0 = \Psi_{\sigma, c}^{-1}(\bar{u}^0), \quad \frac{d\tau}{dt} = e^{\sigma(t)}, \quad \tau(0) = 0. \tag{2.83}$$

Then equation (2.82) has a unique global solution $(\lambda'(\tau), \gamma'(\tau)) : [0, \infty) \rightarrow \mathbb{R}^2 \times \mathcal{H}_\perp$ for any initial data $(\lambda'(0), \gamma'(0)) \in \mathbb{R}^2 \times \mathcal{H}_\perp$. Moreover, if

$$C_D \iota \|(\lambda'_-(\tau_0), \gamma'(\tau_0))\|_{\mathbb{R} \times \mathcal{H}} \leq |\lambda'_+(\tau_0)| \tag{2.84}$$

at some $\tau_0 \geq 0$, then there exists $\tau_I \in (\tau_0, \infty)$, such that for all $\tau > \tau_I$ we have

$$C_D |\lambda'_+(\tau)| > |\lambda'_+(\tau_I)| e^{k(\tau - \tau_I)/2} + \|(\lambda'_-(\tau), \gamma'(\tau))\|_{\mathbb{R} \times \mathcal{H}} / \iota. \tag{2.85}$$

On the other hand, if (2.84) fails for all $\tau_0 \geq 0$, then for all $\tau > 0$

$$|\lambda'_+(\tau)| \lesssim \iota \|(\lambda'_-(\tau), \gamma'(\tau))\|_{\mathbb{R} \times \mathcal{H}} \lesssim e^{C_I \tau} \|(\lambda'_-(0), \gamma'(0))\|_{\mathbb{R} \times \mathcal{H}}. \tag{2.86}$$

Proof We only sketch the proof for (2.86), since the rest is essentially the same as Lemma 2.2. In the same way as for (2.52), we obtain, for any $0 < \tau_0 < \tau < \tau_0 + \eta_I$,

$$\begin{aligned} v(\tau) &\leq (1 + C\iota)v(\tau_0) + C\iota^{3/2}|\lambda'(\tau_0)|, \\ |(\partial_\tau + k)\lambda'_-(\tau)| &\lesssim \iota^3|\lambda'(\tau_0)| + \iota^2v(\tau_0), \end{aligned} \tag{2.87}$$

where $v(\tau) := \sqrt{\iota^2|(\alpha', \mu')|^2 + \langle \mathcal{L}\gamma' | \gamma' \rangle}$ and $(\alpha', \mu') := \langle \gamma'_1 | (S', T')\rho \rangle$. Using that $|\lambda'_+| \lesssim \iota|\lambda'_-| + v$, we deduce from the above estimate

$$v(\tau) + \iota^{1/2}|\lambda'_-(\tau)| \lesssim e^{C_I \tau} [v(0) + \iota^{1/2}|\lambda'_-(0)|], \tag{2.88}$$

and so, using (1.43), we obtain (2.86). □

The above lemma implies that for each $(\lambda'_-(0), \gamma'(0)) \in \mathbb{R} \times \mathcal{H}_\perp$, there is a unique $\check{m}_+(\lambda'_-(0), \gamma'(0)) \in \mathbb{R}$ such that if $\lambda'_+(0) = \check{m}_+$ then (2.84) is not satisfied at any $\tau_0 \geq 0$, and if $\pm(\lambda'_+(0) - \check{m}_+) > 0$ then $\pm\lambda'_+(\tau)$ grows exponentially to ∞ . To see that $\check{\lambda}_+(0, h) \rightarrow \check{m}_+$, we apply Lemma 2.2 to \bar{u}^0 and \bar{u}^1 . Since $\bar{u}^1(0) \rightarrow \bar{u}^0(0)$ as $h \rightarrow 0$, the local wellposedness implies that $\bar{u}^1 \rightarrow \bar{u}^0$ locally uniformly on $[0, T)$,

⁴ Note that the proof of Lemma 2.2 did not use any particular structure of $\langle N(v^p) \rangle$.

⁵ The constant ι_I is chosen here to be the same as in Lemma 2.2, just for convenience. It does not mean that the admissible range of ι_I is exactly the same for these two lemmas.

and so does $(\tilde{\sigma}(\tilde{u}^1), \tilde{c}(\tilde{u}^1))$. Hence for any $S < \infty$, for $|h|$ small enough we could apply Lemma 2.2 to \tilde{u}^0 and \tilde{u}^1 starting at any $\tau \in [0, S]$, if we had

$$2\iota_I \|\llcorner \tilde{v}^\triangleright(\tau)\|_{\mathcal{H}} < |\llcorner \lambda'_+(\tau)|, \tag{2.89}$$

but its conclusion would contradict that both \tilde{u}^j are on \mathcal{M}_0 (specifically between (2.4) and (2.15)). Hence for all $\tau > 0$, as $h \rightarrow 0$ along the sequence,

$$\iota_I \gtrsim \frac{|\check{\lambda}_+(\tau, h)|}{|\check{\lambda}_-(\tau, h)| + \|\check{\gamma}(\tau, h)\|_{\mathcal{H}}} \rightarrow \frac{|\lambda'_+(\tau)|}{|\lambda'_-(\tau)| + \|\gamma'(\tau)\|_{\mathcal{H}}}, \tag{2.90}$$

which implies $\lambda'_+(0) = \check{m}_+$, since otherwise the right hand side will grow in τ at least to $O(1/\iota_I)$.

Therefore $\check{\lambda}_+(0, h) \rightarrow \check{m}_+(\lambda'_-(0), \gamma'(0))$ as $h \rightarrow 0$, without restricting h to a sequence. In other words, m_+ is Gâteaux differentiable at φ^0

$$\check{m}_+(\lambda'_-(0), \gamma'(0)) = \lim_{h \rightarrow 0} \frac{m_+(\varphi^0 + h\varphi') - m_+(\varphi^0)}{h} = m'_+(\varphi^0)(\varphi'). \tag{2.91}$$

The linearity of m'_+ on φ' is clear from the definition of \check{m}_+ , while its boundedness follows from the Lipschitz property of m_+ .

To show the continuity of m'_+ for φ^0 , take any sequence $\varphi_n^0 \rightarrow \varphi^0$ in \mathcal{B}'_{δ_m} , let \tilde{u}_n^0 be the solution starting from $\tilde{u}_n^0(0) = \Psi_{\sigma(0), c(0)}(m_+(\varphi_n^0)g_+ + \varphi_n^0)$ and let $\tilde{v}_n^0 := \Psi_{\sigma, c}^{-1}(\tilde{u}_n^0)$. The local wellposedness implies that $\tilde{v}_n^0 \rightarrow \tilde{v}^0$ in $L^\infty_{\tau} \mathcal{H} \cap \text{St}_{\tau}(0, S)$ for any $S \in (0, \infty)$. Let (2.82)_n be the equation obtained by replacing v^0 with v_n^0 in (2.82).

For any small $\zeta > 0$, Lemma 2.4 allows one to choose $S \gg 1$ such that the solution of (2.82) with $\lambda'_+(\tau) = 1$ and $(\lambda'_-(\tau), \gamma'(\tau)) = 0$ satisfies

$$C_D \iota_I \lambda'_+(\tau) > \|(\lambda'_-(\tau), \gamma'(\tau))\|_{\mathbb{R} \times \mathcal{H}} + e^{k\tau/2}/\zeta \tag{2.92}$$

for all $\tau > S/2$. On the other hand, for any $\varphi' \in \mathcal{B}'_1$, the solution of (2.82) with $\lambda'_+(\tau) = m'_+(\varphi^0)\varphi'$ and $\lambda'_-(\tau)g_- + \gamma'(\tau) = \varphi'$ satisfies

$$\|(\lambda'(\tau), \gamma'(\tau))\|_{\mathbb{R}^2 \times \mathcal{H}} \lesssim \iota_I^{-1} e^{C\iota_I \tau} \tag{2.93}$$

for all $\tau \geq 0$. Since $S \gg 1$ and $k \gg \iota_I$, combining these two estimates yields that the solution of (2.82) with $|\lambda'_+(\tau) - m'_+(\varphi^0)\varphi'| > \zeta$ and $\lambda'_-(\tau)g_- + \gamma'(\tau) = \varphi'$ satisfies

$$\iota_I |\lambda'_+(\tau)| \gtrsim \|(\lambda'_-(\tau), \gamma'(\tau))\|_{\mathbb{R} \times \mathcal{H}} \tag{2.94}$$

for all $\tau > S/2$.

The local uniform convergence of v_n^0 implies that the solution of (2.82)_n with the same initial data also satisfies (2.94) around $\tau = S$ for large n . Moreover, since $(\tilde{\sigma}(\tilde{u}_n^0), \tilde{c}(\tilde{u}_n^0)) \rightarrow (\sigma, c)$ uniformly on $[0, 2S]$ as $n \rightarrow \infty$, we have the same estimate

(2.94) around $\tau = S$ also in the coordinate associated with the solution \vec{u}_n^0 , for large n . Then it implies that

$$|m'_+(\varphi_n^0)\varphi' - m'_+(\varphi^0)\varphi'| \leq \zeta, \quad (2.95)$$

for all $\varphi' \in B'_1$. Hence m'_+ is continuous $\mathcal{B}'_{\delta_m} \rightarrow (\mathbb{R} \times \mathcal{H}_\perp)^*$. This concludes the proof of Theorem 2.1.

3 Extension of the manifold and the 9-set dynamics

3.1 Extension by the backward flow

By the maximal evolution, we can extend \mathcal{M}_0 to an invariant manifold:

$$\mathcal{M}_1 := \bigcup \{ \vec{u}(I) \mid \vec{u}(0) \in \mathcal{M}_0, \vec{u} \in \text{Solution}(I) \} \supset \mathcal{M}_0. \quad (3.1)$$

\mathcal{M}_1 also inherits from \mathcal{M}_0 the invariance for \mathcal{T} and \mathcal{S} . By the property of \mathcal{M}_0 , those solutions are eventually trapped by the ground state, namely

$$\limsup_{t \nearrow T_+} d_W(\vec{u}(t))^2 \leq 2(E(u) - E(W)) \lesssim \delta_m^2, \quad (3.2)$$

where T_+ is the maximal existence time of u .

Conversely, if a solution u satisfies the above condition and $E(\vec{u}) - E(W) \ll \delta_m^2$, then its orbit is included in \mathcal{M}_1 . This is because every solution getting close enough to the ground states is classified by the trichotomy of Theorem 2.1, whereas those solutions which never approach the ground states have been classified into the four sets of scattering to 0 and blowup away from the ground states in [14]. One may wonder what happens if a solution stays around $d_W(\vec{u}) \sim \delta_m$, but such behavior is precluded by the ejection lemma [14, Lemma 3.2] (applied in both time directions) under the energy constraint $E(u) - E(W) \ll \delta_m^2$.

For each point $\varphi \in \mathcal{M}_1$, there is a small neighborhood $O \ni \varphi$ and $T > 0$ such that the nonlinear flow $U_N(T)$ maps O into a small neighborhood of a point on \mathcal{M}_0 , where the trichotomy holds. Then $O \cap \mathcal{M}_1$ is mapped onto $U_N(T)(O) \cap \mathcal{M}_0$. Since $U_N(T)$ is smooth on O , it implies that \mathcal{M}_1 is also a C^1 manifold of codimension 1.

The one-pass theorem in [14] implies that every solution on $\mathcal{M}_1 \cap \mathcal{M}_1^\dagger$ satisfies $d_W(\vec{u}(t)) \lesssim \delta_m$ all over its life, and so it is essentially the same as the center manifold $\mathcal{M}_0 \cap \mathcal{M}_0^\dagger$, and in particular with codimension 2. The rest of \mathcal{M}_1 is split into two parts, scattering to 0 or blowup away from the ground states, in the negative time direction. Each set is non-empty and relatively open in \mathcal{M}_0 .

Therefore, we have all 3×3 combinations of dynamics in $t > 0$ and in $t < 0$: (1) blowup away from the ground states, (2) trapping by the ground states (or by \mathcal{M}_0 for $t > 0$ and by \mathcal{M}_0^\dagger for $t < 0$), and (3) scattering to 0. It was already shown in [14] that the combinations of (1) and (3) have non-empty interior. Moreover, those 9 sets exhaust all possible dynamics in the region $E(u) - E(W) \ll \min(\delta_m, \varepsilon_*)^2$. Thus we obtain Theorem 1.1.

Before going to the next step using the Lorentz transform, it is convenient to consider the space-time maximal extension of each solution of CW.

3.2 Space-time extension and restriction

To solve the equation locally in and out of light cones, and in more general space-time sets, we introduce restricted energy semi-norms. Let \mathcal{B}_d be the totality of Borel sets in \mathbb{R}^d . For any $B \in \mathcal{B}_d$ and any $a \geq 0$, we define two sets $B_{\pm a} \in \mathcal{B}_d$ by

$$\begin{aligned} B_{+a} &:= \{x \in \mathbb{R}^d \mid \exists y \in B, |x - y| \leq a\}, \\ B_{-a} &:= \{x \in \mathbb{R}^d \mid |x - y| \leq a \implies y \in B\}. \end{aligned} \tag{3.3}$$

It is clear that for any $a, b, t \geq 0$, we have

$$\begin{aligned} (B_{+a})_{+b} &= B_{+(a+b)}, & (B_{-a})_{-b} &= B_{-(a+b)}, \\ (B_{-a})_{+a} &\subset B \subset (B_{+a})_{-a}, & (B_{+a})^c &= (B^c)_{-a}. \end{aligned} \tag{3.4}$$

For any $I \subset \mathbb{R}$ and any $F : \mathbb{R} \rightarrow \mathbb{R}$, we define $B_{\pm F}(I) \in \mathcal{B}_{1+d}$ by

$$B_{\pm F}(I) := \{(t, x) \in I \times \mathbb{R}^d \mid x \in B_{\pm F(t)}\}. \tag{3.5}$$

For any $B \in \mathcal{B}_d$, let $V(B) \subset \mathcal{H}$ be the closed subspace defined by

$$V(B) := \{\varphi \in \mathcal{H} \mid \varphi(x) = 0 \text{ a.e. } x \in B\}, \tag{3.6}$$

and then define the restriction of \mathcal{H} onto B by

$$\mathcal{H} \downarrow B := \mathcal{H}/V(B) \simeq V(B)^\perp, \tag{3.7}$$

where \simeq means the isometry, with the quotient norm

$$\|\varphi\|_{\mathcal{H} \downarrow B} := \inf\{\|\psi\|_{\mathcal{H}} \mid \psi = \varphi \text{ on } B\} \quad (\varphi \in \mathcal{H}). \tag{3.8}$$

Henceforth, we denote for brevity

$$\varphi = \psi \text{ on } B \stackrel{\text{def}}{\iff} \varphi - \psi \in V(B). \tag{3.9}$$

We also use the more explicit semi-norms for $\varphi \in \mathcal{H}$:

$$\|\varphi\|_{\mathcal{H}(B)}^2 := \int_B [|\nabla u_1|^2 + |u_2|^2] dx, \quad \|\varphi\|_{\tilde{\mathcal{H}}(B)} := \|\varphi\|_{\mathcal{H}(B)} + \|\varphi_1\|_{L^{2^*}(B)}. \tag{3.10}$$

All of these three semi-norms are increasing for B and invariant for \mathcal{T} , \mathcal{S} , namely

$$\|\varphi\|_{X(B_1)} \leq \|\varphi\|_{X(B_1 \cup B_2)}, \quad \|\mathcal{T}^c \mathcal{S}^s \varphi\|_{X(B)} = \|\varphi\|_{X(e^{-\sigma} B + c)}, \quad (3.11)$$

for $X = \mathcal{H}$, $\tilde{\mathcal{H}}$, and $\mathcal{H} \downarrow$. We have, uniformly for B ,

$$\|\varphi\|_{\mathcal{H}(B)} \leq \|\varphi\|_{\mathcal{H} \downarrow B}, \quad \|\varphi\|_{\tilde{\mathcal{H}}(B)} \lesssim \|\varphi\|_{\mathcal{H} \downarrow B}. \quad (3.12)$$

We may have the reverse inequalities when B is smooth. In particular, we have

$$\begin{aligned} \|\varphi\|_{\mathcal{H} \downarrow \{|x-c| < R\}} &\sim \|\varphi\|_{\tilde{\mathcal{H}}(\{|x-c| < R\})}, \\ \|\varphi\|_{\mathcal{H} \downarrow \{|x-c| > R\}} &\sim \|\varphi\|_{\tilde{\mathcal{H}}(\{|x-c| > R\})} \sim \|\varphi\|_{\mathcal{H}(\{|x-c| > R\})}, \end{aligned} \quad (3.13)$$

uniformly for c and R , where the open region can be replaced with the closure. The extension operator $X_B : \mathcal{H} \rightarrow V(B)^\perp \subset \mathcal{H}$ is nothing but the orthogonal projection to $V(B)^\perp$, such that we have

$$\|X_B \varphi\|_{\mathcal{H}} = \|\varphi\|_{\mathcal{H} \downarrow B}, \quad X_B \varphi = \varphi \text{ on } B. \quad (3.14)$$

Restriction of energy-type functionals is denoted as follows

$$E_B(\varphi) := \frac{\|\varphi\|_{\mathcal{H}(B)}^2}{2} - \frac{\|\varphi_1\|_{L^{2^*}(B)}^{2^*}}{2^*}, \quad K_B(\varphi) := \|\nabla \varphi\|_{L^2(B)}^2 - \|\varphi\|_{L^{2^*}(B)}^{2^*}. \quad (3.15)$$

The finite propagation speed implies that if a solution u of CW satisfies

$$\vec{u}(0) = \psi \in \mathcal{H} \downarrow B \text{ on } B, \quad (3.16)$$

then u is uniquely determined on $B_{-|t|}$ by ψ . More precisely, if $\vec{u}^0, \vec{u}^1 \in C([0, T]; \mathcal{H})$ satisfy $\vec{u}^0(0) = \vec{u}^1(0)$ on B , then

$$0 \leq \forall t \leq T, \text{ a.e. } x \in B_{-|t|}, \quad \vec{u}_0(t, x) = \vec{u}_1(t, x). \quad (3.17)$$

By the Strichartz estimate, there is $C > 0$ such that if $C \|\psi\|_{\mathcal{H} \downarrow B} \leq \varepsilon_S$ then there is a free solution v satisfying $\vec{v}(0) = \psi$ on B and $\|v\|_{\text{St}(\mathbb{R})} \leq \varepsilon_S$, and so is $u \in \text{Solution}(\mathbb{R})$ satisfying $\vec{u}(0) = \psi$ on B and $\|u\|_{\text{St}(\mathbb{R})} \lesssim \varepsilon_S$, which is unique on $B_{-|t|}(\mathbb{R})$.

Now we introduce the space-time maximal extension of a solution of (CW). For any $\varphi \in \mathcal{H}$, $c \in \mathbb{R}^d$ and $R > 0$, consider the local solution in the light cone $\mathcal{K}_{c,R} := \{|x-c| + t < R, t \geq 0\}$ with the initial data $\vec{u}(0) = \varphi$ on $|x-c| < R$. Let $t_+(\varphi, c)$ be the supremum of such R that there is a unique solution u in $\mathcal{K}_{c,R}$ satisfying $\|u\|_{L^{qm}(\mathcal{K}_{c,R})} < \infty$. The uniqueness in cones implies that we have a unique solution u in the space-time region

$$\{(t, x) \in \mathbb{R}^{1+d} \mid 0 \leq t < t_+(\varphi, x)\}, \quad (3.18)$$

as well as the Lipschitz continuity

$$|t_+(\varphi, x) - t_+(\varphi, y)| \leq |x - y|. \quad (3.19)$$

We also write for any strong solution u (either before or after the above space-time maximal extension),

$$t_+(u, x) := t_+(\vec{u}(0), x). \quad (3.20)$$

The maximal existence time is then given by

$$T_+(u) := \inf_{x \in \mathbb{R}^d} t_+(u, x). \quad (3.21)$$

The small data theory in interior and exterior cones implies that for any $\varphi \in \mathcal{H}$, there are $a(\varphi), b(\varphi) > 0$ such that

$$t_+(\varphi, x) \geq \max(a(\varphi), |x| - b(\varphi)). \quad (3.22)$$

The definition of t_+ implies that

$$\|u\|_{L^{qm}(\mathcal{K}_{c, t_+(u, c)})} = \lim_{R \nearrow t_+(u, c)} \|u\|_{L^{qm}(\mathcal{K}_{c, R})} = \infty, \quad (3.23)$$

and so the small data theory implies

$$\liminf_{t \nearrow t_+(u, c)} \|\vec{u}(t)\|_{\tilde{\mathcal{H}}(|x-c| < t_+(u, c) - t)} \gtrsim \varepsilon_S. \quad (3.24)$$

In particular, the number of first blow-up points is bounded in the case of type-II

$$\#\{c \in \mathbb{R}^d \mid t_+(u, c) = T_+(u)\} \lesssim \liminf_{t \nearrow T_+(u)} \sqrt{\|\vec{u}(t)\|_{\mathcal{H}}/\varepsilon_S}. \quad (3.25)$$

Similarly we can define $t_-(\varphi, c) < 0$ to be the maximal extension in the negative direction, and thus a unique solution u in the maximal space-time domain

$$\mathcal{D}(u) := \mathcal{D}(\varphi) := \{(t, x) \in \mathbb{R}^{1+d} \mid t_-(\varphi, x) < t < t_+(\varphi, x)\}, \quad (3.26)$$

satisfying for some $a(\varphi), b(\varphi) > 0$ and for all $x, y \in \mathbb{R}^d$,

$$\pm t_{\pm}(\varphi, x) \geq \max(a(\varphi), |x| - b(\varphi)), \quad |t_{\pm}(\varphi, x) - t_{\pm}(\varphi, y)| \leq |x - y|. \quad (3.27)$$

Since the Lorentz transforms preserve light cones as well as the measure and the topology of \mathbb{R}^{1+d} , the property (3.23) of t_+ is also preserved. Hence, each strong solution u defined on its maximal space-time domain $\mathcal{D}(u)$ is transformed by any Lorentz transform into another solution defined on the maximal domain. This process can produce a solution with no Cauchy time slice, namely $\inf t^+ < \sup t^-$, but we ignore such solutions, in order to keep the dynamical viewpoint in terms of the Cauchy

problem or the flow in \mathcal{H} . In other words, the Lorentz transforms should be restricted to the range where $\inf t^+ < \sup t^-$ is kept.

For the blow-up solutions on the center-stable manifold, we have

Lemma 3.1 *Let $0 < T < \infty$ and $\vec{u} \in S_{\text{solution}}([0, T])$ satisfy*

$$\vec{u}(0) \in \mathcal{M}_0, \quad \|d_W(\vec{u})\|_{L^\infty(0,T)} \leq \delta \lesssim \delta_m. \tag{3.28}$$

Then there exists $c_ \in \mathbb{R}^d$ and $\varepsilon > 0$ such that*

$$\begin{aligned} e^{-\tilde{\sigma}(\vec{u}(t))} + |\tilde{c}(\vec{u}(t)) - c_*| &\lesssim \delta |t - T|, \\ t_+(\vec{u}(0), x) = T + |x - c_*|, \quad t_-(\vec{u}(0), x) &< T - |x - c_*| - \varepsilon. \end{aligned} \tag{3.29}$$

Proof Let $(\sigma(t), c(t)) := (\tilde{\sigma}(\vec{u}(t)), \tilde{c}(\vec{u}(t)))$. First we show $\sigma(t) \rightarrow \infty$ as $t \rightarrow T - 0$. If not, there exist a sequence $t_n \nearrow T$ with $\sup_n \sigma(t_n) < \infty$ and $R > 0$ such that

$$\sup_n \sup_{c \in \mathbb{R}^d} \|\vec{u}(t_n)\|_{\tilde{\mathcal{H}}(|x-c|<R)} \lesssim \delta \ll \varepsilon_S, \tag{3.30}$$

which ensures solvability in the cone $|x - c| + |t - t_n| < R$ for all $c \in \mathbb{R}^d$ and all t_n , thereby extending the solution u beyond T , a contradiction. Hence $\sigma(t) \rightarrow \infty$. Then the modulation equation (1.56) implies convergence $c(t) \rightarrow \exists c_* \in \mathbb{R}^d$ as well as the first estimate of (3.29). Since $\|\vec{u}(t) - \mathcal{T}^c \mathcal{S}^\sigma \vec{W}\|_{\mathcal{H}} \sim d_W(\vec{u}(t)) < \delta$, that behavior of (σ, c) implies that for any $R > 0$,

$$\limsup_{t \nearrow T} \|\vec{u}(t)\|_{\mathcal{H}(|x-c_*|>R)} \lesssim \delta \ll \varepsilon_S, \tag{3.31}$$

which ensures solvability in the exterior cone $|x - c_*| + |t - T'| > R$ for all $T' < T$ close to T , and so $t_+(\vec{u}(0)) \geq T' + |x - c_*| - R$ and $t_-(\vec{u}(0)) \leq T' - |x - c_*| + R$. Letting $R \searrow 0$ and $T' \nearrow T$, we obtain

$$t_+ \geq T + |x - c_*|, \quad t_- \leq T - |x - c_*|. \tag{3.32}$$

It can not be better for t_+ as u blows up at (T, c_*) , so we obtain the identity in (3.29). The finite propagation implies $\|\vec{u}(t)\|_{\mathcal{H}(|x-c_*|>R+|T-t|)} \lesssim \delta$, so that we can solve in a slightly larger exterior cone starting from $t = T' \in (0, T)$. Thus we obtain the last estimate in (3.29). □

3.3 Extension by the Lorentz transform

Next we use the Lorentz transforms to extend the manifold, so that it can include all the ground state solitons $S_{\text{soliton}}(W)$.

Let $\vec{u} \in S_{\text{solution}}([0, T])$, $\vec{u}(0) \in \mathcal{M}_0$ and $T < \infty$. The estimate (3.29) on the blowup surface implies existence of $T' \in (0, T)$ such that $\{|x - c_*| \geq |t - T'|\} \subset \mathcal{D}(u)$. Then for any Lorentz transform centered at (T', c_*) , the solution u is transformed into

another solution on the maximal domain containing the time slice $\{t = T'\}$. In the case where $\vec{u}(0) \in \mathcal{M}_0$ yields $\vec{u} \in \text{Solition}([0, \infty))$, Lorentz transformed solutions are also defined for all large t , because of (3.27).

Let u be any solution with $\vec{u}(0) \in \mathcal{M}_0$ defined on $\mathcal{D}(u)$, and let w be any Lorentz transform of u . The above argument implies that w is defined on some time slice. Let \mathcal{M}_2 be the totality of the maximal orbit of all such solutions w . Then it is invariant by the flow, \mathcal{T} and \mathcal{S} , satisfying

$$\mathcal{M}_2 \supset \mathcal{M}_1 \cup \text{Soliton}(W). \tag{3.33}$$

Every solution on \mathcal{M}_2 is Lorentz transformed to another solution on \mathcal{M}_1 . \mathcal{M}_2 is Lorentz invariant in the following sense: Let u be a solution on \mathcal{M}_2 , extended to $\mathcal{D}(u)$. Then any Lorentz transform of u has non-trivial maximal existence interval, on which the transformed solution belongs to \mathcal{M}_2 .

To see that \mathcal{M}_2 is locally C^1 diffeo to \mathcal{M}_1 , it suffices to see that the Lorentz transform gives a local C^1 mapping around any solution. Let $\vec{u} \in \text{Solition}([-T, T])$ for some $T > 0$, then the local wellposedness yields a neighborhood $O \ni \vec{u}(0)$ and $R > 0$ such that for any $\varphi \in O$ we have

$$\mathcal{D}(\varphi) \supset X := \{(t, x) \in \mathbb{R}^{1+d} \mid |t| < \max(T/2, |x| - R)\}. \tag{3.34}$$

Then there is a neighborhood $U \ni 1$ in the Lorentz group such that every transform in U maps the region X to a set containing $[-T/4, T/4] \times \mathbb{R}^d$. Since the space rotation plays no role, we may restrict to those transforms defined on $(t, x_1) \in \mathbb{R}^2$, which can be parametrized as

$$u^\theta(t, y, z) = u(ct + sy, cy + st, z), \quad c := \cosh \theta, \quad s := \sinh \theta, \quad \theta \in \mathbb{R}, \tag{3.35}$$

where $x = (y, z) \in \mathbb{R}^1 \times \mathbb{R}^{d-1}$. Then there is $\Theta > 0$ such that for any $\vec{u}(0) \in O$ and for any $\theta \in (-\Theta, \Theta)$, we have $\mathcal{D}(u^\theta) \supset [-T/4, T/4] \times \mathbb{R}^d$. This defines a mapping

$$\mathcal{S}_\theta : O \ni \vec{u}(0) \mapsto \vec{u}^\theta(0) \in \mathcal{H} \tag{3.36}$$

for each $\theta \in (-\Theta, \Theta)$. The continuity of \mathcal{S}_θ can be seen by the linear energy identity. For any smooth function u defined on X , we have by the divergence theorem

$$\begin{aligned} E^F(\vec{u}^\theta(0)) &= cE^F(\vec{u}(0)) + sP_1(\vec{u}(0)) + \int_{\frac{sy}{ct} > 1} (\ddot{u} - \Delta u) \frac{t}{|t|} (cu_t + su_y) dxdt, \\ P_1(\vec{u}^\theta(0)) &= sE^F(\vec{u}(0)) + cP_1(\vec{u}(0)) + \int_{\frac{sy}{ct} > 1} (\ddot{u} - \Delta u) \frac{t}{|t|} (su_t + cu_y) dxdt, \end{aligned} \tag{3.37}$$

where $E^F(\varphi) := \|\varphi\|_{\mathcal{H}}^2/2$ denotes the free energy. Applying this to the difference of two solutions u, w starting from O yields

$$\begin{aligned} \|\vec{u}^\theta(0) - \vec{w}^\theta(0)\|_{\mathcal{H}}^2 &\lesssim \|\vec{u}(0) - \vec{w}(0)\|_{\mathcal{H}}^2 \\ &\quad + \|f'(u) - f'(w)\|_{L_t^1 L_x^2(sy/(ct) < 1)} \|\vec{u} - \vec{w}\|_{L_t^\infty \mathcal{H}_x(sy/(ct) < 1)}, \end{aligned} \quad (3.38)$$

where the implicit constants depend on Θ . The standard perturbation argument implies that the last term is also bounded by $\|\vec{u}(0) - \vec{w}(0)\|_{\mathcal{H}}^2 \ll 1$. The existence and continuity of the derivative of \mathcal{S}_θ is shown similarly by applying the energy estimate to the linearized equation

$$(\partial_t^2 - \Delta)u' = f''(u)u', \quad \vec{u}'(0) \in \mathcal{H}. \quad (3.39)$$

Since $(\mathcal{S}_\theta)^{-1} = \mathcal{S}_{-\theta}$ is obvious, we thus conclude that \mathcal{S}_θ is a local C^1 diffeo on O . Therefore \mathcal{M}_2 is also a C^1 manifold with codimension 1 in \mathcal{H} . It is clear from the construction that all these manifolds $\mathcal{M}_0 \subset \mathcal{M}_1 \subset \mathcal{M}_2$ are connected.

For the trichotomy around \mathcal{M}_2 , it is obvious from the energy estimate that every scattering (to 0 as $t \rightarrow \infty$) solution is transformed into another such solution by any Lorentz transform. The solutions in the other part of the neighborhood blow up away from a (much) bigger neighborhood, which is generated from the δ_X neighborhood of $S_{\text{static}}(W)$ by the above extensions. Thus we obtain the 9-set dynamics classification in the region

$$E(u) < \sqrt{(E(W) + \varepsilon^2)^2 + |P(u)|^2}. \quad (3.40)$$

Reduction of this region to $E(u) < E(W) + \varepsilon^2$ is done as in [14] by the Lorentz transform and the identities for (CW)

$$E(\vec{u}^\theta) = cE(\vec{u}) + sP_1(\vec{u}), \quad P_1(\vec{u}^\theta) = sE(\vec{u}) + cP_1(\vec{u}). \quad (3.41)$$

Indeed, if $E(u) < |P(u)|$ then we can transform it (in some space-time region) to another solution with negative energy, which has to blow up in both time directions by the classical argument of Levine [18], or more precisely by [11]. Hence the solution before the transform should also blow up in both directions.

If $E(u) = |P(u)|$ and it is global for $t > 0$, then there is a sequence of solutions $\vec{u}^n \in S_{\text{solution}}([0, \infty))$ given by Lorentz transforms such that $E(u_n) \rightarrow 0$. Then the classical argument of Payne-Sattinger [23] implies that $K(u_n) \geq 0$ as soon as $E(u_n) < E(W)$, and so $E(u_n) \sim \|\vec{u}_n\|_{L^\infty \mathcal{H}}^2 \rightarrow 0$. The small data scattering implies that $\|u_n\|_{L^{qm}(\mathbb{R}^{1+d})} \lesssim E(u_n)^{1/2} \rightarrow 0$, but since the Lorentz transform is measure preserving on \mathbb{R}^{1+d} , it implies that $\|u\|_{L^{qm}(\mathbb{R}^{1+d})} = 0$. In short, all the solutions with $E(u) \leq |P(u)|$ blow up in both time directions except for the trivial solution 0.

If $E(u) > |P(u)|$, then we can transform it to another solution \vec{u} with $P(\vec{u}) = 0$ and $E(\vec{u}) < E(W) + \varepsilon^2$, and so \vec{u} should either scatter to 0 as $t \rightarrow \infty$, blow up away from the ground state in the positive time direction, or live on \mathcal{M}_1 . Each of those properties is transferred back to the original solution u . Note that if $\mathcal{D}(\vec{u})$ contains no time slice then the original solution u must blow up in both time directions. Thus we complete the 9-set dynamics classification slightly above the ground states, and the proof of Theorem 1.2.

Part II: Large radiation

The goal in the rest of paper is to extend the center-stable manifold to the entire energy space \mathcal{H} , together with the dynamics around it, by a simple argument which allows one to reduce the problem to \mathcal{M}_0 in the region $E(u) < E(W) + \varepsilon^2$, using the asymptotic Huygens principle together with the finite speed of propagation.

4 Detaching the radiation

For any $B \in \mathcal{B}_d$ and any $T \in (0, \infty]$, we define a semi-norm \mathcal{R}_B^T in \mathcal{H} by

$$\|\varphi\|_{\mathcal{R}_B^T} := \inf\{\|U(t)\psi\|_{L_t^\infty(0,T;\mathcal{H}|_{B_{+t}})\cap\text{St}(0,T)} \mid \psi = \varphi \text{ on } B^{\mathbb{C}}\}. \tag{4.1}$$

Smallness in \mathcal{R}_B^T will imply that we can detach the exterior component using the wave starting from ψ which is out-going dispersive in the sense of the energy on the interior cone B_{+t} , and also the Strichartz norm on \mathbb{R}^d , both for $0 \leq t \leq T$.

The lower semi-continuity of the norms implies that the infimum in defining \mathcal{R}_B^T is achieved by some $\psi \in \mathcal{H}$ such that $\psi = \varphi$ on $B^{\mathbb{C}}$ and

$$\|\varphi\|_{\mathcal{R}_B^T} = \|U(t)\psi\|_{L_t^\infty(0,T;\mathcal{H}|_{B_{+t}})\cap\text{St}(0,T)} \lesssim \|\psi\|_{\mathcal{H}} \sim \|\varphi\|_{\mathcal{H}|_{B^{\mathbb{C}}}}. \tag{4.2}$$

For the last equivalence, \geq is by definition of $\mathcal{H}|_{B^{\mathbb{C}}}$, while \lesssim follows from

$$\|\psi\|_{\mathcal{H}} \leq \|\psi\|_{\mathcal{H}|_B} + \|\psi\|_{\mathcal{H}|_{B^{\mathbb{C}}}} \leq \|\varphi\|_{\mathcal{R}_B^T} + \|\varphi\|_{\mathcal{H}|_{B^{\mathbb{C}}}} \lesssim \|\varphi\|_{\mathcal{H}|_{B^{\mathbb{C}}}}. \tag{4.3}$$

The following laws of order are trivial by definition

$$\tau \geq 0 \implies \|\varphi\|_{\mathcal{R}_B^T} \leq \|\varphi\|_{\mathcal{R}_B^{T+\tau}}, \quad \|U(\tau)\varphi\|_{\mathcal{R}_{B_{+\tau}}^{T-\tau}} \leq \|\varphi\|_{\mathcal{R}_B^T}, \tag{4.4}$$

as is the invariance $\|T^c S^\sigma \varphi\|_{\mathcal{R}_B^T} = \|\varphi\|_{\mathcal{R}_{e^\sigma B_{+c}}^{e^\sigma T}}$, but it is not invariant under the time inversion $\varphi \mapsto \varphi^\dagger$. The space-time continuity of the norms implies

$$\lim_{R,T \rightarrow +0} \|\varphi\|_{\mathcal{R}_{|x-c|<R}^T} = 0. \tag{4.5}$$

Also note the trivial identity $\|\varphi\|_{\mathcal{R}_{\mathbb{R}^d}^T} = 0$.

The following ‘‘asymptotic Huygens principle’’ plays a crucial role in using the dispersive property of (CW) in the above function space.

Lemma 4.1 *For any free solution $\vec{v} \in C(\mathbb{R}; \mathcal{H})$ and bounded $B \in \mathcal{B}_d$ we have*

$$\lim_{T \rightarrow \infty} \sup_{t>0} \|\vec{v}(T+t)\|_{\mathcal{H}|_{B_{+t}}} = 0. \tag{4.6}$$

Proof Since B is included in some ball, it suffices to prove

$$\lim_{T \rightarrow \infty} \sup_{t > T} \|\vec{v}(t)\|_{\tilde{\mathcal{H}}(|x| < t-T)} = 0. \quad (4.7)$$

Since the statement is obviously stable in the energy norm, we may restrict the initial data to a dense set, say $C_0^\infty(\mathbb{R}^d)$. Multiplying the equation with $(t^2 + r^2)\dot{v} + 2trv_r + (d-1)tv$, we obtain conservation of the conformal energy

$$\int_{\mathbb{R}^d} |tv_t + rv_r + (d-1)v|^2 + |rv_t + tv_r|^2 + (d-1)|v|^2 + (t^2 + r^2)|\nabla^\perp v|^2 dx, \quad (4.8)$$

where $\nabla^\perp v := \nabla v - xv_r$ denotes the derivative in the angular directions. Since in the region $|x| < t - T$

$$\left| \begin{pmatrix} v_t \\ v_r \end{pmatrix} \right| = \left| \frac{1}{t^2 - r^2} \begin{pmatrix} t & -r \\ -r & t \end{pmatrix} \begin{pmatrix} tv_t + rv_r \\ rv_t + tv_r \end{pmatrix} \right| \leq \frac{1}{T} \left| \begin{pmatrix} tv_t + rv_r \\ rv_t + tv_r \end{pmatrix} \right|, \quad (4.9)$$

the L^2 norm of the left tends to 0 uniformly as $T \rightarrow \infty$, as well as those of $v/(t-T)$ and $\nabla^\perp v$, while $\|v\|_{L^{2^*}(\mathbb{R}^d)} \rightarrow 0$ by the free dispersive decay. \square

The asymptotic Huygens principle implies the following decay of \mathcal{R}_B^∞ : For any $\varphi \in \mathcal{H}$ and any bounded $B \in \mathcal{B}_d$,

$$\lim_{\tau \rightarrow \infty} \|U(\tau)\varphi\|_{\mathcal{R}_B^\infty} = 0. \quad (4.10)$$

In other words, every free solution in \mathcal{H} will eventually get into any small ball of \mathcal{R}_B^∞ around 0, as well as every scattering solution of CW in \mathcal{H} . Also, when the solution is around the ground states with large dispersive remainder, we can take the semi-norm \mathcal{R}_B^T small by the following.

Lemma 4.2 *There is $\varepsilon_D > 0$ with the following property. Let $\vec{u} \in \mathcal{S}_{\text{solution}}(I)$, $(\sigma, c) : I \rightarrow \mathbb{R}^{1+d}$, $\vec{F}(t) = U(t)\vec{F}(0) \in C(\mathbb{R}; \mathcal{H})$ and $R \in C(I; \mathcal{H})$ satisfy*

$$\begin{aligned} I \ni \forall t, \vec{u}(t) &= \mathcal{T}^{c(t)} \mathcal{S}^{\sigma(t)} \vec{W} + \vec{F}(t) + R(t), \\ \|F\|_{\text{St}(I)} + \|R\|_{L^\infty(I; \mathcal{H})} &:= \varsigma \leq \varepsilon_D, \end{aligned} \quad (4.11)$$

for some interval I . Then for any $t_0, t_1 \in I$ we have

$$|e^{-\sigma(t_0)} - e^{-\sigma(t_1)}| + |c(t_0) - c(t_1)| \lesssim \varsigma (|t_0 - t_1| + e^{-\sigma(t_0)}). \quad (4.12)$$

If $I = [0, T)$, then there are $t_0 \in (0, T)$ and $B : [t_0, T) \rightarrow \mathcal{B}_d$ such that

$$\sup_{t_0 \leq t < T} \|\mathcal{T}^{c(t)} \mathcal{S}^{\sigma(t)} \vec{W}\|_{\mathcal{H}|_{B(t)}} + \|\vec{F}(t)\|_{\mathcal{R}_{B(t)}^{T-t}} \lesssim \varsigma^{d/2-1}. \quad (4.13)$$

Furthermore, if $T < \infty$, then $\sigma(t) \rightarrow \infty$ and $c(t) \rightarrow \exists c_* \in \mathbb{R}^d$ as $t \rightarrow T - 0$.

Proof Let $\psi(t) := \mathcal{T}^{c(t)} \mathcal{S}^{\sigma(t)} \vec{W}$, $\vec{v}(t) := U(t)(\vec{F}(0) + R(0))$, and $\vec{w}(t) := \vec{u}(t) - \psi(0)$. Then we have

$$\square w = f'(\psi_1(0) + w) - f'(\psi_1(0)), \quad \vec{w}(0) = \vec{v}(0), \tag{4.14}$$

hence by Strichartz

$$\begin{aligned} \|\vec{w} - \vec{v}\|_{L^\infty \mathcal{H} \cap \text{St}(0,S)} &\lesssim \|w\|_{\text{St}_s(0,S)} (\|\psi(0)\|_{\text{St}_m(0,S)} + \|w\|_{\text{St}_m(0,S)})^{2^* - 2} \\ &\lesssim \|w\|_{\text{St}_s(0,S)} (|e^{\sigma(0)} S|^{1/q_m} + \|w\|_{\text{St}_m(0,S)})^{2^* - 2}, \end{aligned} \tag{4.15}$$

and $\|\vec{v}\|_{\text{St}(0,S)} \lesssim \|\vec{F}\|_{\text{St}(0,S)} + \|R(0)\|_{\mathcal{H}} \leq \varsigma$, for any $S \in (0, T)$. Choosing $\varepsilon_D \ll 1$, we deduce that as long as $0 < e^{\sigma(0)} S < \varepsilon_D$

$$\|\vec{w} - \vec{v}\|_{L^\infty \mathcal{H} \cap \text{St}(0,S)} \ll \|\vec{v}\|_{\text{St}(0,S)} \lesssim \varsigma. \tag{4.16}$$

Since $\vec{w}(t) - \vec{v}(t) = \psi(t) - \psi(0) + R(t) - U(t)R(0)$, it implies that for $0 < t < e^{-\sigma(0)} \varepsilon_D$,

$$\begin{aligned} \varsigma &\gtrsim \|\vec{w}(t) - \vec{v}(t)\|_{\mathcal{H}} + \|R(t)\|_{\mathcal{H}} + \|R(0)\|_{\mathcal{H}} \\ &\gtrsim \|\psi(t) - \psi(0)\|_{\mathcal{H}} \sim |\sigma(t) - \sigma(0)| + e^{\sigma(0)} |c(t) - c(0)|. \end{aligned} \tag{4.17}$$

Now define a time sequence inductively by $t_0 = 0$ and $t_{j+1} = t_j + e^{-\sigma(t_j)} \varepsilon_D$. Then applying the above argument from t_j yields

$$|\sigma(t) - \sigma(t_j)| + e^{\sigma(t_j)} |c(t) - c(t_j)| \lesssim \varsigma \tag{4.18}$$

for $t_j \leq t \leq t_j + e^{-\sigma(t_j)} \varepsilon_D$, and induction on j yields the desired estimate (4.12).

If $I = [0, \infty)$, let $B(t) := \{|x - c(0)| < e^{-\sigma(0)} + t/2\}$. Lemma 4.1 implies

$$\|\vec{F}(s)\|_{\mathcal{R}_{B(s)}^\infty} \lesssim \|\vec{F}\|_{L_t^\infty(s, \infty; \mathcal{H}|_{B(s)+t})} + \|F\|_{\text{St}(s, \infty)} \rightarrow 0 \tag{4.19}$$

as $s \rightarrow \infty$, while $\|\psi(s)\|_{\mathcal{H}|_{B(s)}} \lesssim S^{d/2-1}$ for large s , by (4.12) together with the explicit form of W . Thus we obtain (4.13).

If $I = [0, T)$ and $T < \infty$, then $\sigma(t) \rightarrow \infty$ as $t \nearrow T$, since otherwise there is a sequence $t_n \nearrow T$ with $\sup_n \sigma(t_n) < \infty$ and $\delta > 0$ such that

$$\sup_n \sup_{c \in \mathbb{R}^d} \|\vec{u}(t_n)\|_{\tilde{\mathcal{H}}(|x-c| < \delta)} \lesssim \varsigma. \tag{4.20}$$

Choosing $\varepsilon_D \ll \varepsilon_S$, this ensures solvability in the cone $|x - c| + |t - t_n| < \delta$ for all $c \in \mathbb{R}^d$ and all t_n , thereby extending the solution beyond T , a contradiction. Hence $\sigma(t) \rightarrow \infty$. Then (4.12) implies the convergence $c(t) \rightarrow \exists c_* \in \mathbb{R}^d$. Let $B := \{|x - c_*| < R\}$. Then as $s \nearrow T$ we have

$$\|\vec{F}(s)\|_{\mathcal{H}_B^{T-s}} \leq \|\vec{F}\|_{L^\infty(s,T;\mathcal{H}|_{B_{T-s}})} + \|\vec{F}\|_{\text{St}(s,T)} \rightarrow \|\vec{F}(T)\|_{\mathcal{H}|_B}, \quad (4.21)$$

and $\|\psi(s)\|_{\mathcal{H}|_{B^c}} \rightarrow 0$. Choosing R small enough yields (4.13). \square

The following Sobolev-type inequality implies that \mathcal{H}_B^∞ controls L^{2^*} , which is a notable difference from \mathcal{H}_B^T with $T < \infty$.

Lemma 4.3 (Time-Sobolev for the Strichartz norms) *Let $\mathbb{N} \ni d \geq 2$, $\mathbb{Z} \ni k \geq 0$ and $2 \leq q \leq r \leq \infty$. Then for any free solution v we have*

$$\|\partial_t^k v\|_{L_t^\infty(0,\infty;\dot{B}_{r,r}^{\sigma-1/q-k})} \lesssim \|v\|_{L_t^q(0,\infty;\dot{B}_{r,r}^\sigma)}. \quad (4.22)$$

In particular, if $d \geq 3$, $1/q + d/r - \sigma = d/2 - 1$ and $\sigma > 1/q$, then

$$\|\partial_t^k v\|_{L_t^\infty(0,\infty;|\nabla|^k L^{2^*})} \lesssim \|v\|_{L_t^q(0,\infty;\dot{B}_{r,2}^\sigma)}. \quad (4.23)$$

The last inequality applies to any Strichartz norm of the \dot{H}^1 scaling with the condition $1/\sigma < q \leq r$, in particular to $\text{St}_s = L^{q_s} \dot{B}_{q_s,2}^{1/2}$. Hence we have

$$\|\vec{v}\|_{L_t^\infty(0,\infty;(1 \oplus |\nabla|)L^{2^*})} \lesssim \|\vec{v}\|_{\text{St}_s(0,\infty)}. \quad (4.24)$$

Proof Let $v = \sum_{j \in \mathbb{Z}} v_j$ be a standard Littlewood-Paley decomposition in $x \in \mathbb{R}^d$ with $\text{supp } \mathcal{F}v_j \subset \{2^{j-1} < |\xi| < 2^{j+1}\}$. The wave equation and the property of the L-P decomposition imply

$$\|\ddot{v}_j\|_{L_x^r L_t^q} = \|\Delta v_j\|_{L_x^r L_t^q} \sim 2^{2j} \|v_j\|_{L_x^r L_t^q}, \quad (4.25)$$

on $(0, \infty) \times \mathbb{R}^d$. Then by the elementary interpolation inequalities

$$\|\dot{u}\|_{L^q(0,\infty)} \lesssim \|u\|_{L^q(0,\infty)}^{1/2} \|\ddot{u}\|_{L^q(0,\infty)}^{1/2}, \quad \|u\|_{L^\infty(0,\infty)} \lesssim \|u\|_{L^q(0,\infty)}^{1-1/q} \|\ddot{u}\|_{L^q(0,\infty)}^{1/q}, \quad (4.26)$$

we obtain

$$\begin{aligned} \|v_j\|_{L_x^r L_t^q} &\lesssim \|v_j\|_{L_x^r L_t^q}^{1-1/(2q)} \|\ddot{v}_j\|_{L_x^r L_t^q}^{1/(2q)} \sim 2^{j/q} \|v_j\|_{L_x^r L_t^q}, \\ \|\dot{v}_j\|_{L_x^r L_t^\infty} &\lesssim \|v_j\|_{L_x^r L_t^q}^{1/2-1/(2q)} \|\ddot{v}_j\|_{L_x^r L_t^q}^{1/2+1/(2q)} \sim 2^{j(1/q+1)} \|v_j\|_{L_x^r L_t^q}. \end{aligned} \quad (4.27)$$

Using Minkowski and Littlewood-Paley yields

$$\|v\|_{L_t^\infty \dot{B}_{r,r}^{\sigma-1/q}} \leq \|2^{j(\sigma-1/q)} v_j\|_{\ell_j^r L_x^r L_t^q} \lesssim \|2^{j\sigma} v_j\|_{\ell_j^r L_x^r L_t^q} \leq \|v\|_{L_t^q \dot{B}_{r,r}^\sigma}. \quad (4.28)$$

The estimate on the time derivatives follows in the same way. (4.23) follows from the standard Sobolev embedding for the Besov space, $\dot{B}_{r,2}^{\sigma-1/q} \subset \dot{B}_{r,r}^{\sigma-1/q} \subset L^{2^*}$.

Finally, (4.26) can be proved as follows. By the density argument, we may assume that v is nonzero, real analytic, and exponentially decaying. Then \dot{v} has at most countable number of zeros $0 \leq z_1 < z_2 < \dots$ with no accumulation. For each $z_k < z_{k+1}$ we have, denoting $[v]^q := |v|^{q-1}v$,

$$\begin{aligned} \|\dot{v}\|_{L^1(z_k, z_{k+1})}^q &= |v(z_k) - v(z_{k+1})|^q \leq 2^{q-1} |[v(z_k)]^q - [v(z_{k+1})]^q| \\ &\leq 2^{q-1} \int_{z_k}^{z_{k+1}} q|v|^{q-1}|\dot{v}|dt \leq 2^{q-1}q\|v\|_{L^q(z_k, z_{k+1})}^{q-1}\|\dot{v}\|_{L^q(z_k, z_{k+1})}, \end{aligned} \tag{4.29}$$

and similarly,

$$\|\dot{v}\|_{L^\infty(z_k, z_{k+1})}^q \leq \text{Re} \int_{z_k}^{z_{k+1}} q[\dot{v}]^{q-1}\bar{v}dt \leq q\|\dot{v}\|_{L^q(z_k, z_{k+1})}^{q-1}\|\dot{v}\|_{L^q(z_k, z_{k+1})}. \tag{4.30}$$

The same argument yields the second inequality of (4.26). Interpolating the above two estimates by Hölder

$$\|\dot{v}\|_{L^q(z_k, z_{k+1})} \leq 2^{(q-1)/q^2}q^{1/q}\|v\|_{L^q(z_k, z_{k+1})}^{(q-1)/q^2}\|\dot{v}\|_{L^q(z_k, z_{k+1})}^{1/q^2+(1-1/q)^2}\|\dot{v}\|_{L^q(z_k, z_{k+1})}^{(q-1)/q^2}, \tag{4.31}$$

and so $\|\dot{v}\|_{L^q(z_k, z_{k+1})} \leq [2q^{1-1/q}\|v\|_{L^q(z_k, z_{k+1})}\|\dot{v}\|_{L^q(z_k, z_{k+1})}]^{1/2}$. Taking the ℓ^q sum over k , we obtain the first inequality of (4.26). \square

Remark 4.1 It is obvious from the homogeneous nature that the above lemma fails on any bounded set in \mathbb{R} for any $q < \infty$. If such an inequality would hold, then it must be uniform for the rescaling $u(t, x) \mapsto u(\lambda t, \lambda x)$, but the $L_t^q(I)$ norm decays as I shrinks to a point $t_0 \in \mathbb{R}$, while the $L_t^\infty(I)$ norm converges to the value at t_0 .

The following lemma allows us to detach exterior radiation which is small in \mathcal{R}_R^T from any solution of CW.

Lemma 4.4 (Detaching lemma) *Let $B \in \mathcal{B}_d$, $\tilde{T} \geq T > 0$, and $u \in \text{Solution}([0, T]) \cup \text{Solution}([0, T])$ satisfy $\|\tilde{u}(0)\|_{\mathcal{R}_B^{\tilde{T}}} = \varsigma < \varepsilon_S$. Then*

- (I) *There are a free solution v , and two strong solutions $u^{\mathbf{x}}$ and $u^{\mathbf{d}}$ of (CW), defined on $[0, \tilde{T}]$ and on $[0, T]$ respectively, satisfying*

$$\begin{aligned} \tilde{v}(0) &= \tilde{u}(0) \text{ on } B^{\mathbb{G}}, \quad \|\tilde{v}\|_{L_t^\infty(0, \tilde{T}; \mathcal{H}|_{B_{+t}}) \cap \text{St}(0, \tilde{T})} \lesssim \varsigma, \\ \tilde{u}^{\mathbf{x}}(t) &= \tilde{u}(t) \text{ on } (B_{+t})^{\mathbb{G}}, \quad \|\tilde{u}^{\mathbf{x}} - \tilde{v}\|_{(L^\infty \mathcal{H} \cap \text{St})(0, \tilde{T})} \lesssim \varsigma^{2^* - 1}, \\ \tilde{u}^{\mathbf{d}}(t) &= \tilde{u}(t) \text{ on } B_{+t}, \quad \|\tilde{u} - \tilde{u}^{\mathbf{d}} - \tilde{v}\|_{(L^\infty \mathcal{H} \cap \text{St})(0, T)} \lesssim \varsigma, \end{aligned} \tag{4.32}$$

and $\|\tilde{v}(0)\|_{\mathcal{H}} \lesssim \|\tilde{u}(0)\|_{\mathcal{H}|_{B^{\mathbb{G}}}}$. More precisely, there is $w \in C([0, \tilde{T}]; \mathcal{H})$ such that $\tilde{u} = \tilde{u}^{\mathbf{d}} + w$ for $0 \leq t < T$ and $\|w - \tilde{v}\|_{(L^\infty \mathcal{H} \cap \text{St})(0, \tilde{T})} \lesssim \varsigma$. If $\tilde{T} = \infty$, then

$$\|w\|_{L^\infty(0, \infty; (1 \oplus |\nabla|)L^{2^*}(\mathbb{R}^d))} + \|\tilde{u}^{\mathbf{x}}\|_{L^\infty(0, \infty; (1 \oplus |\nabla|)L^{2^*}(\mathbb{R}^d))} \lesssim \varsigma. \tag{4.33}$$

- (II) *There exists $\{\bar{u}^\theta\}_{0 \leq \theta \leq 1}$ in $S_{\text{olution}}([0, T])$ or $S_{\text{olution}}([0, T])$, which is C^1 in θ , such that $\bar{u}^0 = \bar{u}$, $\bar{u}^1 = \bar{u}^{\mathbf{d}}$, $\bar{u}^\theta = \bar{u}$ on $B_{+t}(0, T)$, $\|\bar{u}^\theta(0)\|_{\mathcal{R}_B^{\tilde{T}}} \lesssim \varsigma$, and \bar{u}^θ satisfies (I) for the fixed $u^{\mathbf{d}}$ and some θ -dependent $u^{\mathbf{x}}$ and v for all $\theta \in [0, 1]$.*
- (III) *Although such $u^{\mathbf{d}}$ and $u^{\mathbf{x}}$ in (I) are not unique, we can define a C^1 map $A : \bar{u}(0) \mapsto \bar{u}^{\mathbf{d}}(0) \in \mathcal{H}$ locally around each $\bar{u}(0)$ and fixed B . Moreover, it satisfies*

$$\|A(\varphi^0) - A(\varphi^1) - (\varphi^0 - \varphi^1)\|_{\mathcal{H}} \lesssim \|\varphi^0 - \varphi^1\|_{\mathcal{H} \setminus B^{\mathbb{C}}}. \tag{4.34}$$

Note that no energy bound is required on u , while the condition in $\mathcal{R}_B^{\tilde{T}}$ can be satisfied either by localization as in (4.5) or by dispersion as in (4.10), which is useful respectively for concentrating blow-up and for scattering solutions.

Proof The definition of $\mathcal{R}_B^{\tilde{T}}$ yields a free solution \bar{v} such that $\bar{v}(0) = \bar{u}(0)$ on $B^{\mathbb{C}}$ and

$$\|\bar{u}(0)\|_{\mathcal{R}_B^{\tilde{T}}} = \|\bar{v}\|_{L_t^\infty(0, \tilde{T}; \mathcal{H} \setminus B_{+t}) \cap \text{St}(0, \tilde{T})} \lesssim \|\bar{v}(0)\|_{\mathcal{H}} \sim \|\bar{u}(0)\|_{\mathcal{H} \setminus B^{\mathbb{C}}}. \tag{4.35}$$

Since $\varsigma < \varepsilon_S$, there is a unique $u^{\mathbf{x}} \in S_{\text{olution}}([0, \tilde{T}])$ such that

$$\bar{u}^{\mathbf{x}}(0) = \bar{v}(0), \quad \|\bar{u}^{\mathbf{x}} - \bar{v}\|_{(L^\infty \mathcal{H} \cap \text{St})(0, \tilde{T})} \lesssim \|v\|_{\text{St}(0, \tilde{T})}^{2^*-1} = \varsigma^{2^*-1}, \tag{4.36}$$

which, together with the above estimate on v , implies that

$$\|\bar{u}^{\mathbf{x}}\|_{L_t^\infty(0, \tilde{T}; \mathcal{H} \setminus B_{+t}) \cap \text{St}(0, \tilde{T})} \sim \varsigma. \tag{4.37}$$

In addition, if $\tilde{T} = \infty$ then combining the above with (4.24) yields (4.33) for $u^{\mathbf{x}}$. The propagation speed of CW implies that $\bar{u}^{\mathbf{x}} = \bar{u}$ on $(B_{+t})^{\mathbb{C}} = (B^{\mathbb{C}})_{-t}$. If $\tilde{T} < \infty$, then let w be the solution of

$$(\partial_t^2 - \Delta)w = f'(u^{\mathbf{x}}) - f'(u^{\mathbf{x}} - w), \tag{4.38}$$

with $\bar{w}(\tilde{T}) = (1 - X_{B_{+\tilde{T}}})u^{\mathbf{x}}(\tilde{T})$, where $X_{B_{+\tilde{T}}}$ is the extension operator for $B_{+\tilde{T}}$. Then

$$\|\bar{w}(\tilde{T}) - \bar{v}(\tilde{T})\|_{\mathcal{H}} \leq \|\bar{u}^{\mathbf{x}}(\tilde{T}) - \bar{v}(\tilde{T})\|_{\mathcal{H}} + \|X_{B_{+\tilde{T}}}\bar{v}(\tilde{T})\|_{\mathcal{H}} \lesssim \varsigma, \tag{4.39}$$

and so $\|U(t - \tilde{T})\bar{w}(\tilde{T})\|_{\text{St}(0, \tilde{T})} \lesssim \varsigma$ by the Strichartz estimate. Also we have

$$\begin{aligned} \|\bar{w} - U(t - \tilde{T})\bar{w}(\tilde{T})\|_{\text{St}} &\lesssim \|\square w\|_{\text{St}_s^*} \lesssim \|w\|_{\text{St}} (\|u^{\mathbf{x}}\|_{\text{St}} + \|w\|_{\text{St}})^{2^*-2} \\ &\lesssim \varsigma \|w\|_{\text{St}} + \|w\|_{\text{St}}^{2^*-1}, \end{aligned} \tag{4.40}$$

thereby we obtain $w \in C([0, \tilde{T}]; \mathcal{H})$ satisfying $\|\bar{w} - \bar{v}\|_{L^\infty(0, \tilde{T}; \mathcal{H})} + \|w\|_{\text{St}(0, \tilde{T})} \lesssim \varsigma$ as well as $\bar{w} = 0$ on $B_{+t}([0, \tilde{T}])$ by the finite propagation speed. Let $u^{\mathbf{d}} := u - w \in C([0, T]; \mathcal{H})$. Then we have, for $0 < t < T$, $\bar{u}^{\mathbf{d}} = \bar{u}$ on B_{+t} and

$$\square u^{\mathbf{d}} = f'(u) - f'(u^{\mathbf{x}}) + f'(u^{\mathbf{x}} - w) = \begin{cases} f'(u) = f'(u^{\mathbf{d}}) & \text{on } B_{+t} \\ f'(u - w) = f'(u^{\mathbf{d}}) & \text{on } (B_{+t})^{\mathbb{C}} \end{cases}, \tag{4.41}$$

since $\bar{w} = 0$ on B_{+t} and $\bar{u} = \bar{u}^{\mathbf{x}}$ on $(B_{+t})^{\mathbb{C}}$. $\bar{u}^{\mathbf{d}} - \bar{u} + \bar{v} = \bar{v} - \bar{w}$ has been already estimated above.

To define u^θ in (II), let w^θ be the solution of (4.38) with $\bar{w}^\theta(\tilde{T}) = \theta \bar{w}(\tilde{T})$ and let $u^\theta := u - w^\theta$. Then obviously $w^0 = 0$, $w^1 = w$, $\bar{w} = 0$ on $B_{+t}(0, \tilde{T})$, and so $\square u^\theta = f'(u^\theta)$ in the same way as (4.41). The same estimate as above yields $\|w^\theta - \theta \bar{w}\|_{L^\infty \mathcal{H} \cap \text{St}(0, \tilde{T})} \lesssim \varsigma$, and hence $\|\bar{u}^\theta(0) - (1 - \theta)\bar{u}(0) - \theta \bar{u}^{\mathbf{d}}(0)\|_{\mathcal{H}} \lesssim \varsigma$, and $\|\bar{u}^\theta(0) - \bar{u}(0) + \theta v(0)\|_{\mathcal{H}} \lesssim \varsigma$, which implies that

$$\|\bar{u}^\theta(0)\|_{\mathcal{R}_B^{\tilde{T}}} \leq \|\bar{u}(0)\|_{\mathcal{R}_B^{\tilde{T}}} + \theta \|\bar{v}\|_{L_t^\infty(0, \tilde{T}; \mathcal{H}|_{B_{+t}}) \cap \text{St}(0, \tilde{T})} + C\varsigma \lesssim \varsigma. \tag{4.42}$$

Hence \bar{u}^θ satisfies (I) with the above constructed $u^{\mathbf{d}}$, the free solution $\bar{v}^\theta(t) := \bar{v}(t) + U(t)[\bar{u}^\theta(0) - \bar{u}(0)]$ and the associated nonlinear solution $u^{\mathbf{x}}$ (dependent on θ).

In the case $\tilde{T} = \infty$, we define a sequence $\bar{w}_n \in C([0, n]; \mathcal{H})$ with $\tilde{T} = n$ as above. Then the uniform bound allows us to take a weak limit along a subsequence to $w \in C([0, \infty); \mathcal{H}) \cap \text{St}(0, \infty)$ solving (4.38), the estimates and $\bar{w} = 0$ on $B_{+t}(0, \infty)$. For (II), let w^θ be the solution of the integral equation

$$\bar{w} = \int_{\infty}^t U(t - t')(0, f'(u^{\mathbf{x}} + w) - f'(u^{\mathbf{x}}))dt' \tag{4.43}$$

obtained by the iteration starting from θw . Then $u^\theta := u - w^\theta$ satisfies the desired properties, which is seen by the same argument as above.

To define the map $A : \bar{u}(0) \mapsto \bar{u}^{\mathbf{d}}(0)$ in (III), we perturb $\bar{u}(0)$ around some fixed $\bar{u}^0(0) \in \mathcal{H}$ satisfying the assumption. Let \bar{v}^0 be the free solution chosen as above for \bar{u}^0 . For $\bar{u}(0) \in \mathcal{H}$ close to $\bar{u}^0(0)$, we have

$$\|\bar{u}(0)\|_{\mathcal{R}_B^{\tilde{T}}} \leq \|\bar{u}^0(0)\|_{\mathcal{R}_B^{\tilde{T}}} + C\|\bar{u}(0) - \bar{u}^0(0)\|_{\mathcal{H}} < \varepsilon_S. \tag{4.44}$$

We choose the free solution \bar{v} for \bar{u} as a perturbation from \bar{v}^0 , putting

$$\bar{v}(0) := \bar{v}^0(0) + X_{B^{\mathbb{C}}}(\bar{u}(0) - \bar{u}^0(0)). \tag{4.45}$$

Then in the same way as above, if $\tilde{T} < \infty$, let $\bar{u}^{\mathbf{x}} \in \text{Solution}([0, \tilde{T}])$ with $\bar{u}^{\mathbf{x}}(0) = \bar{v}(0)$, let w be the solution of (4.38), and let $\bar{u}^{\mathbf{d}} = \bar{u} - \bar{w}$. By the Strichartz estimate, we see that the maps $\bar{u}(0) \mapsto \bar{v}(0) = \bar{u}^{\mathbf{x}}(0) \mapsto \bar{u}^{\mathbf{x}}(\tilde{T}) \mapsto \bar{w}(\tilde{T}) \mapsto \bar{w}(0) \mapsto \bar{u}^{\mathbf{d}}(0)$ are C^1 , where $\bar{u}^{\mathbf{x}}$ and \bar{w} are Lipschitz with respect to $\bar{u}(0) \in \mathcal{H}|_{B^{\mathbb{C}}}$, leading to (4.34).

In the case $\tilde{T} = \infty$, we also fix $\bar{w}^0 \in C([0, \infty); \mathcal{H})$ for u^0 , and then let w be the solution of (4.43) obtained by the iteration starting from w^0 . Then $w \in C([0, \infty); \mathcal{H})$ solves $\square w = f'(u^{\mathbf{x}} + w) - f'(u^{\mathbf{x}})$ on $t > 0$, $\bar{w} = 0$ on B_{+t} , and

$$\|\bar{w} - \bar{w}^0\|_{L^\infty \mathcal{H} \cap \text{St}(0, \infty)} \lesssim \|\bar{v}(0) - \bar{v}^0(0)\|_{\mathcal{H}}. \tag{4.46}$$

Moreover, the maps $\vec{u}(0) \mapsto \vec{u}^{\mathbf{x}}(0) \mapsto \vec{w}(0) \mapsto \vec{u}^{\mathbf{d}}(0)$ are C^1 , where $\vec{u}^{\mathbf{x}}$ and \vec{w} are Lipschitz with respect to $\vec{u}(0) \in \mathcal{H}|_B \mathbb{G}$, leading to (4.34). \square

The following lemma is crucial to show that the ground state component is small in the region where the solution is dispersive.

Lemma 4.5 (Reverse Sobolev for the ground state) *For any $d \geq 3$, there exists $C = C(d) > 0$ such that for any $B \in \mathcal{B}_d$,*

$$\|\vec{W}\|_{\tilde{\mathcal{H}}(B)} + \|g_{\pm}\|_{\tilde{\mathcal{H}}(B)} \leq C\|W\|_{L^{2^*}(B)}, \quad \|g_{\pm}\|_{\mathcal{H}|_B} \leq C\|\vec{W}\|_{\mathcal{H}|_B}. \tag{4.47}$$

Remark 4.2 This lemma obviously fails for any other stationary solutions, since they have indefinite sign. To see that, concentrate B at any zero point.

Proof For the first estimate, we may assume that $\|W\|_{L^{2^*}(B)} < 1$, since the left hand side is uniformly bounded. Let $R := \|W\|_{L^{2^*}(B)}^{-2/(d-2)} > 1$. Since $W \sim \langle x \rangle^{2-d}$ and $|\nabla W| \sim |x| \langle x \rangle^{-d}$, we have

$$\begin{aligned} \|\nabla W\|_{L^2(B)}^2 &\leq \int_{B \cap \{|x| < R\}} |\nabla W|^2 dx + \int_{|x| > R} |\nabla W|^2 dx \\ &\lesssim \int_B R^2 |W|^{2^*} dx + R^{2-d} \sim R^{2-d} = \|W\|_{L^{2^*}(B)}^2. \end{aligned} \tag{4.48}$$

The estimate on g_{\pm} follows from that $\rho/W \in L^\infty \cap W^{1,d}(\mathbb{R}^d)$. Indeed, let $\chi := \rho/W$ and let $\varphi = \vec{W}$ on B . Then we have $g_{\pm} = (2k)^{-1/2}(1, \pm k)\chi\varphi_1$ on B , and

$$\|(1, \pm k)\chi\varphi_1\|_{\mathcal{H}} \leq \|\chi\|_{L^\infty} \|\nabla\varphi_1\|_2 + (\|\nabla\chi\|_{L^d} + \|\chi_1\|_{L^d}) \|\varphi_1\|_{L^{2^*}} \lesssim \|\varphi\|_{\mathcal{H}}. \tag{4.49}$$

Hence $\|g_{\pm}\|_{\mathcal{H}|_B} \lesssim \|\vec{W}\|_{\mathcal{H}|_B}$. The estimate in $\tilde{\mathcal{H}}(B)$ is similar. \square

5 Center-stable manifold with large radiation

Now we can extend the center-stable manifold by adding large radiation. Fix $\varsigma_m > 0$ such that $\varsigma_m \ll \delta_m$. Let \mathcal{M}_3 be the totality of $\vec{u}(0) \in \mathcal{H}$ such that for the solution u we can apply the detaching Lemma 4.4 with

$$\tilde{T} = \infty, \quad \vec{u}^{\mathbf{d}}(0) \in \mathcal{M}_0, \quad \varsigma + \|\mathcal{F}_{\vec{u}^{\mathbf{d}}(0)} \vec{W}\|_{\mathcal{H}|_B \mathbb{G}} < \varsigma_m. \tag{5.1}$$

Then $\mathcal{M}_0 \subset \mathcal{M}_3$ by using the trivial case $B = \mathbb{R}^d$ and $u^{\mathbf{d}} = u$. The invariance of \mathcal{M}_3 for \mathcal{T}, \mathcal{S} is inherited from \mathcal{M}_0 , which is also clear from the definition.

For each point $\varphi \in \mathcal{M}_3$, Lemma 4.4 gives a neighborhood $O \ni \varphi$ and a C^1 map $A : O \ni \vec{u}(0) \mapsto \vec{u}^{\mathbf{d}}(0) \in \mathcal{H}$ such that $A(\varphi) \in \mathcal{M}_0$. Reducing O if necessary, we may assume that $A(O)$ is within the domain of M_+ defined in (2.11), and that the last condition of (5.1) holds all over O .

Then we have $\mathcal{M}_3 \cap O = (M_+ \circ A)^{-1}(0)$. Indeed \supset is clear from the definition of \mathcal{M}_3 and M_+ . If $M_+(A(\psi)) > 0$, then the solution u^d starting from $A(\psi)$ blows up in some finite $T > 0$, so does the solution u starting from ψ , because of (4.32),

$$\begin{aligned} \text{dist}_{L^{2*}}(u(t), S_{\text{tatic}}(W)_1) &\geq \text{dist}_{L^{2*}}(u^d(t), S_{\text{tatic}}(W)_1) - \|u(t) - u^d(t)\|_{L^{2*}} \\ &\gg \delta_m - O(\zeta) \sim \delta_m, \end{aligned} \tag{5.2}$$

for t close to T . On the other hand, if $\vec{u}(0) \in \mathcal{M}_3$ with $\vec{u} \in S_{\text{olution}}([0, T))$, then

$$\begin{aligned} \text{dist}_{L^{2*}}(u(t), S_{\text{tatic}}(W)_1) &\leq \text{dist}_{L^{2*}}(u^d(t), S_{\text{tatic}}(W)_1) + \|u(t) - u^d(t)\|_{L^{2*}} \\ &\lesssim \delta_m + \zeta \sim \delta_m \end{aligned} \tag{5.3}$$

for t close to T . Hence $M_+(A(\psi)) > 0$ implies that $\psi \notin \mathcal{M}_3$. If $M_+(A(\psi)) < 0$, then (4.32) with Strichartz implies that the solution u starting from ψ also scatters to 0, contradicting (5.3), and so $\psi \notin \mathcal{M}_3$.

In order to conclude that \mathcal{M}_3 is a C^1 manifold of codimension 1, it now suffices to show that the C^1 functional $M_+ \circ A$ does not degenerate on its zero set \mathcal{M}_3 . Indeed, if $M_+(A(\psi)) = 0$ then $\partial_h M_+(A(\psi + h\mathcal{T}_{A(\psi)}g_+)) \sim 1$, because by the Lipschitz property of A (4.34) and m_+ (2.7), and the last condition of (5.1) together with (4.47), we have

$$\begin{aligned} A(\psi + h\mathcal{T}_{A(\psi)}g_+) &= A(\psi) + h[\mathcal{T}_{A(\psi)}g_+ + O(\zeta_m; \mathcal{H})], \\ M_+(A(\psi + h\mathcal{T}_{A(\psi)}g_+)) &= h + O(h\zeta_m). \end{aligned} \tag{5.4}$$

By Lemma 4.4(II), we can connect each $\varphi \in \mathcal{M}_3$ with some $\psi \in \mathcal{M}_0$ by a C^1 curve, which is included in an enlarged \mathcal{M}_3 for which the last bound in (5.1) is replaced with $C\zeta_m$ for some constant $C > 1$. Including those curves connecting \mathcal{M}_3 to \mathcal{M}_0 , we obtain a slightly bigger manifold $\widetilde{\mathcal{M}}_3$, which is C^1 and connected with codimension 1.

Let \mathcal{M}_4 be the maximal evolution of $\widetilde{\mathcal{M}}_3$ (in the same way as we define \mathcal{M}_1 from \mathcal{M}_0). Then \mathcal{M}_4 is a connected C^1 manifold of codimension 1, which is invariant by the flow, \mathcal{T} and \mathcal{S} , and $\mathcal{M}_4 \supset \mathcal{M}_1 \cup \widetilde{\mathcal{M}}_3$. Every solution $\vec{u} \in S_{\text{olution}}([0, T))$ on \mathcal{M}_4 satisfies

$$\limsup_{t \nearrow T} \text{dist}_{(1 \oplus |\nabla|)L^{2*}}(\vec{u}(t), S_{\text{tatic}}(W)) \lesssim \delta_m. \tag{5.5}$$

Around each point on \mathcal{M}_4 , there is a small open ball which is split into two open sets by \mathcal{M}_4 , such that all the solutions starting from one of them blow up in finite time, near which time

$$\text{dist}_{L^{2*}}(u(t), S_{\text{tatic}}(W)_1) \gg \delta_m, \tag{5.6}$$

and all those starting from the other scatter to 0 as $t \rightarrow \infty$.

On the other hand, if $\vec{u} \in S_{\text{olution}}([0, \infty))$ scatters to $S_{\text{tatic}}(W)$, namely

$$\exists \varphi \in \mathcal{H}, \quad d_W(\vec{u}(t) - U(t)\varphi) \rightarrow 0 \quad (t \rightarrow \infty), \tag{5.7}$$

then $\vec{u}(0) \in \mathcal{M}_4$. This is because Lemma 4.2 implies that we can detach the free radiation $U(t)\varphi$ at some large $t = T$, so that $\vec{u}^d(t) = \mathcal{T}_{\vec{u}(t)}\vec{W} + O(\zeta)$ in \mathcal{H} for all $t \geq T$ with $\zeta \ll \zeta_m \ll \delta_m$. The last condition of (5.1) is ensured by (4.13).

Finally, let \mathcal{M}_5 be the Lorentz extension of \mathcal{M}_4 , defined in the same way as for \mathcal{M}_2 from \mathcal{M}_1 . Note that all the solutions on \mathcal{M}_4 have the space-time maximal regions as in (3.29) in the case of blow-up, since the remainder u^x is globally small in the Strichartz norms. Thus we obtain a connected C^1 manifold $\mathcal{M}_5 \supset \text{Soliton}(W) \cup \mathcal{M}_4 \cup \mathcal{M}_2$ with codimension 1 in \mathcal{H} , which is invariant by the flow, \mathcal{T} , \mathcal{S} and the Lorentz transform. If $\vec{u} \in \text{S}_{\text{olution}}([0, \infty))$ satisfies (1.21), then the scaling invariance and dispersion of the free wave v implies that

$$P(\vec{u}) = \lim_{t \rightarrow \infty} P(\vec{W}_{0,0,p(t)}) + P(\vec{v}) = \lim_{t \rightarrow \infty} p(t)E(W) + P(\vec{v}). \tag{5.8}$$

Hence $p(t)$ converges to some $p_* \in \mathbb{R}^d$, and then $\vec{W}_{\lambda(t),c(t),p(t)} - \vec{W}_{\lambda(t),c(t),p_*} \rightarrow 0$ in \mathcal{H} . Take a Lorentz transform which maps $\vec{W}_{0,0,p_*}$ to \vec{W} , and apply it to \vec{u} . Then we obtain another global solution satisfying (1.21) with $p(t) \equiv 0$, namely scattering to $\text{S}_{\text{tatic}}(W)$, and so belonging to \mathcal{M}_4 . Hence u is on \mathcal{M}_5 .

Since each Lorentz transform defines a local C^1 diffeo around each solution, there is a neighborhood of \mathcal{M}_5 transformed from a neighborhood of \mathcal{M}_4 , such that all solutions starting off the manifold within the neighborhood either scatter to 0 as $t \rightarrow \infty$ or blow up in $t > 0$ away from a bigger neighborhood. Thus we obtain Theorem 1.3.

6 One-pass theorem with large radiation

In this section, we derive one-pass theorems which allow arbitrarily large radiation. For $\varphi \in \mathcal{H}$, we define the ‘‘radiative distance’’ $d_{\mathcal{R}}$ to the ground states for any $\varphi \in \mathcal{H}$ by

$$d_{\mathcal{R}}(\varphi) := \inf_{B \in \mathcal{B}_d, \psi \in \pm \text{S}_{\text{tatic}}(W)} \|\varphi - \psi\|_{\mathcal{H} \downarrow B} + \|\varphi\|_{\mathcal{R}_B^\infty} + \|\psi\|_{\mathcal{H} \downarrow B^c}. \tag{6.1}$$

Obviously $d_{\mathcal{R}} : \mathcal{H} \rightarrow [0, \infty)$ is Lipschitz continuous, and $d_{\mathcal{R}}(\mathcal{T}^c \mathcal{S}^\sigma \varphi) = d_{\mathcal{R}}(\varphi)$. It is not invariant for the time inversion $\varphi \mapsto \varphi^\dagger$. Taking $B = \mathbb{R}^d$ yields

$$d_{\mathcal{R}}(\varphi) \leq \text{dist}_W(\varphi) \sim d_W(\varphi). \tag{6.2}$$

The embeddings (4.24) and $\mathcal{H} \subset (1 \oplus |\nabla|)L^{2^*}$ imply

$$\text{dist}_{(1 \oplus |\nabla|)L^{2^*}}(\varphi, \pm \text{S}_{\text{tatic}}(W)) \lesssim d_{\mathcal{R}}(\varphi). \tag{6.3}$$

By Lemma 4.2, we immediately obtain

Lemma 6.1 *Under the same assumption as in Lemma 4.2 with $I = [0, \infty)$,*

$$\lim_{t \rightarrow \infty} d_{\mathcal{R}}(\vec{u}(t)) \lesssim \zeta^{d/2-1} + \zeta. \tag{6.4}$$

In particular, if $\vec{u} \in \text{S}_{\text{olution}}([0, \infty))$ scatters to the ground states (5.7), then

$$d_{\mathcal{R}}(\vec{u}(t)) + t^{-1}(e^{-\tilde{\sigma}(\vec{u}(t))} + |\tilde{c}(\vec{u}(t))|) \rightarrow 0 \quad (t \rightarrow \infty). \tag{6.5}$$

Proof Combine (4.13) with $d_{\mathcal{R}}(\vec{u}(t)) \lesssim \|\vec{F}(t)\|_{\mathcal{R}_{B(t)}^\infty} + \|\psi(t)\|_{\mathcal{H}|_{B(t)}^{\mathbb{C}}} + \|R(t)\|_{\mathcal{H}}$. □

Smallness of the radiative distance enables the detaching.

Lemma 6.2 *Let $\vec{u} \in \text{S}_{\text{olution}}([0, T])$ satisfy $d_{\mathcal{R}}(\vec{u}(0)) = \varsigma < \varepsilon_S$. Then there exist $B \in \mathcal{B}_d$, a free solution $v, \vec{u}^{\mathbf{x}} \in \text{S}_{\text{olution}}([0, \infty))$ and $\vec{u}^{\mathbf{d}} \in \text{S}_{\text{olution}}([0, T])$, satisfying $\vec{v}(0) = \vec{u}(0)$ on $B^{\mathbb{C}}$, $\vec{u}^{\mathbf{x}} = \vec{u}$ on $(B_{+t})^{\mathbb{C}}$, $\vec{u}^{\mathbf{d}} = \vec{u}$ on B_{+t} , and*

$$\begin{aligned} \|\vec{v}\|_{L_t^\infty(0, \infty; \mathcal{H}|_{B_{+t}}) \cap \text{St}(0, \infty)} + \|\vec{u} - \vec{u}^{\mathbf{d}} - \vec{v}\|_{L^\infty \mathcal{H} \cap \text{St}(0, T)} &\lesssim \varsigma, \\ \|\vec{u}^{\mathbf{x}} - \vec{v}\|_{L^\infty \mathcal{H} \cap \text{St}(0, \infty)} &\lesssim \varsigma^{2^*-1}, \quad \|\vec{u}^{\mathbf{d}}(t)\|_{L_t^\infty(0, T; \mathcal{H}|_{B_{+t}})} \lesssim \varsigma. \end{aligned} \tag{6.6}$$

Moreover, for $0 \leq t < T$ we have

$$d_{\mathcal{R}}(\vec{u}(t)) \lesssim \varsigma + d_W(\vec{u}^{\mathbf{d}}(t)). \tag{6.7}$$

Proof By definition of $d_{\mathcal{R}}$, there exist $B \in \mathcal{B}_d$ and $\psi \in \pm \text{S}_{\text{tatic}}(W)$ such that

$$\varsigma \gtrsim \|\vec{u}(0) - \psi\|_{\mathcal{H}|_B} + \|\vec{u}(0)\|_{\mathcal{R}_B^\infty} + \|\psi\|_{\mathcal{H}|_{B^{\mathbb{C}}}} < \varepsilon_S. \tag{6.8}$$

Applying Lemma 4.4 with $\vec{T} = \infty$ yields $v, u^{\mathbf{x}}, u^{\mathbf{d}}$ and (6.6), where the last inequality follows from the others

$$\|\vec{u}^{\mathbf{d}}(t)\|_{\mathcal{H}|_{B_{+t}^{\mathbb{C}}}} \leq \|\vec{u}^{\mathbf{x}}(t) - \vec{v}(t)\|_{\mathcal{H}|_{B_{+t}^{\mathbb{C}}}} + \|\vec{u}^{\mathbf{d}}(t) - \vec{u}(t) + \vec{v}(t)\|_{\mathcal{H}} \lesssim \varsigma. \tag{6.9}$$

Let $\psi(t) : [0, T] \rightarrow \pm \text{S}_{\text{tatic}}(W)$ such that $d_W(\vec{u}^{\mathbf{d}}(t)) \sim \|\vec{u}^{\mathbf{d}}(t) - \psi(t)\|_{\mathcal{H}}$. Then

$$\begin{aligned} \|\vec{u}(s) - \psi\|_{\mathcal{H}|_{B_{+s}}} &\leq \|\vec{u}^{\mathbf{d}}(s) - \psi\|_{\mathcal{H}} \sim d_W(\vec{u}^{\mathbf{d}}(s)), \\ \|\psi\|_{\mathcal{H}|_{(B_{+s})^{\mathbb{C}}}} &\leq \|\vec{u}^{\mathbf{d}}(s) - \psi\|_{\mathcal{H}} + \|u^{\mathbf{d}}(s)\|_{\mathcal{H}|_{(B_{+s})^{\mathbb{C}}}} \lesssim d_W(\vec{u}^{\mathbf{d}}(s)) + \varsigma, \\ \|\vec{u}(s)\|_{\mathcal{R}_{B_{+s}}^\infty} &\leq \|U(t-s)\vec{u}^{\mathbf{x}}(s)\|_{L_t^\infty(s, \infty; \mathcal{H}|_{B_{+t}}) \cap \text{St}(s, \infty)} \\ &\lesssim \|\vec{v}\|_{L_t^\infty(s, \infty; \mathcal{H}|_{B_{+t}}) \cap \text{St}(s, \infty)} + \|\vec{v}(s) - \vec{u}^{\mathbf{x}}(s)\|_{\mathcal{H}} \lesssim \varsigma. \end{aligned} \tag{6.10}$$

Gathering these three estimates, we obtain (6.7). □

We are now ready to prove the first one-pass theorem in the radiative distance.

Theorem 6.3 *There exist constants $C_* > 1 > \varsigma_* > 0$ such that if $\vec{u} \in \text{S}_{\text{olution}}([0, T])$ satisfies*

$$\max(d_{\mathcal{R}}(\vec{u}(0)), d_{\mathcal{R}}(\vec{u}(T))) =: \varsigma \leq \varsigma_* \ll \delta_*, \tag{6.11}$$

then there is $B \in \mathcal{B}_d$ such that the conclusion of Lemma 4.4(I) holds and

$$\|d_{\mathcal{R}}(\vec{u})\|_{L_t^\infty(0, T)} + \|d_W(\vec{u}^{\mathbf{d}})\|_{L_t^\infty(0, T)} \leq C_* \varsigma. \tag{6.12}$$

On the other hand, if $\vec{u} \in \text{Solution}([0, T])$ satisfies

$$d_{\mathcal{R}}(\vec{u}(0)) =: \varsigma \leq \varsigma_*, \quad d_{\mathcal{R}}(\vec{u}(t_0)) > C_*\varsigma \tag{6.13}$$

at some $t_0 \in (0, T)$, then there is $t_1 \in [t_0, T)$ such that

$$d_{\mathcal{R}}(\vec{u}(t_1)) = \inf_{t_1 \leq t < T} d_{\mathcal{R}}(\vec{u}(t)) \geq \max(\varsigma, \varsigma_*/C_*). \tag{6.14}$$

Moreover, if $K_{B_{+t_1}}(u(t_1)) < 0$ then $T < \infty$ and

$$\liminf_{t \nearrow T} \text{dist}_{L^{2^*}}(u(t), \text{Static}(W)_1) \gtrsim \delta_*, \tag{6.15}$$

otherwise $T = \infty$ and u scatters to 0 as $t \rightarrow \infty$.

Proof We will reduce it to [14] by the detaching Lemma 4.4. Choose $\varsigma_* < \varepsilon_S$ and let B, v, u^x and u^d be as in the above lemma, with $\psi \in \pm \text{Static}(W)$ satisfying (6.8). Combining it with (6.6) yields

$$\|\vec{u}^d(0) - \psi\|_{\mathcal{H}} \leq \|\vec{u}(0) - \psi\|_{\mathcal{H}(B)} + \|\psi\|_{\mathcal{H}(B^c)} + \|\vec{u}^d(0)\|_{\mathcal{H}(B^c)} \lesssim \varsigma, \tag{6.16}$$

which implies $E(\vec{u}^d) = E(W) + O(\varsigma^2)$ since $E' = 0$ on any static solution.

The same argument at $t = T$ yields $\tilde{B} \in \mathcal{B}_d$ and $\tilde{\psi} \in \pm \text{Static}(W)$ such that

$$\|\vec{u}(T) - \tilde{\psi}\|_{\mathcal{H}|\tilde{B}} + \|\vec{u}(T)\|_{\mathcal{R}^\infty_{\tilde{B}}} + \|\tilde{\psi}\|_{\mathcal{H}|\tilde{B}^c} + \|u(T)\|_{L^{2^*}(\tilde{B}^c)} \lesssim \varsigma. \tag{6.17}$$

Let $\hat{B} := B_{+T}$. Using the reverse Sobolev (4.47), we obtain

$$\|\nabla \tilde{\psi}_1\|_{L^2(\tilde{B} \setminus \hat{B})} \lesssim \|\tilde{\psi}_1\|_{L^{2^*}(\tilde{B} \setminus \hat{B})} \lesssim \|\tilde{\psi} - \vec{u}(T)\|_{\mathcal{H}|\tilde{B}} + \|u(T)\|_{L^{2^*}(\tilde{B}^c)} \lesssim \varsigma, \tag{6.18}$$

and, combining it with (6.6) and (6.8),

$$\begin{aligned} \|\vec{u}^d(T) - \tilde{\psi}\|_{\tilde{\mathcal{H}}(\tilde{B})} &\leq \|\vec{u}(T) - \tilde{\psi}\|_{\tilde{\mathcal{H}}(\tilde{B} \cap \hat{B})} + \|\vec{u}^d(T)\|_{\tilde{\mathcal{H}}(\tilde{B} \setminus \hat{B})} + \|\tilde{\psi}\|_{\tilde{\mathcal{H}}(\tilde{B} \setminus \hat{B})} \\ &\lesssim \|\vec{u}(T) - \tilde{\psi}\|_{\mathcal{H}|\tilde{B}} + \|\vec{u}^d(T)\|_{\mathcal{H}|\hat{B}^c} + \varsigma \lesssim \varsigma. \end{aligned} \tag{6.19}$$

Expanding the energy for $\varphi := \vec{u}^d(T) - \tilde{\psi}$ yields

$$E(\vec{u}^d(T)) = E(W) + \frac{1}{2}[\langle L_{\tilde{\psi}}\varphi_1|\varphi_1 \rangle + \|\varphi_2\|_2^2] + o(\|\varphi_1\|_{2^*}^2), \tag{6.20}$$

where $L_{\tilde{\psi}} := -\Delta - f''(\tilde{\psi})$. Since $\|u^d(T)\|_{L^{2^*}(\tilde{B}^c \cup \hat{B}^c)} + \|\tilde{\psi}_1\|_{L^{2^*}(\tilde{B}^c \cup \hat{B}^c)} \lesssim \varsigma$ and (6.19), we have $\|\varphi_1\|_{L^{2^*}(\mathbb{R}^d)} \lesssim \varsigma$ and so

$$E(W) + O(\varsigma^2) = E(\vec{u}^d(T)) = E(W) + \frac{1}{2}\|\varphi\|_{\mathcal{H}}^2 + o(\varsigma^2). \tag{6.21}$$

Thus we obtain

$$d_W(\vec{u}^d(T)) \lesssim \|\vec{u}^d(T) - \vec{\psi}\|_{\mathcal{H}} \lesssim \zeta. \tag{6.22}$$

Choose $\zeta_* \ll \varepsilon_*$ of [14, Theorem 5.1], so that we can apply that one-pass theorem to u^d , both from $t = 0$ forward in time, and from $t = T$ backward in time. Specifically, there are $\varepsilon, \delta > 0$ such that

$$\begin{aligned} \zeta \sim \varepsilon \sim \delta, \quad \varepsilon \leq \varepsilon_*, \quad \sqrt{2}\varepsilon < \delta \leq \delta_*, \\ E(\vec{u}^d) \leq E(W) + \varepsilon^2, \quad \max(d_W(\vec{u}^d(0)), d_W(\vec{u}^d(T))) < \delta \end{aligned} \tag{6.23}$$

so that we can deduce from the theorem that $\max_{0 \leq t \leq T} d_W(\vec{u}^d(t)) < \delta$, and then, by the above lemma, $d_{\mathcal{R}}(\vec{u}) \lesssim \delta \sim \zeta$ for $0 < t < T$.

Next, if $d_{\mathcal{R}}(\vec{u}(t_0)) \geq C_*\zeta$ at some t_0 , with $C_* > 1$ large enough, then the above lemma implies that $d_W(\vec{u}^d(t_0)) > \delta$ and so, by the classification of the dynamics after ejection in [14], we conclude that $d_W(\vec{u}^d(t)) \gtrsim \delta_*$ after some $t_1 \in (t_0, T)$, and then u^d either blows up in finite time, or scatters to 0, so does u by (6.6). The blow up occurs away from the ground states in the sense of (5.2). Moreover, this dichotomy is determined by $\text{sign}(K(u^d(t_1)))$. Choosing ζ_* smaller if necessary, and using (6.6), we have

$$\pm\delta_* \sim K(u^d(t_1)) = K_{B_{t_1}}(u(t_1)) + O(\zeta), \tag{6.24}$$

which implies $\text{sign } K_{B_{t_1}}(u(t_1)) = \text{sign } K(u^d(t_1))$. □

In particular, we can characterize the manifold with large radiation, constructed in the previous section, by using the radiative distance.

Corollary 6.4 *Let $\vec{u} \in \text{S}_{\text{olution}}([0, T])$. If $\vec{u}(0) \in \mathcal{M}_3$ then $\sup_{0 < t < T} d_{\mathcal{R}}(\vec{u}(t)) \lesssim \delta_m$. Conversely, if $\sup_{0 < t < T} d_{\mathcal{R}}(\vec{u}(t)) < \min(\zeta_m, \zeta_*/C_*)$ then $\vec{u}(0) \in \mathcal{M}_3$.*

Proof Let $\vec{u}(0) \in \mathcal{M}_3$, then we can apply the detaching Lemma 4.4 with (5.1), so

$$d_{\mathcal{R}}(\vec{u}(0)) \leq \|\vec{u}^d(0) - \psi\|_{\mathcal{H}|_B} + \|\vec{u}(0)\|_{\mathcal{R}_B^\infty} + \|\psi\|_{\mathcal{H}|_B^c} \lesssim \delta_m + \zeta_m \lesssim \delta_m, \tag{6.25}$$

where $\psi := \mathcal{F}_{\vec{u}^d(0)} \vec{W}$. Since $\vec{u}^d(0) \in \mathcal{M}_0$, using (6.7) we obtain $d_{\mathcal{R}}(\vec{u}(t)) \lesssim d_W(\vec{u}^d(t)) + \delta_m \lesssim \delta_m$.

If $\sup_{0 < t < T} d_{\mathcal{R}}(\vec{u}(t)) < \min(\zeta_*, \zeta_m/C_*)$, then the above theorem implies that

$$d_{\mathcal{R}}(\vec{u}(0)) + \sup_{0 < t < T} d_W(\vec{u}^d(t)) < \zeta_m \ll \delta_m. \tag{6.26}$$

Hence $\vec{u}^d(0) \in \mathcal{M}_0$ and the definition of $d_{\mathcal{R}}$ implies that $\vec{u}(0) \in \mathcal{M}_3$. □

We can choose those distance parameters such as $\zeta_m = \delta_m/C$ and $\zeta_* = \zeta_m/C$ with some large absolute constant $C > 1$, provided that δ_m is chosen much smaller than the energy parameter $\varepsilon_* > 0$ of the one-pass theorem in [14].

The above one-pass theorem does not preclude oscillation between $\zeta < d_{\mathcal{R}} < C_*\zeta$. In the case of d_W in [14], it was possible to exclude such oscillations completely thanks to the convexity in time of d_W^2 near $\text{S}_{\text{tatic}}(W)$, which is not inherited by $d_{\mathcal{R}}$. However, we can make an exact version of the above one-pass theorem by the flow.

Theorem 6.5 (One-pass theorem with large radiation) *There exist constants $C_* > 1 > \zeta_* > 0$, and an open set $X(\zeta) \subset \mathcal{H}$ for each $\zeta \in (0, \zeta_*]$ satisfying:*

- (1) $X(\zeta)$ is increasing, i.e. $\zeta_1 < \zeta_2 \implies X(\zeta_1) \subset X(\zeta_2)$.
- (2) Its boundary is in $\zeta \leq d_{\mathcal{R}} \leq C_*\zeta$, namely

$$d_{\mathcal{R}}(\varphi) < \zeta \implies \varphi \in X(\zeta) \implies d_{\mathcal{R}}(\varphi) < C_*\zeta. \quad (6.27)$$

- (3) No solution can return to it, namely for any $\vec{u} \in S_{\text{olution}}([0, T])$

$$\vec{u}(0) \in X(\zeta), \exists t_0 \in (0, T), \vec{u}(t_0) \notin X(\zeta) \implies \forall t \in [t_0, T), \vec{u}(t) \notin X(\zeta). \quad (6.28)$$

Moreover, such u either blows up in finite time or scatters to 0 in $t > t_0$.

Proof Let $\zeta_* > 0$ and $C_* > 1$ be as in the previous theorem, though we will modify them at the end of proof. For $0 < \zeta \leq \zeta_*/C_*^2$, let $X(\zeta)$ be the totality of the initial data $\vec{u}(0)$ of any solution $\vec{u} \in S_{\text{olution}}((T_-, T_+))$ with $T_- < 0 < T_+$ satisfying

$$\inf_{T_- < t \leq 0} d_{\mathcal{R}}(\vec{u}(t)) < \zeta, \quad \inf_{0 \leq t < T_+} d_{\mathcal{R}}(\vec{u}(t)) < C_*\zeta. \quad (6.29)$$

Since $d_{\mathcal{R}}$ is continuous in \mathcal{H} , the local wellposedness implies that $X(\zeta)$ is open. Since the left quantity is non-increasing while the right quantity is non-decreasing along the flow, the no-return property (3) is obvious. At the exit time t_0 we have

$$d_{\mathcal{R}}(\vec{u}(t_0)) = \inf_{t_0 \leq t < T_+} d_{\mathcal{R}}(\vec{u}(t)) = C_*\zeta, \quad (6.30)$$

and the previous theorem implies that such a solution u either scatters to 0 or blows up in $t > t_0$. It also implies that $d_{\mathcal{R}}(\vec{u}(0)) \leq C_*^2\zeta$. By definition we have $d_{\mathcal{R}} \geq \zeta$ on $X(\zeta)^{\text{G}}$. Hence replacing ζ_* with ζ_*/C_*^2 and then C_* with C_*^2 , we obtain the desired properties of $X(\zeta)$. \square

Appendix A: Concentration blowup in the interior of blowup region

Here we observe that type-II blow-up is not always on the dynamical boundary between the scattering to 0 and blow-up. More precisely, we have

Proposition 6.1 *Let $0 < T < \infty$, $\vec{u}^0 \in S_{\text{olution}}([0, T]) \cap L^\infty([0, T]; \mathcal{H})$. Then for any $\delta > 0$, there is $\vec{u}^1 \in S_{\text{olution}}([0, T])$ with the following property: $\vec{u}^1(t) - \vec{u}^0(t)$ has a strong limit as $t \nearrow T$, and for any $t \in [0, T)$ and any $\psi \in \mathcal{H}$ with $\|\psi\|_{\mathcal{H}} < \delta$, the solution starting from $\vec{u}^1(t) + \psi$ blows up in positive finite time.*

In other words, for any blow-up with bounded energy norm, there is another solution with the same blow-up profile, whose orbit is in the interior of the blow-up set of initial data, with arbitrarily large distance from the exterior.

Proof Fix $R \geq 1 + T$ such that $\|\bar{u}^0(0)\|_{\mathcal{H}(|x|>R)} \ll 1$. Let u^2 be the solution for the initial data $\bar{u}^2(0) = \Gamma(x/R)\bar{u}^0(0)$, where Γ is a smooth radial function on \mathbb{R}^d satisfying $\Gamma(x) = 1$ for $|x| \leq 3$ and $\Gamma(x) = 0$ for $|x| \geq 4$. Then the finite speed of propagation implies that for $0 < t < T$ and as long as u^2 exists,

- (1) $\bar{u}^2(t) = \bar{u}^0(t)$ on $|x| < 3R - t$,
- (2) $\|\bar{u}^2(t)\|_{\mathcal{H}(|x|>R+t)} \ll 1$,
- (3) $\text{supp } \bar{u}^2(t) \subset \{|x| < 4R + t\}$.

Since the regions for (1) and for (2) cover $[0, T) \times \mathbb{R}^d$, we deduce that u^2 extends beyond $t < T$. Moreover, both u^0 and u^2 extend to $|x| > R + t$ for all $t > 0$ by the smallness in the exterior cone. Hence $\bar{u}^2(t) - \bar{u}^0(t)$ has a strong limit in \mathcal{H} as $t \nearrow T$.

Now fix $\delta > 0$. Since u^2 is bounded in \mathcal{H} for $0 < t < T$,

$$M := \sup\{E_{|x|<5R}(\bar{u}^2(t) + \psi) \mid t \in [0, T), \|\psi\|_{\mathcal{H}} < \delta\} \tag{7.1}$$

is finite. Then we can find a strong radial solution u^3 such that

- (1) $\text{supp } \bar{u}^3(t) \subset \{|x| > 6R - t\}$.
- (2) $\sup\{E_{|x|>5R}(\bar{u}^3(t) + \psi) \mid t \in [0, T), \|\psi\|_{\mathcal{H}} < \delta\} < -M - 1$.

Indeed, it is easy to satisfy (1) and (2) at $t = 0$ by using a very flat radial smooth function, since for any $\varphi, \psi \in \mathcal{H}$ and any $0 < \varepsilon \ll 1$,

$$E_{|x|>5R}(\varphi + \psi) \leq (1 + \varepsilon)E_{|x|>5R}(\varphi) + C_\varepsilon(\|\psi\|_{\mathcal{H}}^2 + \|\psi_1\|_{2^*}^2). \tag{7.2}$$

(1) is preserved for $t > 0$ by the finite speed of propagation. For such initial data, the solution may blow up in finite time, but we can delay the blow-up time as much as we like by the rescaling \bar{S}^σ with $\sigma \rightarrow -\infty$, which makes both (1) and (2) easier. This yields $u^3 \in \text{S}_{\text{olution}}([0, 2T])$ with the above properties.

Now let u^1 be the strong solution for the initial data

$$\bar{u}^1(0) = \bar{u}^2(0) + \bar{u}^3(0). \tag{7.3}$$

Then the finite propagation property together with the disjoint supports of u^2 and u^3 implies that $\bar{u}^1 = \bar{u}^2$ for $|x| < 6R - t$, $\bar{u}^1 = \bar{u}^3$ for $|x| > 4R + t$, so $\bar{u}^1 = \bar{u}^2 + \bar{u}^3$ for $0 < t < T$, and $\bar{u}^1(t) - \bar{u}^0(t)$ has a strong limit in \mathcal{H} as $t \nearrow T$. Moreover, for any $t \in [0, T)$ and any $\psi \in \mathcal{H}$ satisfying $\|\psi\| < \delta$ we have

$$E(\bar{u}^1(t) + \psi) = E_{|x|<5R}(\bar{u}^2(t) + \psi) + E_{|x|>5R}(\bar{u}^3(t) + \psi) < -1, \tag{7.4}$$

hence the solution starting from $\bar{u}^1(t) + \psi$ has to blow up in finite time because of the negative energy, see [11, 18]. □

Appendix B: Table of notation

$\triangleleft X^{\triangleright}$	$= X^1 - X^0$	(2.12)
\bar{u}	$= (u, \dot{u})$ vector in the phase space	(1.2)
φ^\dagger	$= (\varphi_1, -\varphi_2)$ time inversion	(1.17)
$\langle \cdot \cdot \rangle$	L^2 inner product	(1.36)
\mathcal{B}_d	Borel sets in \mathbb{R}^d	
(CW)	The critical wave equation	(1.1)
$\mathcal{H}, \mathcal{H}_\perp$	Energy space, its subspace	(1.2),(1.35)
$S_{\text{olution}}(I)$	Solutions of (CW) on I	(1.27)
$E(\bar{u}), P(\bar{u})$	Total energy and momentum	(1.4),(1.5)
$E_B(\varphi), K_B(\varphi)$	Restricted energy functionals	(3.15)
$U(t)$	Free propagator	(1.12)
$\mathcal{T}^c, \mathcal{S}^\sigma, \mathcal{S}_a^\sigma$	Invariant translation and scaling	(1.28)
$\text{St}, \text{St}_*^*, q_s, q_m$	Strichartz norms and exponents	(1.22)
$W, S_{\text{tatic}}(W)$	Ground states	(1.8),(1.9)
$S_{\text{oliton}}(W)$	Ground solitons	(1.10)
dist_W, d_W	Distances to the ground states	(1.16),(1.59)
L_+, \mathcal{L}	Linearized operators around W	(1.30),(1.32)
$N(v), \underline{N}(v)$	Higher order terms	(1.31),(1.54)
ρ, k, P_\perp	Ground state of L_+	(1.34)
g_\pm, Λ_\pm	(un)stable modes of $J\mathcal{L}$	(1.39),(1.40)
$v, \lambda, \gamma, \lambda_\pm$	Components of u around W	(1.38),(1.39)
(α, μ)	Parameters to define the orthogonality	(1.42)
$(\tilde{\sigma}, \tilde{c}), \mathcal{T}_\varphi, \tilde{\lambda}$	Local coordinates by the orthogonality	(1.52), Lemma 1.4
$Z = (Z_1, Z_2)$	Modulation operator in the equation	(1.54)
τ	Rescaled time variable	(1.53)
$\ \varphi\ _E, \nu(\tau)$	Linearized energy norms	(1.43),(2.42)
$\mathcal{B}_\delta, \mathcal{B}_\delta^+, \mathcal{B}'_\delta$	Small balls for different components	(1.45), (3.3)
$\mathcal{N}_\delta, \mathcal{N}_{\delta_1, \delta_2}$	Neighborhoods of $S_{\text{tatic}}(W)$	(1.45),(2.2)
$\Phi_{\sigma, c}, \Psi_{\sigma, c}$	Local coordinates around $S_{\text{tatic}}(W)$	(1.44)
$\mathcal{M}_0 \sim \mathcal{M}_5$	Local manifold and its extensions	(2.8),(3.1),(3.33),(5.1)
m_+, M_+	Functions defining the local manifold	Theorem 2.1,(2.11)
a_W, b_W	Positive constants	(1.48),(1.49)
ε_S	Small Strichartz norm for scattering	(1.24)
δ_Φ, δ_m	Small distances from $S_{\text{tatic}}(W)$	Lemma 1.4, Theorem 2.1
ι_I	Smallness in the ignition lemma	Lemma 2.2
η_I	τ -length for uniform Strichartz bound	Lemma 2.3
ζ_m, ζ_*	Smallness in radiative distance	(5.1), Theorems 6.3,6.5
ε_*, δ_*	Smallness in the one-pass theorem	[14, Theorem 5.1]
$\kappa(\delta)$	Variational bound on K	[14, Lemma 4.1]
B_{+a}, B_{-a}	Fattened and thinned sets by radius a	(3.3),(3.5)
$\mathcal{H} \lfloor B, \tilde{\mathcal{H}}(B)$	Restrictions of \mathcal{H} to B	(3.7),(3.10)
X_B	Extension operator from B to \mathbb{R}^d	(3.14)
$\mathcal{D}(u), t_\pm(\varphi, x)$	Maximal space-time domain of solution	(3.26)
\mathcal{R}_B^T	Seminorm measuring radiation	(4.1)
$u^{\mathbf{d}}, u^{\mathbf{x}}$	Detached interior and exterior solutions	Lemma 4.4
$d_{\mathcal{R}}(\varphi)$	Radiative distance to the ground states	(6.1)

References

1. Beceanu, M.: A centre-stable manifold for the energy-critical wave equation in \mathbb{R}^3 in the symmetric setting. [arXiv:1212.2285](#)
2. Bizon, P., Chmaj, T., Tabor, Z.: On blowup for semilinear wave equations with a focusing nonlinearity. *Nonlinearity* **17**(6), 2187–2201 (2004)
3. Donninger, R., Huang, M., Krieger, J., Schlag, W.: Exotic blowup solutions for the u^5 focusing wave equation in \mathbb{R}^3 . [arXiv:1212.4718](#)
4. Donninger, R., Krieger, J.: Nonscattering solutions and blow up at infinity for the critical wave equation. [arXiv:1201.3258v1](#)
5. Duyckaerts, T., Kenig, C., Merle, F.: Universality of blow-up profile for small radial type II blow-up solutions of energy-critical wave equation. *J. Eur. Math. Soc.* **13**(3), 533–599 (2011)
6. Duyckaerts, T., Kenig, C., Merle, F.: Universality of the blow-up profile for small type II blow-up solutions of energy-critical wave equation: the non-radial case. [arXiv:1003.0625](#), to appear in JEMS
7. Duyckaerts, T., Kenig, C., Merle, F.: Profiles of bounded radial solutions of the focusing, energy-critical wave equation. [arXiv:1201.4986](#), to appear in GAFA
8. Duyckaerts, T., Kenig, C., Merle, F.: Classification of radial solutions of the focusing, energy-critical wave equation. [arXiv:1204.0031](#)
9. Duyckaerts, T., Merle, F.: Dynamic of threshold solutions for energy-critical wave equation. *Int. Math. Res. Pap. IMRP* **2007**(4), 67, Art. ID rpn002 (2008)
10. Hillairet, M., Raphaël, P.: Smooth type II blow up solutions to the four dimensional energy critical wave equation. [arXiv:1010.1768](#)
11. Kenig, C., Merle, F.: Global well-posedness, scattering and blow-up for the energy-critical focusing non-linear wave equation. *Acta Math.* **201**(2), 147–212 (2008)
12. Krieger, J., Nahas, J.: Instability of type II blow up for the quintic nonlinear wave equation on \mathbb{R}^{3+1} . [arXiv:1212.4628](#)
13. Krieger, J., Nakanishi, K., Schlag, W.: Global dynamics away from the ground state for the energy-critical nonlinear wave equation. *Am. J. Math.* **135**(4), 935–965 (2013)
14. Krieger, J., Nakanishi, K., Schlag, W.: Global dynamics of the nonradial energy-critical wave equation above the ground state energy. *DCDS* **33**, 6 (2013)
15. Krieger, J., Nakanishi, K., Schlag, W.: Threshold phenomenon for the quintic wave equation in three dimensions. [arXiv:1209.0347](#)
16. Krieger, J., Schlag, W.: On the focusing critical semi-linear wave equation. *Am. J. Math.* **129**(3), 843–913 (2007)
17. Krieger, J., Schlag, W., Tataru, D.: Slow blow-up solutions for the $H^1(\mathbb{R}^3)$ critical focusing semilinear wave equation. *Duke Math. J.* **147**(1), 1–53 (2009)
18. Levine, H.A.: Instability and nonexistence of global solutions to nonlinear wave equations of the form $Pu_{tt} = -Au + \mathcal{F}(\nabla)$. *Trans. Am. Math. Soc.* **192**, 1–21 (1974)
19. Nakanishi, K., Schlag, W.: Global dynamics above the ground state energy for the focusing nonlinear Klein-Gordon equation. *J. Diff. Eq.* **250**, 2233–2299 (2011)
20. Nakanishi, K., Schlag, W.: Global dynamics above the ground state energy for the cubic NLS equation in 3D. *Calc. Var. PDE* **44**(1–2), 1–45 (2012)
21. Nakanishi, K., Schlag, W.: Global dynamics above the ground state for the nonlinear Klein-Gordon equation without a radial assumption. *Arch. Ration. Mech. Anal.* **203**(3), 809–851 (2012)
22. Nakanishi, K., Schlag, W.: Invariant manifolds around soliton manifolds for the nonlinear Klein-Gordon equation. *SIAM J. Math. Anal.* **44**(2), 1175–1210 (2012)
23. Payne, L.E., Sattinger, D.H.: Saddle points and instability of nonlinear hyperbolic equations. *Israel J. Math.* **22**(3–4), 273–303 (1975)
24. Szpak, N.: Relaxation to intermediate attractors in nonlinear wave equations. *Theor. Math. Phys.* **127**, 817 (2001)