

# *Arabidopsis* ABCB21 is a Facultative Auxin Importer/Exporter Regulated by Cytoplasmic Auxin Concentration

Yoshihisa Kamimoto<sup>1</sup>, Kazuyoshi Terasaka<sup>2</sup>, Masafumi Hamamoto<sup>1</sup>, Kojiro Takanashi<sup>1</sup>, Shoju Fukuda<sup>1</sup>, Nobukazu Shitan<sup>1</sup>, Akifumi Sugiyama<sup>1</sup>, Hideyuki Suzuki<sup>3</sup>, Daisuke Shibata<sup>3</sup>, Bangjun Wang<sup>4,6</sup>, Stephan Pollmann<sup>5,7</sup>, Markus Geisler<sup>4,6</sup> and Kazufumi Yazaki<sup>1,\*</sup>

<sup>1</sup>Laboratory of Plant Gene Expression, Research Institute for Sustainable Humanosphere, Kyoto University, Gokasho Uji, 611-0011 Japan

<sup>2</sup>Graduate School of Pharmaceutical Sciences, Nagoya City University, Japan

<sup>3</sup>Kazusa DNA Research Institute, 2-6-7 Kazusakamatari, Kisarazu, Chiba, 292-0818 Japan

<sup>4</sup>Institute of Plant Biology, Basel–Zurich Plant Science Center, University of Zurich, CH-8008 Zurich, Switzerland

<sup>5</sup>Ruhr-Universität Bochum, Lehrstuhl für Pflanzenphysiologie, Germany

<sup>6</sup>Present address: University of Fribourg, Department of Biology, Plant Biology, CH-1700 Fribourg, Switzerland.

<sup>7</sup>Present address: Centro de Biotecnología y Genómica de Plantas, Campus de Montegancedo, 28223 Pozuelo de Alarcón, Madrid, Spain.

\*Corresponding author: E-mail, yazaki@rsh.kyoto-u.ac.jp; Fax, +81-774-38-3623.

(Received February 2, 2012; Accepted October 31, 2012)

The phytohormone auxin is critical for plant growth and many developmental processes. Members of the P-glycoprotein (PGP/ABCB) subfamily of ATP-binding cassette (ABC) transporters have been shown to function in the polar movement of auxin by transporting auxin over the plasma membrane in both monocots and dicots. Here, we characterize a new *Arabidopsis* member of the ABCB subfamily, ABCB21/PGP21, a close homolog of ABCB4, for which conflicting transport directionalities have been reported. ABCB21 is strongly expressed in the abaxial side of cotyledons and in junctions of lateral organs in the aerial part, whereas in roots it is specifically expressed in pericycle cells. Membrane fractionation by sucrose density gradient centrifugation followed by Western blot showed that ABCB21 is a plasma membrane-localized ABC transporter. A transport assay with *Arabidopsis* protoplasts suggested that ABCB21 was involved in IAA transport in an outward direction, while naphthalene acetic acid (NAA) was a less preferable substrate for ABCB21. Further functional analysis of ABCB21 using yeast import and export assays showed that ABCB21 mediates the 1-*N*-naphthylphthalamic acid (NPA)-sensitive translocation of auxin in an inward direction when the cytoplasmic IAA concentration is low, whereas this transporter mediates outward transport under high internal IAA. An increase in the cytoplasmic IAA concentration by pre-loading of IAA into yeast cells abolished the IAA uptake activity by ABCB21 as well as ABCB4. These findings suggest that ABCB21 functions as a facultative importer/exporter controlling auxin concentrations in plant cells.

**Keywords:** ABC protein • ABCB21 • *Arabidopsis thaliana* • Auxin transport • GUS expression • Pericycle cell • Transport direction.

**Abbreviations:** ABC, ATP-binding cassette; BA, benzoic acid; DTT, dithiothreitol; 5-FI, 5-fluoroindole; GUS,  $\beta$ -glucuronidase; LR, lateral root; NAA, naphthalene acetic acid; NBD, nucleotide-binding domain; NPA, 1-*N*-naphthylphthalamic acid; PGP, P-glycoprotein; PMSF, phenylmethylsulfonyl fluoride; RNAi, RNA interference; RT-PCR, reverse transcription-PCR.

## Introduction

Higher plants develop various organs after germination; i.e. in aerial parts, the shoot meristem produces leaves and stems, while in underground parts, a branching root system is developed through the formation of lateral roots (LRs) from the primary root. LR formation is an important developmental process that contributes to the establishment of root conformation to enable the plant to absorb water and nutrients efficiently, and physically to sustain the aerial parts (Park et al. 2005). In *Arabidopsis thaliana*, for example, LRs are generated from several pericycle cells adjacent to the xylem parenchyma, which starts to divide in a highly regulated manner, to form new root tissue toward the lateral axis. The first trigger of cell division at the pericycle leading to LR formation is the event of auxin entry into the cell (Casimiro et al. 2001). Auxin is also deeply involved in vertical axis formation by the polar transport via cell-to-cell movement of IAA in xylem parenchyma cells in plants.

Membrane transport of auxin is regulated by several types of transporters, e.g. PIN proteins (major facilitator superfamily; MFS) and AUX1/LAX H<sup>+</sup> symporters (Schnabel and Frugoli 2004, Xu et al. 2005, Carraro et al. 2006, Wiśniewska et al. 2006, Bainbridge et al. 2008). In addition, ATP-binding cassette

(ABC) transporters have been discovered as the third major family responsible for the membrane transport of auxin. Out of approximately 120 ABC transporters in a plant, mostly the P-glycoprotein (PGP)/ABCB (hereafter referred to as ABCB) subclass (Jasinski et al. 2003, Verrier et al. 2008) is reported thus far as being relevant for auxin transport (Noh et al. 2001, Luschnig 2002, Noh et al. 2003, Geisler et al. 2005, Terasaka et al. 2005). First, two ABCBs, ABCB1 and ABCB19, in *A. thaliana* were shown to bind specifically the auxin efflux inhibitor 1-N-naphthylphthalamic acid (NPA) (Murphy et al. 2002). T-DNA insertion mutants of *abcb19* showed pleiotropic auxin-related phenotypes, and characterization of the double mutant *abcb1/abcb19* and antisense *abcb1* plants strongly suggested that these ABC transporters are necessary for the polar auxin transport.


Previously, we identified another plasma membrane ABCB-type transporter, (PGP4)/ABCB4, in *A. thaliana*, which was specifically expressed in the root epidermis, and demonstrated its involvement in basipetal shootward auxin transport, i.e. from the root tip to the root–shoot transition zone (Terasaka et al. 2005). Most plant ABCB members form pairs sharing high sequence similarity with each other (Geisler and Murphy 2006). The paralog of ABCB4 in the Arabidopsis genome is ABCB21 (68% nucleotide identity and 79% amino acid identity), but no biochemical and molecular biological characterization was made on this ABCB member. Besides, for ABCB4, conflicting transport directionalities, either inward or outward transport activity, have been reported, depending on the experimental systems (Santelia et al. 2005, Terasaka et al. 2005, Cho et al. 2007, Lewis et al. 2007, Kubeš et al. 2012). In this report, we describe the biochemical characterization of ABCB21 comparing the transport property of ABCB4 as well as ABCB1, and suggest that ABCB21 functions as facultative transporter that mediates import or export of IAA depending on the cytoplasmic IAA concentration.

## Results

### Expression analysis of ABCB21

ABCB21 locates at chromosome 3 and is divided into 11 exons, and the deduced polypeptide consists of 1,293 amino acids (Fig. 1a). This ABC protein is grouped in cluster II of the ABCB subfamily, which is separate from cluster I where ABCB1 and ABCB19 are classified (Fig. 1b). Due to the high sequence similarity between ABCB4 and ABCB21, a possible transcriptional correlation was evaluated employing a microarray analysis, in which the *abcb4* mutant and wild-type Arabidopsis were compared (Supplemental Fig. S1). In *abcb4*, 158 genes showed a >2-fold elevated expression (Supplemental Table S1a), whereas 21 genes revealed less than half that level of expression (Supplemental Table S1b), but no ABC protein genes, including ABCB21, were observed in these lists.

To evaluate the organ-specific expression of ABCB21, a real-time PCR analysis was conducted, and this indicated that




**Fig. 1** Molecular characterization of ABCB21. (a) Exon–intron structure of the ABCB21 gene. Exons are shown as shaded boxes and sequences coding for two nucleotide-binding domains (NBD) are shown as white boxes. The region used to prepare the RNAi construct is indicated with arrowheads. (b) Phylogenetic tree of the Arabidopsis ABCB subfamily, which is classified into three clusters. ABCB21 belonging to cluster II is highlighted. Numbers indicate bootstrap values.

the ABCB21 transcript was detectable in all Arabidopsis tissues examined (Fig. 2a). This expression pattern is different from that of ABCB4 that is almost solely expressed in the root, in particular in the epidermis and the root cap (Terasaka et al. 2005), while strong expression of ABCB21 was also seen in leaf tissues.

We analyzed the response of ABCB21 expression to various phytohormones and abiotic stresses using 14-day-old seedlings of Arabidopsis. While ABCB21 expression was decreased upon treatments with benzyladenine, ABA and gibberellic acid, as well as cold or dark treatments, no increase in the gene expression was observed even by auxins at a high concentration in this primary screening (Supplemental Fig. S2). However, a time course experiment with a low concentration (300 nM) of IAA showed that the seedlings responded to exogenously added IAA in a different pattern from that of ABCB4, i.e. the ABCB4 expression increased rapidly and then decreased gradually, whereas ABCB21 expression decreased transiently and then returned to a steady-state level gradually (Fig. 2b), suggesting a different physiological role for ABCB21 from that for ABCB4.

A detailed cell type-specific expression analysis of ABCB21 was performed using a *ProABCB21:β-glucuronidase (GUS)* transformant at the T<sub>3</sub> generation, in which a 0.75 kb sequence upstream of the ABCB21 initiation codon was fused to a GUS reporter gene. The *ProABCB21:GUS* transformants were



**Fig. 2** Expression profile of ABCB21 in Arabidopsis. (a) Organ-specific expression of ABCB21 monitored by real-time RT-PCR. The experiment was repeated three times, and error bars indicate the standard deviation. (b) Time course of auxin response of ABCB21 and ABCB4 gene expression. Seedlings were treated with 300 nM IAA, and sampled at 0, 0.5, 1, 3 and 7 d after treatment. Total RNA (10 µg) prepared from whole seedlings was probed with <sup>32</sup>P-labeled ABCB21 (top panel) and ABCB4 fragments (middle panel). Loading controls are shown by β-actin (bottom panel). The experiment was repeated twice to yield similar results.

histochemically stained with X-Gluc at different developmental stages. The promoter length used in this experiment was 0.75 kb, because another gene (At3g62160) was found upstream of the promoter region. At the aerial part, ABCB21 was expressed in the cotyledon, which was almost specifically located at the abaxial side (Fig. 3a–c). Strong expression was also observed at junctions between the cauline leaf and stem, and the petal and inflorescence stem (Fig. 3d–f). Weak expression was seen in vascular tissues in petals (Fig. 3g). In young seedlings shortly after hatching, GUS staining was prominent in the boundary regions of hydrathodes of cotyledons (Supplementary Fig. S3). The most remarkable cell type-specific expression of ABCB21 was observed in the root tissue, where GUS staining was restricted to the pericycle; in particular, the strongest staining was seen at the part adjacent to the xylem parenchyma, but not at the root cap (Fig. 3h–m). In summary, it appears that despite their high sequence similarity, the expression pattern and responsiveness to auxin of ABCB21 and ABCB4 are highly complementary.

### Membrane localization of ABCB21

Membrane localization of ABCB21 was analyzed by sucrose density gradient fractionation of microsomal membranes followed by Western blot, in which a specific peptide antibody directed against ABCB21 was used. The ABCB21 polypeptide peaks at the 32/38% fraction that coincides with that of the

plasma membrane H<sup>+</sup>-ATPase AHA, and the fractionation pattern is apparently different from those of the H<sup>+</sup>-pyrophosphatase (AVP1) and of the endoplasmic reticulum luminal binding protein (BiP), which were used as markers of vacuolar and endoplasmic reticulum membrane proteins, respectively (Fig. 4). These results suggest that ABCB21 is located on the plasma membrane and not on vacuolar membranes or the endoplasmic reticulum.


### ABCB21 is a facultative auxin importer/exporter

To analyze the transport function of ABCB21 in planta, we obtained an insertion (WiscDsLox1C2) mutant of the ABCB21 gene from the Wisconsin mutant collection. However, reverse transcription-PCR (RT-PCR) analysis of mutant seedlings indicated that the transcript corresponding to the 3'-truncated ABCB21 open reading frame was clearly detected and the coding sequence for the first ATPase domain remained (Supplementary Fig. S4). Adding to the absence of the *P*-value of this gene in the microarray analysis of the loss-of-function mutant (not shown) and its low germination rate, we did not use the strain for further studies.

As the second best option, we therefore generated ABCB21 RNA interference (RNAi) lines, in which an intron-spliced hairpin RNA (RNAi) construct targeted to the ABCB21 sequence was introduced (Fig. 1a). Among 18 kanamycin-resistant transformants, two lines, designated *abcb21-ir1* and *abcb21-ir2*, showed strong silencing at the protein level and were used for further studies (Fig. 5a). Importantly, in these RNAi lines, ABCB4 expression was not significantly influenced.

Having the proposed import directionality for ABCB4 (Santelia *et al.* 2005, Terasaka *et al.* 2005), we performed auxin import and export assays in ABCB21 RNAi plant (line *ir-1*) mesophyll protoplasts in the presence of 100 nM external IAA (Fig. 5b). Surprisingly, in import assays, the ABCB21 RNAi line showed significantly increased IAA uptake compared with the vector control, indicating, together with the plasma membrane localization, that the transport directionality of ABCB21 may be pointing outward in Arabidopsis protoplasts (Fig. 5b). Using naphthalene acetic acid (NAA) double labeling in the same assay, we found no significant difference in NAA loading between the RNAi line (*ir-1*) and the vector control (Fig. 5c), which is in line with the chemical feature of NAA, unlike IAA, to be able to by-pass carrier-mediated uptake by diffusion (Marchant *et al.* 1999). However, reduced NAA export capacities have also been recently reported for ABCB4 using tobacco BY2 cells (Kubeš *et al.* 2012). In agreement with this proposed efflux directionality, the export of IAA by the RNAi line was reduced in export assays where outside concentrations of auxins were monitored (Fig. 5d), while the transport activity for NAA was not at a significant level (Fig. 5e).

The transport activity of ABCB21 for IAA was further investigated at the cellular level in yeast. As previously reported for ABCB1 (Geisler *et al.* 2005), the plasma membrane localization of ABCB21 expressed in yeast cells was confirmed by membrane




**Fig. 3** GUS staining of the root tissues of *ProABCB21:GUS* transformants. Transgenic plants were grown on 1/2 MS medium containing 1% sucrose, and the plant was stained with X-Gluc for 24 h. (a) GUS activity in *ProABCB21:GUS* cotyledons (whole mount). (b) Magnification of (a). (c) Cross-section of a cotyledon. (d) GUS activity between a cauline leaf and stem junction. (e) GUS activity between a silique and stem junction. (f) GUS activity between a flower and stem junction. (g) GUS activity in the vein of petals. (h) GUS activity in *ProABCB21:GUS* roots (whole mount). (i) Cross-section of a root. (j) Longitudinal section of a root. (k) Enlargement of (i). (l) Representative GUS activity in the root tip. (m) Minority of GUS activity in the root tip. The experiment was repeated three times with similar results. Bars = 100  $\mu$ m, except for (k) where it is 40  $\mu$ m.

fractionation using sucrose density gradient centrifugation followed by Western blot analysis (Fig. 6a). The IAA-sensitive strain *gef1* (for Glycerol/ethanol Fe-requiring) lacks a putative chloride channel protein. In a growth test using 5-fluorindole (5-FI), a toxic analog of a potential IAA precursor, which provides a useful tool for evaluating auxin transport (Gaxiola et al. 1998), ABCB21, like ABCB4 (Santelia et al. 2005), showed a significant hypersensitivity compared with the vector control (Fig. 6b), supporting the assumption that ABCB21, like its closest homolog, might function as an auxin analog importer in yeast.

In order to clarify these seemingly conflicting transport data in more detail, we performed yeast IAA transport studies for ABCB21, in direct comparison with ABCB1 and ABCB4, as both import and export assays. As explained in detail in the Materials and Methods, in import assays external IAA is added at time point 0 resulting in high external IAA but low cytoplasmic auxin concentrations during this assay. In export assays, prior to export, IAA loading followed by removal of external IAA is performed, resulting in low external but high cytoplasmic auxin levels. In import assays (Fig. 7a), as reported, ABCB1 clearly reveals efflux activity for IAA, while ABCB21 as well as

ABCB4 show uptake activity in this assay. Use of the unspecific diffusion control, benzoic acid (BA), employed as double labeling in the same assay, demonstrated transport specificity of this assay. However, surprisingly, in export assays (Fig. 7c), all three




**Fig. 4** Membrane localization of ABCB21. Fractionation of total microsomes from wild-type *Arabidopsis* roots was done on a non-continuous sucrose gradient consisting of 0, 20, 32, 38 and 45% (w/v) sucrose. Membrane fractions were collected from the interfaces between different sucrose concentrations and analyzed via SDS-PAGE and protein gel blotting. Blots were probed with antisera against ABCB21, plasma membrane H<sup>+</sup>-ATPase (AHA), vacuolar H<sup>+</sup>-PPase (AVP1) and endoplasmic reticulum BiP.


ABCBs were capable of exporting IAA, resulting in lower IAA retentions, which again was not found with the diffusion control, BA.

To test the idea that the cytoplasmic IAA concentration indeed alters the transport directionality of ABCB4 and ABCB21, we repeated the import assays after pre-loading with IAA. Strikingly, pre-loading with 10 μM IAA at pH 4.5 completely blocked the ABCB4/ABCB21-mediated IAA import to the vector control level (Fig. 7b). IAA pre-loading also reduced ABCB1-catalyzed IAA export, which is, however, most probably caused by competition of unlabeled IAA with radiolabeled IAA. These findings strongly support our hypothesis that internal IAA regulates the transport directionalities of ABCB21 and ABCB4. Interestingly, BA export seems to be reduced in ABCB21- and ABCB4-expressing yeast, despite this not being significant, arguing that internal IAA might influence the specificity of transport substrates for ABCB21 and ABCB4 (Fig. 7b).

Finally, we also tested NPA binding of those ABCB members. Fig. 7d and e reveal that all three can bind to NPA to a similar extent, and that NPA suppressed the IAA transport function of



**Fig. 5** Transport assay of ABCB21 knock-down protoplast cells. (a) Protein gel blot analyses of the *abc21* RNAi line. The experiment was repeated three times and yielded similar results. (b) Import assay: IAA uptake into protoplasts of the *abc21* RNAi line compared with that of the vector control. Values are mean activities ± SEs, four individual measurements ( $n = 4$ ). (c) Import assay: NAA uptake into protoplasts of the *abc21* RNAi line compared with that of the vector control. Values are mean activities ± SEs, four individual measurements ( $n = 4$ ). (d) Export assay: IAA export from protoplasts of the *abc21* RNAi line compared with that of the vector control. Values are mean activities ± SEs, four individual measurements ( $n = 4$ ). (e) Export assay: NAA export from protoplasts of the *abc21* RNAi line compared with that of the vector control. Values are mean activities ± SEs, four individual measurements ( $n = 4$ ).



**Fig. 6** Membrane localization of ABCB21 in yeast and spot assay. (a) ABCB21 and ABCB4 co-migrate with the plasma membrane marker, H<sup>+</sup>-ATPase, in continuous sucrose gradients judged by Western detection. (b) ABCB21 was expressed in the 5-fluorindole (5-FI)-sensitive *gef1* mutant strain; ABCB4 was used for comparison. Cells were 5-fold diluted five times, and 5 μl of each dilution were spotted onto a SD-URA media plate supplemented with 50 μM 5-FI. Growth at 30°C was assessed after 3–4 d. Assays were performed with three independent transformants.

ABCB21 as well as of ABCB4, as was reported before for ABCB1 (Geisler et al. 2005).

In summary, these data imply that ABCB21, in analogy to its closest homolog, ABCB4, is functioning as an NPA-sensitive, plasma membrane-localized auxin transporter. However, in contrast to the strict auxin exporter, ABCB1, the transport directionality of ABCB21 and ABCB4 as importers/exporters is facultative and probably determined by cytoplasmic auxin levels.

### Phenotype observation of *abcb21* knockdown mutants

We have carefully looked for morphological differences in *abcb21* RNAi lines compared with wild-type plants, especially in the root phenotype. These RNAi lines tend to show retardation of the germination process, but they grow normally and revealed basically no significant morphological differences (Supplementary Fig. S5). In 10-day-old seedlings of the T<sub>3</sub> generation, for example, the main root length of the wild


type and RNAi line was  $22.3 \pm 4.8$  and  $21.7 \pm 5.5$  mm ( $n = 15$ ), respectively, whereas the number of LR was  $2.2 \pm 1.4$  and  $2.5 \pm 1.8$  ( $n = 15$ ) for the wild type and the RNAi line, respectively. The lack of a visible phenotype for *abcb21* RNAi lines may be a result of the putative functional redundancy of other uncharacterized ABCB transporters expressed in these tissues, or phenotype differences may only be observed under special stress conditions.

### Discussion

Thus far, membrane transporters of auxin have been studied intensively from the viewpoint of the polar movement of the IAA molecule along the vertical axis of the plant body, to which PIN proteins provide a dominant contribution. Similarly, both ABCB1 and ABCB19, as well as auxin importers of the AUX1/LAX family are also relevant for the downward movement of auxin in parenchyma cells. Another ABC protein of the B family, ABCB4, is expressed specifically at the root epidermis, which mediates the upward movement of the IAA molecule. In this study, we have demonstrated that a new member of the ABCB subfamily, ABCB21, is expressed in the abaxial side of cotyledons, in junctions of lateral organs in the aerial part and in pericycle cells adjacent to the protoxylem poles of Arabidopsis root, and further this ABC transporter recognized IAA as a transport substrate. Also, this transporter protein binds NPA, resulting in inhibition of IAA transport.

The initiation of the LR formation process is triggered by the entrance of the IAA molecule from the xylem parenchyma into pericycle cells (Laskowski et al. 1995, Himanen et al. 2002). In the microarray analysis of the *abcb21* RNAi lines, the expression of *LAX3*, *PIN5* and *SHY2* involved in LR development was significantly down-regulated (Bouchard et al. 2006, Blakeslee et al. 2007) (Supplementary Table S2). We then observed how ABCB21 expression is altered during the LR initiation process with ABCB21 promoter–GUS transformants (Supplementary Fig. S6). At the dividing pericycle cell, ABCB21 expression disappeared when LR initiation occurs, which is phase II to III according to the literature (Vanneste et al. 2005). This suggests that ABCB21 may be relevant for LR initiation, but once the initiation occurs the role of this transporter can be terminated. We thus expected that knock-down lines would show phenotype alterations in root development, but no clear difference was observed (Supplementary Fig. S5). This does, however, occur relatively frequently in the knock-down lines, or even in knock-out mutants of ABC transporter genes, because there is a high redundancy of this gene family in plants. If an uncharacterized transporter of functional redundancy is identified, the double knock-out mutant might provide a more informative phenotype as reported for ABCB1 and ABCB19 (Noh et al. 2001).

The *abcb21* loss-of-function line (Wisconsin No. CS848985) showed a very low germination rate, and germinated plants did not show a clear auxin-related phenotype either, including LR



**Fig. 7** Transport assay of ABCB21 in yeast. (a) Import assay (high external IAA): IAA and the diffusion control, BA, were used, and the retention was measured in yeast cells (strain JK93da) expressing ABCB21, and ABCB1 and ABCB4 as controls. Bars represent the same as (b). (b) Cold IAA (10 μM) was pre-loaded and the import assay was conducted. Control 1 (pNEV) and control 2 (pDR196). (c) Export assay (low external IAA): these yeast transformants were pre-treated with IAA on ice and then export activity was measured (see the Materials and Methods for details). As diffusion control, BA is used. (d) Specific binding of NPA to microsomes prepared from yeast expressing ABCB21, ABCB1 or ABCB4. (e) NPA inhibition of IAA retention (import assays) in yeast expressing ABCB21 and its closest homolog, ABCB4. Data shown are mean activities ± SEs, 3–5 individual measurements with four replicates each. Significant differences from the vector control ( $P < 0.05$ ) are indicated by asterisks.

development. This suggests that the main phenotype of ABCB21 loss of function is the failure of seed development, and the minority of seeds that could somehow compensate the ABCB21 function, by as yet unidentified IAA transporters, can germinate. Thus, we analyzed the biochemical function of ABCB21 in the root using RNAi lines. The possibility cannot be eliminated that the RNAi construct for suppression of ABCB21 might influence the expression of other auxin transporters, but at least a microarray analysis of RNAi lines revealed that the expression of other ABC transporters was not strongly

influenced. While we could not find phenotype alterations in our experiments, our data strongly support that ABCB21 is a transporter regulating auxin movement in various tissues.

Conflicting reports on the transport directionality of ABCB4 have been reported (Terasaka *et al.* 2005, Cho *et al.* 2007, Lewis *et al.* 2007). Likewise, the closest ABCB4 homolog, ABCB21, seems to export IAA from mesophyll protoplast but, based on yeast assays, functions as an auxin importer (Figs. 5, 7). One possible explanation for this behavior might be the molecular environment of the assay systems where different

endogenous proteins may influence the transport function. For instance, the immunophilin-like protein TWISTED DWARF1 (TWD1) can modulate the function of ABCBs (Geisler et al. 2003, Bouchard et al. 2006), i.e. when co-expressed with ABCB1, TWD1 showed an opposite effect on ABCB1-mediated auxin efflux activity in yeast and HeLa cells (Bouchard et al. 2006). As another example, coordinated auxin transport by ABCB–PIN interactions has also been reported (Noh et al. 2003, Blakeslee et al. 2007), i.e. ABCB4 showed different auxin transport directionalities depending on its PIN partner (Cho et al. 2007). Obviously, these interacting partners are absent in the yeast expression system.

In this study carefully comparing ABCB1, B4 and B21 transport activity in yeast, we suggest another mechanism in that the cytoplasmic IAA concentration is a key factor determining the directionality of IAA transport (Figs. 5, 7): when cytoplasmic auxin concentrations are low (approximately fM order) such as in the yeast loading assays that were conducted in the absence of non-labeled IAA, both ABCB21 and B4, unlike ABCB1, import auxin (Fig. 7). However, when cytoplasmic auxin concentrations are kept relatively high (approximately nM order), either by addition of non-labeled IAA to the protoplast uptake buffer at time point 0 (Fig. 5b) or by yeast loading prior to removal of the extracellular supernatant during the export assays (Fig. 7c), both ABCB21 and B4 catalyze in the export direction like ABCB1. This proposed mechanism is supported by IAA pre-loading experiments that blocked ABCB21- and ABCB4-mediated IAA import (Fig. 7b). In that respect, ABCB21 resembles ABCB4 as a root-localized auxin efflux transporter with auxin uptake activity at low auxin concentrations (Kubeš et al. 2012).

This model is in line with the identification of putative cytoplasmic regulatory, structural elements in ABCB4 apparently absent in constitutive exporters, ABCB1 and B19 (Yang and Murphy 2009, Kubeš et al. 2012). Taken together, an ABCB protein may show such plasticity for the transport function depending on its subcellular expression altering cytoplasmic auxin concentrations. However, at this time, we cannot exclude the possibility that additional regulatory components, such as protein kinases or protein–protein interactions, further tune ABCB21 directionality (Cho et al. 2007, Yang and Murphy 2009).

## Materials and methods

### Plant material and growth conditions

*Arabidopsis thaliana* plants (ecotype Columbia) were grown on soil in growth chambers with 100 or 120  $\mu\text{mol m}^{-2} \text{s}^{-1}$  light in a 16 h light/8 h dark cycle at 21°C. The sterile growth condition was produced as described elsewhere (Terasaka et al. 2005). To analyze ABCB21 expression under various growth regulators, seeds were grown on a nylon mesh (20  $\mu\text{m}$  pore) over Murashige and Skoog (MS) medium for 14 d under the same light cycle described above. Roots were subjected to various

treatments by gentle transfer of the mesh to the new medium. Treatments were stopped by immediate freezing of seedlings in liquid N<sub>2</sub>. Sequence data for the genes in this study can be found in the GenBank/EMBL data libraries under the following accession numbers: PGP/ABCB21 (At3g62150), PGP/ABCB4 (At2g47000) and  $\beta$ -tubulin (At5g12250).

### Cloning of ABCB21 cDNA

The full-length coding sequence of ABCB21 cDNA (At3g62150) with 12 and 17 bp 5'- and 3'-untranslated regions, respectively, was isolated by RT–PCR with two primers, ABCB21-Full-Fw and ABCB21-Full-Rv (Supplementary Table S3). The PCR product was subcloned in the pENTR1A vector (Invitrogen).

### Expression analysis

Total RNA was isolated with the RNeasy Plant Mini-Kit (Qiagen). Reverse transcription was done with SuperScript III reverse transcriptase (Invitrogen), followed by incubation with RNase H (Invitrogen).

Real-time PCR was performed with the Roter-Gene 3000A (Corbett Research), using Platinum SYBR Green qPCR SuperMix-UDG (Invitrogen) according to the manufacturers' instructions. Briefly, the PCR mixture consisted of 10 ng of cDNA template, 5 pmol primers, 1  $\mu\text{l}$  of fluorescent probe provided with the above kit, and 12.5  $\mu\text{l}$  of Platinum Quantitative PCR SuperMix-UDG in a total volume of 25  $\mu\text{l}$ . The standard reaction condition was as follows: 95°C for 10 min, 40 cycles of 95°C for 15 s, 50°C for 30 s, 72°C for 30 s. The primers (ABCB21-RT-Fw and ABCB21-RT-Rv) are listed in Supplementary Table S3. Agarose gel electrophoresis, RNA transfers onto Hybond-N+ membranes (GE Healthcare) and hybridization with the 1.1 kb ABCB21 fragment (positions +2,001 to +3,063) were performed using standard procedures. The primers (ABCB21-northern-Fw and ABCB21-northern-Rv) are listed in Supplementary Table S3.

### Promoter:GUS construct and tissue-specific expression

To create the ABCB21 promoter:reporter construct (ProABCB21:GUS), two PCR primers flanking the regions from –753 to +15 bp were made (ABCB21Pro-Fw and ABCB21Pro-Rv) (Supplementary Table S3). PCR was performed using genomic DNA as a template and KOD-Plus polymerase (TOYOBO). The PCR product was subcloned into the entry vector pDONR221 to transfer the promoter region of ABCB21 into the binary vector pGWB203 by the Gateway system (Invitrogen). BP and LR reactions were performed according to standard procedures. *Agrobacterium tumefaciens* GV3101 (pMP90) was used for transformation via the floral dip method (Clough and Bent 1998). The T<sub>2</sub> generation were histochemically stained to detect GUS activity as described elsewhere (Terasaka et al. 2005). Tissue sections (10  $\mu\text{m}$  thick) embedded in Technovit 7100 (Heraeus Kulzer) were prepared by Leica microtome (RM2155) and observed with a Zeiss microscope (Axioskop2).



## Sucrose density gradient fractionation of plant membranes

Sucrose gradient fractionation was performed basically according to the method described by van den Brule *et al.* (2002) with the following modifications: Arabidopsis roots (10 g) were homogenized in 3 vols. of 50 mM HEPES-KOH, pH 7.5, 5 mM EDTA, 2 mM dithiothreitol (DTT), 250 mM sucrose and 1 mM phenylmethylsulfonyl fluoride (PMSF) using a mortar and pestle. The homogenate was centrifuged for 15 min to remove the debris, and the microsomal membrane fraction was collected by ultracentrifugation at  $100,000\times g$  for 40 min. The pellet was resuspended in 3 ml of gradient buffer (10 mM Tris-MES, pH 7.0, 1 mM DTT, 250 mM sucrose and 1 mM PMSF) and centrifuged at  $100,000\times g$  for 3 h on a non-continuous sucrose gradient from 20% to 45% (w/v) in the same buffer using a Beckman SW41Ti rotor. Gradient fractions were collected from the interface between different sucrose concentrations, and used for immunoblot analysis.

## Western blotting

A keyhole limpet hemocyanin conjugate of an oligopeptide of ABCB21, at position 1 (n-MDSVIESEGLKVDSNPRAD-c), was injected into rabbits according to a standard protocol (Hayashikasei Co. Ltd.). After the third boost, the antiserum was recovered and used for immunoblot analysis. The immunoblot procedure is as described previously (Yazaki *et al.* 2006). For each membrane marker, antibodies against plasma membrane  $H^+$ -ATPase (AHA), vacuolar  $H^+$ -pyrophosphatase (AVP1) and endoplasmic reticulum luminal BiP were used. For the Western blot of Fig. 5a, we used young whole plants.

## Transport experiments with protoplast

Intact Arabidopsis mesophyll protoplasts were prepared from rosette leaves of plants grown on soil under white light ( $100\ \mu\text{mol m}^{-2}\ \text{s}^{-1}$ , 8 h light/16 h dark, 22°C), and auxin import and export experiments were performed as described (Geisler *et al.* 2005). Briefly, intact protoplasts were isolated and loaded by incubation with  $1\ \mu\text{l ml}^{-1}$  [ $^3\text{H}$ ]IAA (specific activity  $7.4 \times 10^{11}\ \text{Bq mmol}^{-1}$ , American Radiolabeled Chemicals) and 4-[ $^3\text{H}$ ]NAA ( $9.3 \times 10^{11}\ \text{Bq mmol}^{-1}$ , American Radiolabeled Chemicals) in the presence of 100 nM IAA on ice. Import was started by incubation at 25°C and halted by silicon oil centrifugation. For export assays, loading was performed for 10 min on ice, allowing equal loading, and external radioactivity was removed by Percoll gradient centrifugation. Export was started by incubation at 25°C and halted by silicon oil centrifugation. Exported/imported radioactivity was determined by scintillation counting of the supernatant/protoplast interfaces and is presented as the relative export/import of the initial export/uptake (export/import prior to temperature incubation).

## Expression of ABCB21 in yeast and transport assay

The ABCB21 cDNA-pENTR1A was transferred into the destination vector, pDR196GW, which has a Gateway cassette

(*attR1-CmR*, *ccdB-attR2*) in the yeast shuttle vector pDR196 (Rentsch *et al.* 1995). The resulting plasmid, pDR196GW-ABCB21, was used to transform the yeast *gef1* (Gaxiola *et al.* 1998) or JK93da (Santelia *et al.* 2005) strains by the lithium acetate method (Ito *et al.* 1983). For IAA loading experiments, JK93da transformants were grown to  $\text{OD}_{600} = 1$ , washed and incubated at 30°C with  $1\ \mu\text{l ml}^{-1}$  5-[ $^3\text{H}$ ]IAA (specific activity  $7.4 \times 10^{11}\ \text{Bq mmol}^{-1}$ , American Radiolabeled Chemicals) and [ $^3\text{H}$ ]BA ( $9.3 \times 10^{11}\ \text{Bq mmol}^{-1}$ , American Radiolabeled Chemicals) in SD media (pH 5.5). For IAA efflux experiments, yeast cells were loaded at 0°C for 15 min with  $1\ \mu\text{l ml}^{-1}$  5-[ $^3\text{H}$ ]IAA (specific activity  $7.4 \times 10^{11}\ \text{Bq mmol}^{-1}$ , American Radiolabeled Chemicals) and [ $^3\text{H}$ ]BA ( $9.3 \times 10^{11}\ \text{Bq mmol}^{-1}$ , American Radiolabeled Chemicals) in SD media (pH 4.5), followed by two washing steps. Finally, export was started by 10 min incubation in SD media (pH 5.5) at 30°C. Aliquots of 1 ml were collected, filtered twice with cold water, and the retained radioactivity was quantified by scintillation counting. Some import assays were conducted with a pre-loading step with 10  $\mu\text{M}$  IAA for 10 min in SD media (pH 4.5) or in the presence of 10  $\mu\text{M}$  NPA in the loading media. NPA binding experiments were carried out as described in Kim *et al.* (2010).

## ABCB21 RNAi transformants

To prepare the RNAi construct, a gene-specific 240 bp fragment from ABCB21 cDNA (corresponding to nucleotide positions +1,908 to +2,148) was subcloned into the destination vector pGWB80 via the donor vector pDONR221 (Invitrogen). The PCR primers (ABCB21ir-Fw and ABCB21ir-Rv) used are listed in Supplementary Table S3. PCR and in vitro BP and LR recombinations were carried out as described above. Kanamycin-resistant  $T_3$  plants were used in the experiments.

## Supplementary data

Supplementary data are available at PCP online.

## Funding

This work was supported by The Ministry of Education, Culture, Sports, Science and Technology Grant-in-Aid for Scientific Research [17027016, 19060016, 20061018, and 23012024 (to K.Y.)]; the Swiss National Science Foundation, *Pole de Recherche* of the University of Fribourg and FOBI [all to M.G.]; the PM Project by NEDO [to S.D. and H.S.].

## Acknowledgments

We would like to thank Dr. W. Frommer (Carnegie Institution, Stanford, CA, USA) for pDR196, Dr. Tsuyoshi Nakagawa (Shimane University) for pGWB vectors, Dr. R.A. Gaxiola (University of Connecticut) for the yeast strain *gef1*, Dr. M. Boutry (Université Catholique de Louvain, Belgium) for anti- $H^+$ -ATPase antibodies, Dr. N. Koizumi (Osaka Prefecture

University, Japan) for anti-BiP antibodies, Dr. M.H. Sato (Kyoto Prefectural University, Japan) for anti-V-PPase antibodies, Dr. Hiroo Fukuda and Dr. Kuninori Iwamoto (The University of Tokyo, Japan) for ABCB21 Gene Chip technical support, and V. Vincenzetti (University of Zürich) and L. Charrier (University of Fribourg) for excellent technical support. DNA sequencing was conducted with the Life Research Support Center in Akita Prefectural University.

## References

- Bainbridge, K., Guyomarc'h, S., Bayer, E., Swarup, R., Bennett, M., Mandel, T. et al. (2008) Auxin influx carriers stabilize phyllotactic patterning. *Genes Dev.* 22: 810–823.
- Blakeslee, J.J., Bandyopadhyay, A., Lee, O.R., Mravec, J., Titapiwatanakun, B., Sauer, M. et al. (2007) Interactions among PIN-FORMED and P-glycoprotein auxin transporters in *Arabidopsis*. *Plant Cell* 19: 131–147.
- Bouchard, R., Bailly, A., Blakeslee, J.J., Oehring, S.C., Vincenzetti, V., Lee, O.R. et al. (2006) Immunophilin-like TWISTED DWARF1 modulates auxin efflux activities of *Arabidopsis* P-glycoproteins. *J. Biol. Chem.* 281: 30603–30612.
- Carraro, N., Forestan, C., Canova, S., Traas, J. and Varotto, S. (2006) *ZmPIN1a* and *ZmPIN1b* encode two novel putative candidates for polar auxin transport and plant architecture determination of maize. *Plant Physiol.* 142: 254–264.
- Casimiro, I., Marchant, A., Bhalerao, R.P., Beeckman, T., Dhooge, S., Swarup, R. et al. (2001) Auxin transport promotes *Arabidopsis* lateral root initiation. *Plant Cell* 13: 843–852.
- Cho, M., Lee, S.H. and Cho, H.T. (2007) P-glycoprotein4 displays auxin efflux transporter-like action in *Arabidopsis* root hair cells and tobacco cells. *Plant Cell* 19: 3930–3943.
- Clough, S.J. and Bent, A.F. (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* 16: 735–743.
- Gaxiola, R.A., Yuan, D.S., Klausner, R.D. and Fink, G.R. (1998) The yeast CLC chloride channel functions in cation homeostasis. *Proc. Natl Acad. Sci. USA* 95: 4046–4050.
- Geisler, M., Blakeslee, J.J., Bouchard, R., Lee, O.R., Vincenzetti, V., Bandyopadhyay, A. et al. (2005) Cellular efflux of auxin catalyzed by the *Arabidopsis* MDR/PGP transporter AtPGP1. *Plant J.* 44: 179–194.
- Geisler, M., Kolukisaoglu, H.U., Bouchard, R., Billion, K., Berger, J., Saal, B. et al. (2003) TWISTED DWARF1, a unique plasma membrane-anchored immunophilin-like protein, interacts with *Arabidopsis* multidrug resistance-like transporters AtPGP1 and AtPGP19. *Mol. Biol. Cell.* 14: 4238–4249.
- Geisler, M. and Murphy, A.S. (2006) The ABC of auxin transport: the role of p-glycoproteins in plant development. *FEBS Lett.* 580: 1094–1102.
- Himanen, K., Boucheron, E., Vanneste, S., Engler, J.A., Inzé, D. and Beeckman, T. (2002) Auxin-mediated cell cycle activation during early lateral root initiation. *Plant Cell* 14: 2339–2351.
- Ito, H., Fukuda, Y., Murata, K. and Kimura, A. (1983) Transformation of intact yeast cells treated with alkali cations. *J. Bacteriol.* 153: 163–168.
- Jasinski, M., Ducos, E., Martinoia, E. and Boutry, M. (2003) The ATP-binding cassette transporters: structure, function, and gene family comparison between rice and *Arabidopsis*. *Plant Physiol.* 131: 1169–1177.
- Kim, J.Y., Henrichs, S., Bailly, A., Vincenzetti, V., Sovero, V., Mancuso, S. et al. (2010) Identification of an ABCB/P-glycoprotein-specific inhibitor of auxin transport by chemical genomics. *J. Biol. Chem.* 285: 23309–23317.
- Kubeš, M., Yang, H., Richter, G.L., Cheng, Y., Młodzińska, E., Wang, X. et al. (2012) The *Arabidopsis* concentration-dependent influx/efflux transporter ABCB4 regulates cellular auxin levels in the root epidermis. *Plant J.* 69: 640–654.
- Laskowski, M.J., Williams, M.E., Nusbaum, H.C. and Sussex, I.M. (1995) Formation of lateral root meristems is a two-stage process. *Development* 121: 3303–3310.
- Luschnig, C. (2002) Auxin transport: ABC proteins join the club. *Trends Plant Sci.* 7: 329–332.
- Lewis, D.R., Miller, N.D., Splitt, B.L., Wu, G. and Spalding, E.P. (2007) Separating the roles of acropetal and basipetal auxin transport on gravitropism with mutations in two *Arabidopsis* multidrug resistance-like ABC transporter genes. *Plant Cell* 19: 1838–1850.
- Marchant, A., Kargul, J., May, S.T., Müller, P., Delbarre, A., Perrot-Rechenmann, C. et al. (1999) AUX1 regulates root gravitropism in *Arabidopsis* by facilitating auxin uptake within root apical tissues. *EMBO J.* 18: 2066–2073.
- Murphy, A.S., Hoogner, K.R., Peer, W.A. and Taiz, L. (2002) Identification, purification, and molecular cloning of N-1-naphthylphthalamic acid-binding plasma membrane-associated aminopeptidases from *Arabidopsis*. *Plant Physiol.* 128: 935–950.
- Noh, B., Bandyopadhyay, A., Peer, W.A., Spalding, E.P. and Murphy, A.S. (2003) Enhanced gravi- and phototropism in plant *mdr* mutants mislocalizing the auxin efflux protein PIN1. *Nature* 423: 999–1002.
- Noh, B., Murphy, A.S. and Spalding, E.P. (2001) Multidrug resistance-like genes of *Arabidopsis* required for auxin transport and auxin-mediated development. *Plant Cell* 13: 2441–2454.
- Park, S., Li, J., Pittman, J.K., Berkowitz, G.A., Yang, H., Undurraga, S. et al. (2005) Up-regulation of a H<sup>+</sup>-pyrophosphatase (H<sup>+</sup>-PPase) as a strategy to engineer drought-resistant crop plants. *Proc. Natl Acad. Sci. USA* 102: 18830–18835.
- Rentsch, D., Laloï, M., Rouhara, I., Schmelzer, E., Delrot, S. and Frommer, W.B. (1995) *NTR1* encodes a high affinity oligopeptide transporter in *Arabidopsis*. *FEBS Lett.* 370: 264–268.
- Santelia, D., Vincenzetti, V., Azzarello, E., Bovet, L., Fukao, Y., Düchtig, P. et al. (2005) MDR-like ABC transporter AtPGP4 is involved in auxin-mediated lateral root and root hair development. *FEBS Lett.* 579: 5399–5406.
- Schnabel, E.L. and Frugoli, J. (2004) The PIN and LAX families of auxin transport genes in *Medicago truncatula*. *Mol. Genet. Genomics* 272: 420–432.
- Terasaka, K., Blakeslee, J.J., Titapiwatanakun, B., Peer, W.A., Bandyopadhyay, A., Makam, S.N. et al. (2005) PGP4, an ATP binding cassette P-glycoprotein, catalyzes auxin transport in *Arabidopsis thaliana* roots. *Plant Cell* 17: 2922–2939.
- van den Brûle, S., Müller, A., Fleming, A.J. and Smart, C.C. (2002) The ABC transporter SpTUR2 confers resistance to the antifungal diterpene sclareol. *Plant J.* 30: 649–662.
- Vanneste, S., Rybel, B.D., Beemster, G.T.S., Ljung, K., Smet, I.D., Isterdael, G.V. et al. (2005) Cell cycle progression in the pericycle is not sufficient for SOLITARY ROOT/JAA14-mediated lateral root initiation in *Arabidopsis thaliana*. *Plant Cell* 17: 3035–3050.

- Verrier, P.J., Bird, D., Burla, B., Dassa, E., Forestier, C., Geisler, M. et al. (2008) Plant ABC proteins—a unified nomenclature and updated inventory. *Trends Plant Sci.* 13: 151–159.
- Wiśniewska, J., Xu, J., Seifertová, D., Brewer, P.B., Růžička, K., Blilou, I. et al. (2006) Polar PIN localization directs auxin flow in plants. *Science* 312: 883.
- Xu, M., Zhu, L., Shou, H. and Wu, P. (2005) A *PIN1* family gene, *OsPIN1*, involved in auxin-dependent adventitious root emergence and tillering in rice. *Plant Cell Physiol.* 46: 1674–1681.
- Yang, H. and Murphy, A.S. (2009) Functional expression and characterization of Arabidopsis ABCB, AUX 1 and PIN auxin transporters in *Schizosaccharomyces pombe*. *Plant J.* 59: 179–191.
- Yazaki, K., Yamanaka, N., Masuno, T., Konagai, S., Shitan, N., Kaneko, S. et al. (2006) Heterologous expression of a mammalian ABC transporter in plant and its application to phytoremediation. *Plant Mol. Biol.* 61: 491–503.