

A Randomized Controlled Trial of the Effectiveness of Computer-Assisted Cognitive Remediation (CACR) in Adolescents with Psychosis or at High Risk of Psychosis

Laurent Holzer, Sebastien Urben, Christina Moses Passini and Laure Jaughey

Service Universitaire de Psychiatrie de l'Enfant et de l'Adolescent, Lausanne, Switzerland

Michael H. Herzog

Ecole Polytechnique Fédérale de Lausanne (EPFL), Switzerland

Olivier Halfon and Sandrine Pihet

Service Universitaire de Psychiatrie de l'Enfant et de l'Adolescent, Lausanne, Switzerland

Background: Computer assisted cognitive remediation (CACR) was demonstrated to be efficient in improving cognitive deficits in adults with psychosis. However, scarce studies explored the outcome of CACR in adolescents with psychosis or at high risk. **Aims:** To investigate the effectiveness of a computer-assisted cognitive remediation (CACR) program in adolescents with psychosis or at high risk. **Method:** Intention to treat analyses included 32 adolescents who participated in a blinded 8-week randomized controlled trial of CACR treatment compared to computer games (CG). Cognitive abilities, symptoms and psychosocial functioning were assessed at baseline and posttreatment. **Results:** Improvement in visuospatial abilities was significantly greater in the CACR group than in CG. Other cognitive functions, psychotic symptoms and psychosocial functioning improved significantly, but at similar rates, in the two groups. **Conclusion:** CACR can be successfully administered in this population; it proved to be effective over and above CG for the most intensively trained cognitive ability.

Keywords: Psychosis, adolescent, cognitive remediation, computer, controlled trial.

Introduction

Cognitive impairments are recognized as a core feature of schizophrenia also present in other psychotic disorders, as well as in adolescents and patients at high risk of psychosis. In view of their stability over time, their independence of other symptoms, their clear

impact on functional outcome, and poor alleviation by available psychopharmacological treatments (Palmer, Dawes and Heaton, 2009), cognitive impairments constitute a key target for additional intensive treatments.

In this perspective, cognitive remediation therapy, which can be defined as “a behavioural training-based intervention that aims to improve cognitive processes (attention, memory, executive function, social cognition or metacognition) with the goal of durability and generalization” (Wykes, Huddy, Cellard, McGurk and Czobor, 2011, p. 472). Cognitive remediation therapy was demonstrated to be effective at enhancing cognitive functioning (see meta-analyses of McGurk, Twamley, Sitzer, McHugo and Mueser, 2007 and Wykes et al., 2011).

In particular, cognitive enhancement therapy represents an evidence-based developmental cognitive rehabilitation approach for enhancing cognitive functioning (Uhlmann and Swanson, 2004). More specifically, Eack and colleagues (e.g. Eack et al., 2009, 2010) conducted, in a young adult sample suffering from schizophrenia or schizoaffective disorders, a 2-year randomized controlled trial comparing cognitive enhancement therapy to enriched supportive therapy (illness management and psychoeducation approach). Results indicated positive effects on social cognition, cognitive style, social adjustment and symptomatology as a result of cognitive enhancement therapy. These positive effects were shown to be sustained either at 1-year follow-up (Anderson, 2004) or at 2-year follow up (Eack et al., 2009). Furthermore, Eack et al. (2010) demonstrated that this type of treatment might have a neuroprotective effect as they showed a greater preservation of grey matter volume after 2 years of illness.

Another well-developed form of cognitive remediation is the computer-assisted cognitive remediation (CACR), which provides a standardized training with immediate feedback adapted to suit psychotic patients (Medalia, Aluma, Tryon and Merriam, 1998). A variety of CACR programs improve cognitive deficits in adult psychotic patients, with smaller size effects on symptoms and psychosocial functioning (Grynszpan et al., 2011; McGurk et al., 2007; Wykes et al., 2011), and some new adaptations are currently under evaluation (NEUROCOM trial; Wykes, Reeder, Corner, Williams and Everitt, 1999).

Treatment of cognitive impairments during adolescence, a period of high brain plasticity, may reduce disabilities in adulthood associated with early-onset psychosis. There is a general lack of research on cognitive remediation in adolescents (Wykes et al., 2011). A study observed (non significantly) larger improvements in cognitive functioning, psychiatric symptoms and psychosocial functioning in adolescents with early-onset psychosis receiving cognitive remediation training, compared to a control group (Ueland and Rund, 2004); and Wykes et al. (2007) found significantly larger improvements in the cognitive remediation group for cognitive flexibility only, in young early onset patients with schizophrenia. Note that these programs were not computer-assisted. Many arguments could be offered in favour of CACR, more specifically when working with adolescents. Indeed, the use of computerized technology is an everyday reality related to self-perceived competence that enhances the probability to engage in this form of cognitive remediation (Bremer and Rauch, 1998). In addition, computer activities were thought to improve chances to acquire new compensatory strategies, an important component of CRT (Kurtz, Seltzer, Shagan, Thime and Wexler, 2007). Finally, prolonged multimedia stimulation is believed to favour neural plasticity (Hogarty et al., 2004). Therefore, the effectiveness and feasibility of a specific CACR program need to be confirmed in adolescents with psychosis or presenting a high risk. In such a case, a CACR

program could easily be applied and generalized to everyday clinical practice, thus offering the possibility of being highly beneficial for adolescent health care.

In the current study, it was hypothesized that adolescents with psychosis or with high risk would be able to successfully complete the CACR program and would show significant improvements on cognitive tasks, negative and positive symptoms, and psychosocial functioning, compared to participants in the control condition who played computer games (CG). Here we present the results following a CACR program. However, the results of the 6-month follow-up have already been presented in detail elsewhere (Urban, Pihet, Jaugey, Halfon and Holzer, 2012). We can summarize the results as follows: with regard to the cognitive abilities no amelioration was found in the control group, while in the CACR group, significant improvements in inhibition and reasoning abilities were observed. Furthermore, symptoms were observed to decrease significantly in the control group and marginally in the CACR group. Finally, the enhancements in cognitive abilities were not related to the amelioration of symptoms.

Method

Sample

Inclusion criteria were: (1) diagnosis of psychotic disorder according to the DSM-IV (APA, 1994) using the French version of *Diagnostic Interview for Genetic Studies* (DIGS; Nurnberger et al., 1994; Preisig, Fenton, Matthey, Berney and Ferrero, 1999) or diagnosis of at high risk of psychosis using the Structured Interview for Prodromal Symptoms (SIPS) and the Scale of Prodromal Symptoms (SOPS; Miller et al., 1999); (2) score below the 10th percentile in at least one of five domains of the Repeatable Battery for the Assessment of Neuropsychological Status (RBANS; Randolph, Tierney, Mohr and Chase, 1998). The 10th percentile best differentiated patients with psychotic disorders from patients with other diagnoses (Holzer et al., 2007). Exclusion criteria included: (1) mental retardation (IQ < 70), defined as the need for special education, assessed through a screening of the medical records about the activities before the enrolment in the study; (2) known neurological disease or developmental disability; (3) severe visual or motor disorder incompatible with computer use; (4) transient exclusion criteria: an acute clinical state that could disrupt the training, or a planned absence for more than 2 weeks during the period of intervention. Figure 1 presented the flow diagram.

Thirty-two adolescents ($n = 20$ psychotic; $n = 12$ at risk) were randomized to CACR ($n = 18$) or CG ($n = 14$); 28 participants completed the study (15 in CACR, 13 in CG). As presented in Table 1, groups did not differ significantly in age, gender, ethnicity, number of school years completed, proportion of at risk participants, duration of illness, duration of untreated psychosis, and medication.

Procedure

A blinded 8-week trial of CACR treatment was compared to CG, with random assignment to groups, and assessments at baseline and post-intervention (week 9) of primary (cognitive abilities) and secondary (symptoms and psychosocial functioning) outcomes. Thus, the




Figure 1. (Colour online) Flow diagram of the study (n = whole sample, n with psychosis/ n at risk)

present study followed the guidelines regarding the complex intervention (Medical Council Research, www.mrc.ac.uk/complexinterventionsguidance)

Approval was received from the local ethics committee for human research, and informed consent was obtained from participants and their guardians. Participants were recruited from the Day Care Unit for Adolescents (DCUA) in Lausanne, Switzerland, while they were outpatients. The DCUA of Lausanne accommodates 15–20 adolescents (age ranging from 13 to 18 years) presenting with psychosis (more than half of the patients), mood disorders, anxiety disorders and conduct disorders. The mean duration of stay is about 5 months and the usual treatment program encompasses individual medical and psychological follow-up with special-school attendance, and occupational and work therapy. No longer able to attend school or apprenticeships, adolescents who are admitted (only after medical indication) neither required acute treatment in an inpatient unit nor could rely on a simple outpatient-clinic setting. Despite their relatively severe psychiatric disorders, adolescents attending the DCUA

Table 1. Demographic and clinical characteristics, and treatment compliance of the Computer Assisted Cognitive Remediation (CACR) and Computer Games (CG) groups

	CACR (<i>n</i> = 18)	CG (<i>n</i> = 14)	Group comparison test ¹
Age, mean years (<i>SD</i>)	15.4 (1.3)	15.7 (1.4)	$Z = 1.38, p = .442$
Gender, % (<i>n</i>) male	50 (9)	64 (9)	$\chi^2_{(1)} = 0.74, p = .328$
Ethnicity, % (<i>n</i>) caucasian	89 (16)	79 (11)	$\chi^2_{(1)} = 2.49, p = .357$
Education, mean years (<i>SD</i>)	7.8 (1.2)	8.3 (1.4)	$Z = 0.10, p = .283$
High risk of psychosis, % (<i>n</i>)	39 (7)	36 (5)	$\chi^2_{(1)} = 0.44, p = .547$
Duration of illness, mean months (<i>SD</i>)	31.8 (34.3)	33.2 (36.1)	$Z = 0.10, p = .892$
Untreated psychosis <2 months, % (<i>n</i>) ²	50 (6)	55 (6)	
Untreated psychosis ≤1 year, % (<i>n</i>) ²	33 (4)	18 (2)	
Untreated psychosis >1 year, % (<i>n</i>) ²	17 (2)	27 (3)	$\chi^2_{(2)} = 0.83, p = .662$
No antipsychotic medication, % (<i>n</i>)	40 (7)	40(7)	
Atypical antipsychotic, % (<i>n</i>)	55 (10)	43 (6)	
Typical antipsychotic, % (<i>n</i>)	5 (1)	7 (1)	$\chi^2_{(2)} = 0.14, p = .844$
Missed sessions, mean (<i>SD</i>) ³	2.1 (3.4)	1.3 (2.4)	$Z = 0.32, p = .786$
Session length, mean minutes (<i>SD</i>) ³	45.8 (7.9)	48.4 (7.5)	$Z = 0.95, p = .354$
Program duration, mean days (<i>SD</i>) ³	93.5 (33.4)	93.5 (34.2)	$Z = 0.23, p = .818$
Motivation, mean (<i>SD</i>) ³	4.1 (0.7)	4.5 (0.5)	$Z = 1.75, p = .079$

¹For gender, ethnicity, high risk, untreated psychosis and medication: Chi-square test; for other variables: Mann-Whitney test.

²Missing information for 9 patients (6 in CACR and 3 in CG).

³Data provided only for the participants who finished the program ($N = 15$ in CACR and 13 in CG).

are clinically sufficiently stable to engage themselves in computerized task sessions and their presence 7 hours a day, 5 days a week, represents great availability for training sessions.

Clinical assessment was performed by LH and a senior child and adolescent psychiatrist, who was blind to group assignment during the study. Neuropsychological assessment was performed by one of the two neuropsychologists blind to diagnostic status at baseline and blind to group assignment during the study. After informed consent was given by both the patients and their parents, each adolescent was randomly assigned to the CACR group or the control group (videogames). A computer-generated randomization list was drawn up by the statistician. The group assignment was known only by the CACR trainer and videogames provider (and the adolescent). To ensure balance between groups during the trial, a blocked randomization was used. Thus randomization was completed by the statistician, using blocks of four patients with identical diagnoses (at risk versus with psychosis), with a 1:1 allocation ratio.

Treatment: computer-assisted cognitive remediation (CACR)

The original Captain's Log[®] software (see Sanford and Brown, 1988) consists of five modules; the software consists of 35 multi-level "brain-training" exercises designed to develop and remediate attention, concentration, memory, eye-hand coordination, basic numeric concepts, problem-solving/reasoning skills, self-esteem and self-control. The six modules encompass: (a) Attention Skills: Developmental (eight programs) is designed to

train attention – general, alternating, focused and sustained – visual and auditory processing speed, response inhibition, visual scanning, categorization, and working memory; (b) Visual Motor Skills comprises seven programs to train eye-hand coordination, visual scanning, visual tracking, alternating and divided attention, fine motor control, response inhibition, and processing speed. As in the first module (Attention Skills: Developmental) the programs offer appropriate presentations for children, teenagers and adults; (c) Conceptual Skills consists of seven effective programs designed to train basic reasoning, short-term and working memory, perceptual discrimination, sequencing and categorization; (d) Numeric Concepts/Memory Skills consists of five effective programs designed to train basic reasoning, numeric skills, short-term and working memory, perceptual discrimination, sequencing and categorization; (e) Attention Skills: The Next Generation presents challenging exercises to develop higher level cognitive skills – auditory attention and discrimination, listening skills, divided attention, visual scanning, short-term memory, and faster mental processing speed. All of the three programs can also be used to develop problem-solving/reasoning skills; (f) Logic Skills is a new module consisting of five programs that focus on higher level executive functioning, organization, categorization, pattern recognition, sequencing and closure.

As the whole program is likely to provide more than 500 hours of cognitive training, a selection of specific tasks is needed. In order to limit variation in remediation tasks that might hamper comparability and generalization of findings, we selected a limited number of tasks to be administered to all patients in a standardized manner. Selection was inspired by the Bellucci, Glaberman and Haslam study (2003). Out of 35 possible tasks, 12 were selected for the training program. From the “Attention skills: developmental” module we selected “auditory discrimination/rhythm” (trains working memory, auditory processing speed, sustained attention), “colour discrimination/inhibition” (trains visual scanning, response inhibition, general attention, central processing speed and working memory); from the “Visual motor skills” module we selected “visual timing” (trains fine motor control and visual perception), “visuospatial memory concentration” (trains visuospatial categorization and general attention), “visual tracking/discrimination” (trains visual tracking and visual perception); from the Conceptual skills module we selected “conceptual discrimination” (trains the conceptual abilities of perception, classification and recognition), “size discrimination” (trains selected attention, visual tracking skills), “symbolic display match” (develops complex conceptual reasoning and processing speed); from the “Numeric concept/memory skills” we chose “numeric classifications” (trains visuospatial classification and perception, working memory, general attention), “numeric distinctions” (trains visuospatial sequencing, conceptual reasoning, working memory, immediate memory), and from the “Attention skills: the next generation” module, “symbol search” (trains processing speed, visual scanning, self-control, short term memory); finally, from the Logic skills module “sequential logic” (develops conceptual reasoning and visuospatial sequencing) was selected.

Nearly all of the exercises in Captain’s Log are non-language-based (apart from the task instructions). The trainer selected the tasks and provided the directions for the patient and assisted him/her during all the sessions with encouragement and positive feedback. The Captain’s Log program began with an assessment phase and starts training for all people at the same difficulty level. Progression is based on the person’s level of skill and speed of learning, with an increase in scores required before the next difficulty level is reached. Therefore, progress through levels is determined by the program, rather than by trainers. To summarize, participants received 16 45-minute individual sessions, with a frequency of two

sessions per week for 8 weeks. Research psychologists guiding the CACR selected the tasks to match the cognitive deficits of the participant,¹ translated the instructions to the participant (as exercises are not language-based, except for the instructions, the original English version was used), and provided encouraging and positive feedback.

Control: computer game (CG)

A set of various videogames² (essentially action videogames that require attention and visuo-motor skills) was offered to patients assigned to the control group with two half-hour sessions weekly for 8 weeks. Violent videogames were avoided as deleterious influences on aggressiveness have been described (Anderson, 2004; Uhlmann and Swanson, 2004). The videogames program differed from the CACR program only in content (videogames instead of Captain's Log software) while the setting was the same (location, computer duration, frequency, trainer). The same trainer accompanied the patient during all the sessions. The trainer selected videogames and provided encouragement and positive feedback. The research psychologists provided a supportive climate during gaming. CG and CACR sessions took place in the same room, were conducted by the same psychologists, and had similar duration and frequency (see Compliance for details).

Measures

Primary outcomes. Cognitive functioning was measured by the Repeatable Battery for the Assessment of Neuropsychological Status (RBANS; Randolph et al., 1998), which has two psychometrically equivalent alternate forms: Intraclass correlation for the test-retest reliability for the total score is .83 and ranges from .51 to .72 for the domain scores. The RBANS assesses five domains: (1) immediate memory; (2) visuospatial/constructional; (3) language; (4) attention; (5) delayed memory. An adapted version of the RBANS for use in adolescents was employed (Holzer et al., 2007). For most of the patients, the same trained neuropsychologist conducted the baseline and post-intervention assessments.

Secondary outcomes. Symptoms were assessed with the Positive and Negative Symptom Scale (PANSS; Kay, Fiszbein and Opler, 1987), comprising three subscales (positive symptoms; negative symptoms; general psychopathology). Psychosocial functioning was evaluated by the Social and Occupational Functioning Assessment Scale (SOFAS; APA, 1994) and the Health of Nation Outcome Scale for Children and Adolescents (HoNOSCA; Gowers et al., 1999).

¹All patients presented at least three impaired cognitive abilities, identified as treatment targets, based on a complete neuropsychological assessment; given that all Captain's Log exercises simultaneously train a set of cognitive abilities (e.g. for the first exercise: working memory, auditory processing speed and focused attention), each participant received a unique combination of exercises matching his/her impairments and preferences.

²Computer games did not include any specific educational or strategic content; for example, the three most played games were "Chicken Little", "Tetris", and "Sonic".

Engagement in treatment. For each participant, at the end of each session, trainers rated: 1) general motivation for treatment; and 2) engagement in the training tasks, using a 5-point scale ranging from 1 (very low) to 5 (very high). The two items were then averaged (Cronach's $\alpha = .90$) into a single motivation score. The effective work time (EWT, i.e. time spent training or playing on the computer, outside explanations or discussion with the trainer, in minutes) was also recorded.

Statistical analysis

The analyses were performed with SPSS (version 20) software. We adopted an intention to treat (ITT) analyses in order to include the 32 participants who were randomly assigned either to the CACR or CG groups. According to the guideline proposed by Howell (2008), we first analysed the pattern of the missing data, with Little's MCAR test revealing that the missing data could be considered as missing completely at random (MCAR, $\chi^2(57) = 70.77, p = .104$). We could thus estimate the missing points with the regression procedure included in the SPSS software. We therefore analysed the completed data of the 32 participants for the treatment effects analyses.

Given the presence of significant deviations from normality, non-parametric tests were used. Change scores from baseline to post-intervention were computed for cognitive functions, symptoms and psychosocial functioning so that a positive score indicates improvement and a negative one deterioration. Group comparisons were conducted on change scores using Mann-Whitney tests. Baseline to post-intervention progresses were tested with Wilcoxon signed rank tests on the 32 randomized participants constituting the ITT sample.

Study's statistical power

First, the number of participants was chosen, in keeping with the only two previous studies exploring the outcome of cognitive remediation on young people. Wykes and colleagues (2007) enrolled 31 participants, and 25 adolescents participated in the Ueland and Rund (2004) study, so with 32 participants included in the ITT analyses, we have more participants than previous studies exploring the outcome of cognitive remediation in young people with schizophrenia.

Second, a power analyses was conducted (with G*Power 3.1.3 software) to compute the probability of observing large size effect (those with clinical significance as observed in Bellucci et al., 2003) on the cognitive performances and clinical status of the ITT sample ($N = 32$) for two-tails tests at a level of significance of .05. A power above .8 could be considered sufficient to detect possible changes (Cohen, 1992). These analyses were computed regarding the assessment of the treatment outcomes on the primary measures (cognitive assessment: RBANS scores) and secondary measures (clinical evaluations: PANSS, SOFAS and HoNOSCA scores). The Mann-Whitney tests (assessing the group effect on the change scores of the cognitive and clinical measures, post-intervention minus baseline scores) had a power ($1-\beta$) to detect changes of .56, which is not acceptable. In contrast, the power ($1-\beta$) of the Wilcoxon (estimating time effect on the scores of the baseline compared to the post-intervention) is of .98 to detect changes, which is acceptable. Furthermore, 2 (time: baseline vs post-intervention) by 2 (groups: CACR vs CG) analyses of variance (ANOVA) allowing to assess the interaction of time and group effect had a power of .88, which could be

considered as acceptable. So, in this context, all negative results of the Mann-Whitney tests (those perhaps due to a lack of statistical power) were re-examined by the interaction effect in the corresponding ANOVA in order to prevent type II error.

Results

Treatment compliance

Four patients interrupted their participation in the program: 3 in the CACR group (two due to lack of interest and one due to transfer to another care centre), and 1 in the CG group (due to poor attendance at the DCUA, resulting in transfer). They differed significantly from program completers only on having finished less school years (completers: $mean = 8.2$, $SD = 1.1$; dropouts: $mean = 6.3$, $SD = 1.3$; $Z = 2.52$, $p = .012$).

Treatment compliance was overall very high and similar in both groups (see Table 1): on average, only 2.3 sessions were missed (but later attended), sessions lasted the expected 45 minutes (47.2 min), and participants took around 3 months (92.4 days) to complete the full program. Motivation was also high (4.3), with a marginal advantage to the CG group ($p = .079$), probably due to the playful nature of the task and the free choice of games given to participants. Overall, the acceptance of the CACR intervention was therefore very satisfactory.

Equivalence between groups at baseline

The 32 participants constituting the ITT sample were compared on baseline measures in function of treatment. Mann-Whitney tests revealed no significant differences (see Table 2), apart from lower language abilities for the CACR group. Given that language was not part of the trained cognitive abilities and neither CACR exercises nor CG were language-based, we considered these differences unlikely to bias the assessment of potential effects.

Treatment effects

ITT analyses ($N = 32$) were conducted in order to assess differential treatment effects by group. Thus, Mann-Whitney tests were conducted on the change scores for RBANS, PANSS, SOFAS and HoNOSCA (baseline and post-intervention scores are presented in Table 2). Visuospatial abilities improved significantly more in CACR ($mean = +4.9$, $SD = 10.3$) than in CG patients ($mean = -3.8$, $SD = 16.0$; $Z = 2.47$, $p = .013$), corresponding to a large effect size ($d = 0.62$). This effect was found only in 47% of the patients in the CACR group, and 39% in the CG group³ scored below the 10th percentile on visuospatial abilities at baseline. No other significant group differences were found.

Given that improvement did not differ between groups on most variables, baseline and post-intervention scores were then compared using Wilcoxon signed rank tests on the ITT sample ($N = 32$). As reported in Table 2, improvements were significant for attention, immediate and delayed memory, general psychopathology, and social-occupational functioning. Both groups thus improved on most measures.

³Computed on the sample who finished the study ($N = 28$).

Table 2. Mean and (*SD*) of study variables at baseline and post-intervention for the Computer Assisted Cognitive Remediation (CACR) and Computer Games (CG) groups, and tests of group differences at baseline and of time effects

Variable	CACR group (n = 18)		CG group (n = 14)		Group diff. at baseline ¹	Time effect ²	Group diff. on changes scores ³
	Baseline	Post-interv.	Baseline	Post-interv.			
RBANS-Total	78.8 (8.8)	83.6 (9.4)	85.7 (9.7)	92.3 (14.4)	1.90	2.86**	1.03
RBANS-IM	87.9 (10.3)	93.6 (9.5)	93.3 (13.2)	101.5 (19.6)	1.18	2.68*	0.81
RBANS-VC	87.9 (15.4)	92.6 (13.2)	96.3 (17.2)	92.5 (13.4)	1.30	0.22	2.47*
RBANS-L	84.2 (12.0)	85.1 (19.8)	95.4 (13.6)	98.5 (14.9)	2.31*	0.43	0.53
RBANS-A	74.2 (11.6)	79.8 (15.9)	67.4(12.3)	81.7 (16.8)	1.43	2.76**	1.68
RBANS-DM	82.1 (17.31)	92.0 (16.4)	90.0 (20.6)	96.2 (18.7)	1.37	2.65**	0.82
PANSS-Total	67.8 (19.4)	64.6 (17.5)	70.1 (23.7)	60.9 (22.7)	0.15	2.79**	1.33
PANSS-PS	14.1 (6.2)	13.7 (5.0)	13.7 (5.9)	12.4 (4.9)	0.04	1.55	0.55
PANSS-NS	18.9 (8.1)	17.8 (7.9)	18.9 (7.7)	16.6 (8.6)	0.07	1.94	0.82
PANSS-GP	34.2 (8.8)	32.6 (11.0)	37.5 (13.0)	31.4 (12.1)	0.62	2.65*	1.71
SOFAS	40.83 (20.1)	52.1 (9.3)	42.2 (15.8)	52.8 (10.7)	0.76	2.93*	0.80
HoNOSCA	14.5 (7.4)	18.2 (4.8)	15.8 (8.8)	16.2 (6.0)	0.80	1.13	0.91

RBANS: Repeatable Battery for the Assessment of Neuropsychological Status; IM: immediate memory; VC: visuospatial/constructional, L: language; A: attention; DM: delayed memory; PANSS: positive and negative symptom scale; PS: positive symptoms; NS: negative symptoms; GP: general psychopathology; SOFAS: Social and Occupational Functioning Assessment Scale; HoNOSCA: Health of Nation Outcome Scale for Children and Adolescents

¹Mann-Withney test (Z) comparing CACR and CG on baseline scores; * $p < .05$, ** $p < .01$

²Wilcoxon test (Z) comparing baseline and post-intervention scores for the whole sample; * $p < .05$, ** $p < .01$

³Mann-Withney test (Z) comparing CACR and CG on change scores; * $p < .05$, ** $p < .01$

Subsequent analyses were conducted on the sample of participants who completed the trial, in order not to bias the analysis. One reason why visuospatial abilities may have improved more in CACR than in CG could be that they were particularly intensively trained in CACR exercises. To test this, we calculated for each participant the proportion of CACR exercises that involved each cognitive function. Indeed, on average, visual and visuospatial abilities were trained by 97% of the performed exercises ($SD = 10.2$), corresponding to the highest training intensity with attention (94% of exercises, $SD = 9.2$). This constitutes a significantly (Friedman test: $\chi^2_{(5)} = 53.7, p < .001$) more intensive training than for memory (66%, $SD = 12.7$), processing speed (55%, $SD = 18.8$), response inhibition (49%, $SD = 15.2$), and conceptual reasoning (63%, $SD = 16.9$), supporting our hypothesis.

We further examined the associations (using Spearman's rank correlation³) between cognitive improvements and motivation or compliance: larger gains in attention were significantly related to longer sessions ($\rho = .53, p = .044$) and higher motivation ($\rho = .67, p = .007$), the latter also being nonsignificantly related to progress in visuospatial abilities ($\rho = .31, p = .266$) and delayed memory ($\rho = .41, p = .130$). This result suggests that motivation for CACR treatment enhanced cognitive change. No significant correlations emerged in the CG group.

Discussion and conclusion

The findings with respect to the feasibility of CACR program are encouraging. All but three of the participants completed the program, attending the sessions with high motivation. The results demonstrated a superior improvement in CACR for the most intensively trained cognitive function, visuospatial abilities, and gains in both the control and treatment group for attention, immediate and delayed memory, and general psychopathology, as well as social-occupational functioning. A more intense CACR training might even further improve CACR over CG, consistent with previous evidence highlighting the importance of treatment intensity (McGurk et al., 2007; Medalia and Richardson, 2005). Although duration of intervention was not found to predict efficacy in the latest meta-analyses (Grynszpan et al., 2011; Wykes et al., 2011), according to a recent review, "30–40 hours of training and 3 months of trial duration is viewed as a minimum" (Keefe et al., 2011; p. 1059). Further research is definitely needed to determine the "minimum dose" of treatment in adolescents.

Motivation was also found to foster cognitive improvement in the CACR group, in line with recent findings showing that increased intrinsic motivation leads to better learning in schizophrenia (Choi and Medalia, 2009). CACR efficacy may be further improved by increasing the motivation-enhancing features, such as letting participants choose between a range of exercises, as was the case with CG. It is unclear whether the improvements in attention and delayed memory, symptoms and social-occupational functioning, which were also observed in the CG control group, are caused by CG and CACR, the individual support received from research psychologists, practice effects, or by the general participation in activities at the day clinic, which included psychiatric rehabilitation in both groups. According to two recent meta-analyses (McGurk et al., 2007; Wykes et al., 2011), cognitive remediation combined with active psychiatric rehabilitation achieves the highest gains in psychosocial functioning for adult patients with schizophrenia. Despite its use of this combination, the

present study found no specific effect of CACR on psychosocial functioning, suggesting this meta-analytical result might not hold for adolescents, as is the case for patients with heterogeneous diagnoses, for which lower efficacy has been documented (Wykes et al., 2011). However, this result might also be due to the lack of explicit transfer training in the present implementation, which is done in the NEUROCOM trial assessing the effect on cognitive and everyday functioning of a 16-week program of cognitive training (i.e. attention, executive function, learning and memory) included in a comprehensive psychosocial program on first-episode schizophrenia patients (Wykes et al., 1999). An additional treatment-as-usual control group could help disentangle some of these effects.

Limitations of the current work include a small sample size that may have precluded the detection of the smaller effects typically found for symptom and psychosocial functioning improvement (McGurk et al., 2007; Medalia and Richardson, 2005; Wykes et al., 2011) and the inclusion of adolescents with high risk of psychosis whose cognitive deficits may ameliorate spontaneously. Given the preliminary nature of these encouraging results, further studies on larger samples are needed to confirm the reported improvements, as type I error may have hampered a clear interpretation of the results.

In summary, CACR can be successfully administered to adolescents with psychosis or with high risk, yielding significant improvements in visuospatial abilities, which are important in everyday life situations including navigating familiar and new environments, visual decision making under time pressure (for example when driving a car), and inferring goals of others from analysing facial expressions.

References

- Anderson, C. A.** (2004). An update on the effects of playing violent video games. *Journal of Adolescence*, 27, 113–122.
- APA** (1994). *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (4th edn.). Washington, DC: American Psychiatric Association.
- Bellucci, D. M., Glaberman, K. and Haslam, N.** (2003). Computer-assisted cognitive rehabilitation reduces negative symptoms in the severely mentally ill. *Schizophrenia Research*, 59, 225–232.
- Bremer, J. and Rauch, P. K.** (1998). Children and computers: risks and benefits. *Journal of the American Academy of Child and Adolescent Psychiatry*, 37, 559–560.
- Choi, J. and Medalia, A.** (2009). Intrinsic motivation and learning in a schizophrenia spectrum sample. *Schizophrenia Research*, 118, 12–19.
- Cohen, J.** (1992). A power primer. *Psychological Bulletin*, 112, 155–159.
- Eack, S. M., Greenwald, D. P., Hogarty, S. S., Cooley, S. J., DiBarry, A. L., Montrose, D. M., et al.** (2009). Cognitive enhancement therapy for early-course schizophrenia: effects of a two-year randomized controlled trial. *Psychiatric Services*, 60, 1468–1476.
- Eack, S. M., Hogarty, G. E., Cho, R. Y., Prasad, K. M., Greenwald, D. P., Hogarty, S. S., et al.** (2010). Neuroprotective effects of cognitive enhancement therapy against gray matter loss in early schizophrenia: results from a 2-year randomized controlled trial. *Archives of General Psychiatry*, 67, 674–682.
- Gowers, S. G., Harrington, R. C., Whitton, A., Lelliott, P., Beevor, A., Wing, J., et al.** (1999). Brief scale for measuring the outcomes of emotional and behavioural disorders in children. Health of the Nation Outcome Scales for Children and Adolescents (HoNOSCA). *The British Journal of Psychiatry*, 174, 413–416.

- Grynszpan, O., Perbal, S., Pelissolo, A., Fossati, P., Jouvent, R., Dubal, S., et al. (2011). Efficacy and specificity of computer-assisted cognitive remediation in schizophrenia: a meta-analytical study. *Psychological Medicine*, 41, 163–173.
- Hogarty, G. E., Flesher, S., Ulrich, R., Carter, M., Greenwald, D., Pogue-Geile, M., et al. (2004). Cognitive enhancement therapy for schizophrenia: effects of a 2-year randomized trial on cognition and behavior. *Archives of General Psychiatry*, 61, 866–876.
- Holzer, L., Chinet, L., Jaugey, L., Plancherel, B., Sofia, C., Halfon, O., et al. (2007). Detection of cognitive impairment with the Repeatable Battery for the Assessment of Neuropsychological Status (RBANS) in adolescents with psychotic symptomatology. *Schizophrenia Research*, 95, 48–53.
- Howell, D. C. (2008). The treatment of missing data. In W. Outhwaite and S. Turner (Eds.), *Handbook of Social Science Methodology* (pp. 208–224). London: Sage.
- Kay, S. R., Fiszbein, A. and Opler, L. A. (1987). The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia Bulletin*, 13, 261–276.
- Keefe, R. S., Vinogradov, S., Medalia, A., Silverstein, S. M., Bell, M. D., Dickinson, D., et al. (2011). Report from the working group conference on multisite trial design for cognitive remediation in schizophrenia. *Schizophrenia Bulletin*, 37, 1057–1065.
- Kurtz, M. M., Seltzer, J. C., Shagan, D. S., Thime, W. R. and Wexler, B. E. (2007). Computer-assisted cognitive remediation in schizophrenia: what is the active ingredient? *Schizophrenia Research*, 89, 251–260.
- McGurk, S. R., Twamley, E. W., Sitzler, D. I., McHugo, G. J. and Mueser, K. T. (2007). A meta-analysis of cognitive remediation in schizophrenia. *The American Journal of Psychiatry*, 164, 1791–1802.
- Medalia, A., Aluma, M., Tryon, W. and Merriam, A. E. (1998). Effectiveness of attention training in schizophrenia. *Schizophrenia Bulletin*, 24, 147–152.
- Medalia, A. and Richardson, R. (2005). What predicts a good response to cognitive remediation interventions? *Schizophrenia Bulletin*, 31, 942–953.
- Miller, T. J., McGlashan, T. H., Woods, S. W., Stein, K., Driesen, N., Corcoran, C. M., et al. (1999). Symptom assessment in schizophrenic prodromal states. *The Psychiatric Quarterly*, 70, 273–287.
- Nurnberger, J. I., Blehar, M. C., Kaufmann, C. A., Yorkcooler, C., Simpson, S. G., Harkavyfriedman, J., et al. (1994). Diagnostic Interview for Genetic-Studies: rationale, unique features, and training. *Archives of General Psychiatry*, 51, 849–859.
- Palmer, B. W., Dawes, S. E. and Heaton, R. K. (2009). What do we know about neuropsychological aspects of schizophrenia? *Neuropsychology Review*, 19, 365–384.
- Preisig, M., Fenton, B. T., Matthey, M. L., Berney, A. and Ferrero, F. (1999). Diagnostic Interview for Genetic Studies (DIGS): inter-rater and test-retest reliability of the French version. *European Archives of Psychiatry and Clinical Neuroscience*, 249, 174–179.
- Randolph, C., Tierney, M. C., Mohr, E. and Chase, T. N. (1998). The Repeatable Battery for the Assessment of Neuropsychological Status (RBANS): preliminary clinical validity. *Journal of Clinical and Experimental Neuropsychology*, 20, 310–319.
- Sanford, J. A. and Brown, R. J. (1988). *Captain's Log Cognitive System*. Richmond, VA: Brain Train, Inc.
- Ueland, T. and Rund, B. R. (2004). A controlled randomized treatment study: the effects of a cognitive remediation program on adolescents with early onset psychosis. *Acta Psychiatrica Scandinavica*, 109, 70–74.
- Uhlmann, E. and Swanson, J. (2004). Exposure to violent video games increases automatic aggressiveness. *Journal of Adolescence*, 27, 41–52.
- Urban, S., Pihet, S., Jaugey, L., Halfon, O. and Holzer, L. (2012). Computer-assisted cognitive remediation in adolescents with psychosis or at risk for psychosis: a 6-month follow-up. *Acta Neuropsychiatrica*, 24, 328–335.

- Wykes, T., Huddy, V., Cellard, C., McGurk, S. R. and Czobor, P.** (2011). A meta-analysis of cognitive remediation for schizophrenia: methodology and effect sizes. *The American Journal of Psychiatry*, 168, 472–485.
- Wykes, T., Reeder, C., Corner, J., Williams, C. and Everitt, B. T.** (1999). The effects of neurocognitive remediation on executive processing in patients with schizophrenia. *Schizophrenia Bulletin*, 25, 291–307.