

Effect of average litter weight in pigs on growth performance, carcass characteristics and meat quality of the offspring as depending on birth weight

C. E. Pardo^{1,2}, M. Kreuzer² and G. Bee^{1†}

¹Agroscope Liebefeld-Posieux, Research Station ALP, 1725 Posieux, Switzerland; ²ETH Zurich, Institute of Agricultural Sciences, Universitätstrasse 2, 8092 Zurich, Switzerland

(Received 28 January 2013; Accepted 4 June 2013; First published online 29 July 2013)

Offspring born from normal litter size (10 to 15 piglets) but classified as having lower than average birth weight (average of the sow herd used: 1.46 ± 0.2 kg; mean \pm s.d.) carry at birth negative phenotypic traits normally associated with intrauterine growth restriction, such as brain-sparing and impaired myofiber hyperplasia. The objective of the study was to assess long-term effects of intrauterine crowding by comparing postnatal performance, carcass characteristics and pork quality of offspring born from litters with higher (>1.7 kg) or lower (<1.3 kg) than average litter birth weight. From a population of multiparous Swiss Large White sows (parity 2 to 6), 16 litters with high ($H = 1.75$ kg) or low ($L = 1.26$ kg) average litter birth weight were selected. At farrowing, two female pigs and two castrated pigs were chosen from each litter: from the H-litters those with the intermediate ($H_I = 1.79$ kg) and lowest ($H_L = 1.40$ kg) birth weight, and from L-litters those with the highest ($L_H = 1.49$ kg) and intermediate ($L_I = 1.26$ kg) birth weight. Average birth weight of the selected H_I and L_I piglets differed ($P < 0.05$), whereas birth weight of the H_L - and L_H -piglets were similar ($P > 0.05$). These pigs were fattened in group pen and slaughtered at 165 days of age. Pre-weaning performance of the litters and growth performance, carcass and meat quality traits of the selected pigs were assessed. Number of stillborn and pig mortality were greater ($P < 0.05$) in L- than in H-litters. Consequently, fewer ($P < 0.05$) piglets were weaned and average litter weaning weight decreased by 38% ($P < 0.05$). The selected pigs of the L-litters displayed catch-up growth during the starter and grower–finisher periods, leading to similar ($P > 0.05$) slaughter weight at 165 days of age. However, H_L -gilts were more feed efficient and had leaner carcasses than H_I , L_H - and L_I -pigs (birth weight class \times gender interaction $P < 0.05$). Meat quality traits were mostly similar between groups. The marked between-litter birth weight variation observed in normal size litters had therefore no evident negative impact on growth potential and quality of pigs from the lower birth weight group.

Keywords: litter birth weight, litter size, postnatal growth, carcass characteristics, pig

Implications

In pig litters of normal size (10 to 15 pigs), large differences in average litter weight are generally observed. The question arises whether offspring born from normal litter size but classified as having lower than average litter birth weight suffered an intrauterine insult. The observed greater number of stillborns and higher pre-weaning mortality is indicative of impaired prenatal development of these littermates. Despite post-weaning catch-up growth found in offspring born from low birth weight litters, increasing average litter weight should be targeted to enhance sow reproduction performance and offspring viability.

Introduction

In the last 10 years, sow prolificacy markedly increased in many pig breeds as, for instance, the Swiss Large White populations (Suisag, 2012). With a litter size ranging from 5 to 20 piglets born, a distinct linear decrease in average litter birth weight and a concomitant increase in the proportion of low birth weight pigs can be observed (Foxcroft *et al.*, 2007; Fix *et al.*, 2010). Although average litter birth weight is still moderately inverse to litter size when only litters with 10 to 15 newborns are included in the analysis (Foxcroft *et al.*, 2007), the between-litter variance in average litter birth weight can still reach a maximum of 1 kg. These observations led to the assumption that different patterns of prenatal losses in gestating sows may cause differences in average litter birth weight. On the basis of the compelling

† E-mail: giuseppe.bee@agroscope.admin.ch

evidence delivered by van der Waaij *et al.* (2010), greater mortality rates between implantation on day 13 and up to day 40 of pregnancy had detrimental effects on placenta weight, thus impairing development of the fetuses.

Impaired placental efficiency is predicted to result in prenatal programming effects not only of individual pigs but, to some extent, of all offspring in the litter (Wu *et al.*, 2006; Foxcroft *et al.*, 2007). Because the early stages of gestation mark the onset of myogenesis (Zhao *et al.*, 2011), it is reasonable to assume that differences in placental efficiency caused by different prenatal survival patterns will affect myogenesis and ultimately postnatal growth efficiency (Dwyer *et al.*, 1993).

On the basis of these assumptions, the aim of the study was to compare litter characteristics, as well as postnatal growth performance, carcass characteristics and meat quality traits of offspring originating from litters with 10 to 15 pigs born with either high- or low-average litter birth weight. Of special interest was the question of how the assumed different pattern of prenatal losses could affect postnatal development of the progeny. Therefore, pigs of both genders that (1) either fell within the mid-birth weight range of their respective litters in the two average birth weight groups or (2) had a similar birth weight but originated from different average litter birth weight groups were investigated.

Material and methods

Animals and treatments

All procedures involving animals were approved by the Swiss Cantonal Committee for Animal Care and Use. From the herd of the Agroscope Liebefeld Posieux research station, a total of 16 multiparous Swiss Large White sows (parity 2 to 6) were selected. Two criteria were considered for sow selection: (1) the number of piglets born per litter had to be in the range of 10 to 15 piglets and (2) the average litter birth weight had to be either >1.7 kg (H) or <1.3 kg (L). Allocation to these two treatment groups was performed on the day of farrowing considering the number of total piglets born and the average litter birth weights. Sows were artificially inseminated three times on 2 consecutive days with unfrozen semen from 10 Swiss Large White boars (Suisag, Sempach, Switzerland). During the gestation and lactation periods, all sows were fed standard diets formulated according to Swiss nutrient recommendations for pigs (Agroscope Liebefeld Posieux Research station (ALP), 2012). Within 12 h after birth, the number of piglets born alive, the number of stillborn and their individual BW were determined. As requested by the Swiss animal welfare law, male pigs were castrated only after isoflurane anesthesia induction within 4 days after birth. All piglets of the H- and L-sows stayed with the dam until weaning at day 27 (± 1.2 days), and no cross-fostering was permitted. Suckling piglets had access to creep feed from day 10 until the end of lactation. At weaning, where available, two castrates and two female pigs were chosen from each litter for assessing growth performance during the starter and grower–finisher periods and

Table 1 Ingredients and nutrient composition of the experimental diets

Item	Starter diet	Grower diet	Finisher diet
Ingredient composition (g/kg)			
Barley	287.4	68.0	100.0
Corn	60.0	140.0	140.0
Wheat	474.0	436.0	494.0
Dried sugar beet pulp	16.0	100.0	100.0
Soybean meal	4.0	236.0	146.0
Potato protein	85.0	–	–
Rapeseed	24.0	–	–
NaCl	4.3	3.0	3.2
Dicalcium phosphate	15.5	0.9	2.2
Calcium carbonate	–	7.3	7.0
Calcium formate	14.2	–	–
L-lysine HCl	5.3	1.4	1.5
D,L-methionine	1.1	0.2	–
L-threonine	1.5	0.3	–
L-tryptophan	0.7	–	–
Vitamin–mineral premix ¹	4.0	4.0	4.0
Pellan ²	3.0	3.0	3.0
Analyzed and calculated nutrient and energy content, expressed per kg dry matter			
Total ash (g)	53	50	45
Crude fiber (g)	28	50	47
CP (g)	173	209	180
Ether extract (g)	37	27	25
N-free extract (g ³)	708	664	703
Digestible energy (MJ)	16.1	15.6	15.6

¹Supplied the following micronutrients per kilogram of diet: 20 000 IU of vitamin A, 200 IU of vitamin D₃, 39 mg of vitamin E, 2.9 mg of riboflavin, 2.4 mg of vitamin B₆, 0.01 mg of vitamin B₁₂, 0.2 mg of vitamin K₃, 10 mg of pantothenic acid, 1.4 mg of niacin, 0.48 mg of folic acid, 199 g of choline, 0.052 mg of biotin, 52 mg of Fe as FeSO₄, 0.16 mg of I as Ca(IO₃)₂, 0.15 mg of Se as Na₂Se, 5.5 mg of Cu as CuSO₄, 81 mg of Zn as ZnO₂, and 15 mg of Mn as MnO₂.

²Pellet binder (Mikro-Technik, GmbH & Co. KG, Bürgstadt, Germany).

³Calculated as dry matter – total ash – crude fiber – CP – ether extract.

carcass characteristics and meat quality traits at slaughter. From the H-sows, those with the intermediate (H_i; $n = 16$) and those with the lowest (H_L; $n = 14$) birth weight were selected. Selection from the L-sows included those with the intermediate (L_i; $n = 16$) and those with the highest (L_H; $n = 15$) birth weight, similar to that of the H_L-piglets. Runt piglets with a birth weight of <0.8 kg were not considered for selection. Subsequently, BW and average daily gain (ADG) of all piglets were determined during lactation. From weaning until slaughter, the pigs selected were reared in group pens (eight pigs/pen) equipped with automatic feeder and individual pig recognition system (Schauer Maschinenfabrik GmbH. & Co KG, Prambachkirchen, Austria) as described previously (Bee *et al.*, 2004). Whenever possible, pigs of the same litter stayed in the same pen. Individual *ad libitum* intake of the starter (weaning to 28 kg BW), grower (28.1 to 63.5 kg BW) and finisher diets (63.6 kg BW until slaughter) was monitored and recorded by the control software (MLP, Schauer) linked to each feeder. The composition of these three diets is specified in Table 1. Pigs were weighed weekly. Growth performance data in the table

are presented for the preweaning, starter and grower–finisher (28.1 kg to slaughter) periods.

Slaughter procedure and carcass measurements

Pigs were slaughtered at 165 ± 1.5 days of age in the research station's abattoir. The last 15 h before slaughter, feed was withheld from the pigs. The animals were stunned in a CO₂ stunner (87% CO₂) for 100 s, immediately exsanguinated, scalded, mechanically de-haired, eviscerated and split along the mid-line within 30 min. Weights of hot carcass (total and left side), kidneys, spleen, liver, lung, heart and brain were assessed. Carcasses were then chilled at 2°C for 24 h. One day before post mortem, the total and left carcass side were weighed again to determine cold carcass weight and dissected into major primal cuts (loin, ham, shoulder and belly) and total lean meat was calculated as described previously (Bee *et al.*, 2002). Back fat thickness was determined at the 10th rib level.

Meat quality measurements

Temperature and pH were measured 30 min and 24 h post mortem in the *m. longissimus dorsi* (LM; at the 10th rib level) and the dark portion of the *m. semitendinosus* (ST) using a pH meter (WTW pH 197-S; WTW, Weilheim, Germany) equipped with an insertion glass spearhead electrode (Metrohm, 6.0226.100) combined with a temperature probe (Pt 1000, Metrohm AG, Herisau, Switzerland). After the second assessment of pH and temperature, the entire ST was excised and weighed. Samples of the LM, harvested between the 10th and the 13th rib level, were cut into four 1.5 cm-thick slices free of subcutaneous adipose tissue and bone and labeled A, B, C and D. From the dark portion of the ST, two (E and F) longitudinal slices, weighing ~70 g each, were prepared. The A, C and E slices were used to assess drip loss (48 h) using the plastic bag method by Honikel (1998). After a 10 min bloom, the L* (lightness) values were determined on the B, D and F slices, using a spectrophotometer (model CM-2600d, Minolta, Dietikon, Switzerland). Three measurements on each sample were carried out and averaged. Afterwards, the B and D slices were weighed, vacuum-packed and stored at -20°C. Later the frozen samples were thawed for 24 h at 2°C and then weighed to determine thaw loss. After 1 h at room temperature, the slices were cooked on a preheated (190°C to 195°C) grill plate (Beer Grill AG, Zurich, Switzerland) to an internal temperature of 69°C recorded by a thermometer probe. The slices were cooked for 1.5 min on one side, then turned and cooked for 3.5 min, and finally 2 min on the initial side. Slices were reweighed to determine cooking loss. Approximately 2 h after cooking when meat samples reached room temperature, shear force was determined (five cores per sample) using a Stable Micro System TA.XT2 Texture Analyzer (Godalming, Surrey, UK) equipped with a 2.5-mm-thick Warner–Bratzler shear blade. Shear force values were expressed as the average of the five measurements.

Statistical analysis

Data were analyzed using the mixed procedure (SAS Inst. Inc., Cary, NC, USA). The statistical model used for analyzing

the data from the reproductive performance of the sows and litter characteristics included the average weight of the litter (H- v. L-sows) as fixed effect and the sire nested within sows as random effect. Growth performance, carcass characteristics and meat quality traits of the pigs selected were analyzed with a model including birth weight group (H_i, H_L, L_H, L_i), gender and the two-way interactions. Sire nested within sow was included as a random effect. The pig was the experimental unit. Comparisons among least square means were made with the PDIFF statement using the Tukey–Kramer adjustment. Means were considered statistically different at $P < 0.05$ and referred to as tendencies at $P < 0.10$. The results are presented in the tables as least square means together with the pooled standard error.

Results

Reproductive performance of the sows

In both groups, the sows selected were on average in their third parity (Table 2). Litter size and number of piglets born alive were similar in the two groups, although the number of stillborns was greater ($P = 0.01$) in L- than in H-litters. Piglet survival rate during lactation was lower ($P < 0.01$) in L- compared with H-litters and, consequently, fewer ($P = 0.02$) piglets were weaned from L-litters. Despite a similar female-to-male ratio at birth in both experimental groups, L-sows weaned fewer ($P < 0.07$) female pigs (data not shown). As planned, the average litter birth weight of total piglets born

Table 2 Reproductive performance of sows classified as having high (H > 1.7 kg)- or low (L < 1.3 kg)-average litter birth weight and litter sizes of between 10 and 15 total pigs born

Trait	Litter birth weight		s.e.m.	P-value
	H	L		
Number of sows	8	8		
Average parity, n	3.1	3.5	0.5	0.61
Number of piglets				
Total born	13.4	14.1	0.4	0.22
Born alive	13.1	12.2	0.5	0.25
Stillborn	0.3	1.9	0.4	0.01
Weaned	12.0	8.5	0.9	0.02
Survival rate ¹	0.89	0.61	0.06	<0.01
Total litter weight				
At birth (kg) ²	23.0	15.3	0.7	<0.01
At weaning (kg)	103.9	64.3	8.6	<0.01
Average body weight				
At birth (kg) ²	1.75	1.26	0.04	<0.01
At weaning (kg)	8.60	7.71	0.43	0.17
Standard deviation				
Of birth weight (kg) ²	0.29	0.22	0.02	0.06
Of weaning weight (kg)	1.42	1.00	0.32	0.17
Average daily gain				
Per piglet (kg)	0.24	0.24	0.01	0.84
Of the litter (kg)	2.89	1.84	0.21	0.02

¹Calculated from birth to weaning.

²Calculations included piglets born alive only.

Table 3 Effect of birth weight class and gender on growth performance during lactation, starter and grower–finishing periods of the selected piglets¹

Trait	Birth weight class				Gender			P-value		
	H _I	H _L	L _H	L _I	Barrow	Gilts	s.e.m.	B	G	B × G
Body weight (kg)										
At birth	1.79 ^c	1.40 ^b	1.49 ^b	1.26 ^a	1.50	1.47	0.04	<0.01	0.31	0.30
At weaning	9.70 ^b	7.94 ^a	8.06 ^{ab}	7.37 ^a	8.08	8.46	0.73	<0.01	0.22	0.55
At start of grower period	27.7	27.4	28.3	27.9	27.9	27.7	0.9	0.81	0.65	0.36
At slaughter	103.9	100.3	101.7	99.2	102.9	99.6	3.8	0.44	0.10	0.09
Pre-weaning period										
Daily gain (g/day)	281 ^b	228 ^a	248 ^{ab}	230 ^a	240	254	21	0.01	0.22	0.53
Starter period										
Daily gain (g/day)	377	366	396	379	381	377	20	0.39	0.67	0.80
Daily feed intake (g/day)	644	627	658	636	642	641	28	0.51	0.98	0.96
Total feed intake (kg)	30.8	33.4	33.8	34.6	33.5	32.8	1.7	0.10	0.51	0.19
Gain to feed (g/kg)	585	584	601	595	595	588	17	0.78	0.49	0.77
Duration (days)	48.4	53.6	51.4	54.7	52.6	51.4	3.1	0.05	0.42	0.06
Grower–finisher period										
Daily gain (g/day)	866	879	840	847	882	834	31	0.64	0.01	0.34
Daily feed intake (g/day)	2307	2301	2333	2348	2435	2209	81	0.90	<0.01	0.07
Total feed intake (kg)	203.1	190.5	203.3	196.9	206.7	190.2	10.6	0.47	0.01	0.03
Gain to feed (g/kg)	376 ^b	383 ^b	360 ^a	361 ^a	361	378	7	<0.01	<0.01	0.05
Duration (days)	88.1	82.7	87.2	83.8	84.9	86.1	2.9	0.05	0.43	0.07

¹H_I and H_L = littermates with the intermediate ($n = 16$) and lowest ($n = 14$) birth weight within their litter, born from sows giving birth to litters classified as having high (>1.7 kg)-average litter birth weight; L_H and L_I = littermates with the greatest ($n = 15$) and intermediate ($n = 16$) birth weight within their litter born from sows giving birth to litters classified as having low (<1.3 kg)-average litter birth weight; $n = 31$ for barrow; $n = 30$ for gilts; B = birth weight class, G = gender.

^{a,b,c}Least square means within a row for the main factor birth weight class without a common superscript differ ($P < 0.05$).

alive clearly differed ($P < 0.01$), with the H-litters being 490 g heavier than the L-litters. Concomitantly, total litter weight at birth was lower ($P < 0.01$) for the L-litters than the H-litters. Compared with the H-litters, the L-litters birth weight of female pigs, but not male pigs, were less ($P < 0.01$) variable (data not shown). This resulted in an overall lower standard deviation in L-litters. Regardless of gender, ADG expressed per piglet was similar in both litter groups during lactation, but the high variation prevented the average weaning BW to be significantly different between groups. When calculated for the whole litter, total weaning weight and ADG of the litter were markedly lower ($P \leq 0.02$) in L- than in H-litters.

Birth weight and growth performance of the selected offspring

At birth, H_L- and L_H-piglets had similar average BW, whereas H_I-piglets were 530 g heavier ($P < 0.05$) than L_I-piglets (Table 3). Considering birth weight class within litter groups, H_L-piglets were 390 g lighter ($P < 0.05$) than in H_I-piglets, and L_I-piglets were 230 g lighter ($P < 0.05$) than in L_H-piglets. At weaning, the H_I-piglets were still the heaviest group ($P < 0.05$ against H_L and L_I). In addition, the difference in BW was greater within the H- than within the L-litter piglet groups (H_I v. H_L = 1.76 kg and L_H v. L_I = 0.69 kg). This effect was related to the faster ($P < 0.05$) growth of H_I- compared with H_L-piglets during lactation. During the starter and grower–finisher periods, average daily feed intake (ADFI) and ADG were similar for all groups. Conversely, when comparing

pigs born with similar birth weight but originating either from H- or L-litters (H_L v. L_H), the H_L-pigs were more ($P < 0.05$) feed efficient in the grower–finisher period. At 165 days of age, the slaughter BW did not differ among groups. However, the time needed to reach the target BW in the grower–finisher period was shorter ($P < 0.05$) in H_L- than in H_I-pigs (with the L-litter pigs remaining intermediate) and appeared to be a compensation for the corresponding numerical longer starter period of the H_L-pigs.

Gender differences were not observed in the starter period for any of the variables measured. Regardless of the birth weight class, gilts tended to have a lower ($P = 0.10$) slaughter weight at 165 days than barrows as a result of the lower ($P \leq 0.01$) ADG and ADFI in the grower–finisher period. Nevertheless, gilts were more feed efficient and total feed intake was proportionately lower in the same period than that of the barrows ($P \leq 0.01$ for each). Birth weight class × gender interactions ($P \leq 0.05$) were observed for the gain-to-feed ratio and total feed intake in the grower–finisher period. The H_L-female pigs were the most feed efficient of all the other pig categories (Figure 1). In addition, compared with H_L-barrows, H_L-gilts consumed markedly less ($P < 0.05$) of the grower and finisher diets.

Carcass characteristics and morphometric data of the fattened offspring

Birth weight had little impact on the carcass characteristics (Table 4). Belly percentage was lower ($P < 0.05$) in H_L-pigs




Figure 1 Effect of birth weight and gender on total feed intake and gain to feed ratio during the grower–finisher period. H₁ and H₂ = littermates with the intermediate and lowest birth weight within their litter, born from sows giving birth to litters classified as having high (>1.7 kg) average litter birth weight; L₁ and L₂ = littermates with the greatest and intermediate birth weight within their litter born from sows giving birth to litters classified as having low (<1.3 kg) average litter birth weight. ^{a,b,c}Least square means without a common superscript differ ($P < 0.05$).

Table 4 Effect of birth weight class and gender on carcass characteristics¹

Trait	Birth weight class				Gender			P-value		
	H ₁	H ₂	L ₁	L ₂	Barrows	Gilts	s.e.m.	B	G	B × G
Hot carcass weight (kg)	83.9	80.7	82.4	80.3	83.1	80.6	3.2	0.48	0.16	0.06
Carcass length (cm)	96.2	94.9	95.4	95.3	95.1	95.8	1.4	0.65	0.32	0.22
Carcass yield (%)	80.7	80.4	81.0	81.0	80.7	80.9	0.6	0.69	0.48	0.05
Lean meat (%)	56.2	55.6	54.7	54.8	54.1	56.5	0.7	0.25	<0.01	0.01
Loin (%)	25.2	24.8	24.4	24.4	24.2	25.2	0.4	0.16	<0.01	0.01
Ham (%)	18.8	18.6	18.3	18.3	18.1	18.9	0.3	0.50	<0.01	0.04
Shoulder (%)	12.2	12.2	12.0	12.0	11.9	12.3	0.2	0.84	<0.01	0.41
Belly (%)	18.6 ^a	18.6 ^a	19.3 ^b	19.1 ^{ab}	18.8	18.9	0.3	0.05	0.46	0.28
Total subcutaneous fat (%)	12.6	12.8	13.2	13.1	14.0	11.9	0.5	0.65	<0.01	0.03
10th rib fat thickness (mm)	19.8	20.0	20.3	20.1	21.8	18.3	1.2	0.98	<0.01	0.16
Omental fat (%)	1.27	1.30	1.45	1.35	1.48	1.20	0.99	0.33	<0.01	0.27

¹H₁ and H₂ = littermates with the intermediate ($n = 16$) and lowest ($n = 14$) birth weight within their litter, born from sows giving birth to litters classified as having high (>1.7 kg)-average litter birth weight; L₁ and L₂ = littermates with the greatest ($n = 15$) and intermediate ($n = 16$) birth weight within their litter, born from sows giving birth to litters classified as having low (<1.3 kg)-average litter birth weight; $n = 31$ for barrows; $n = 30$ for gilts; B = birth weight class, G = gender. ^{a,b}Least square means within a row for the main factor birth weight class without a common superscript differ ($P < 0.05$).

and H₁-pigs compared with L₁-pigs, with intermediate values for L₂-pigs. As a result of the greater ($P < 0.01$) percentages of loin, ham and shoulder, and the lower ($P < 0.01$) percentages of omental and back fat, carcasses of gilts were leaner ($P < 0.01$) compared with those of barrows. However, the gender difference for lean meat, loin, ham and total subcutaneous fat percentages was mainly evident between female pigs and barrows born from H-litters, whereas L₁- and L₂-female pigs had similar percentages of loin, ham and subcutaneous fat than barrows (birth weight class × gender interaction; $P \leq 0.03$). Relative weight of internal organs and brain expressed as a percentage of hot carcass weight did not differ among birth weight classes and gender (Table 5). By contrast, the relative ST weight tended ($P < 0.10$) to be lower in L₁- than in H₂-pigs and was lower ($P < 0.01$) in barrows than gilts.

Quality of the meat from the fattened offspring

Birth weight class effects on the assessed meat quality traits were observed only for LM temperature at 24 h post mortem where temperature was lower ($P < 0.05$) in L₂- than in H₁-pigs,

with intermediate values in L₁-pigs and H₂-pigs (Table 6). With respect to gender, pH and temperature at 30 min post mortem of the ST was greater ($P \leq 0.05$) in barrows than in gilts, whereas this was not the case in the LM. At 24 h post mortem, the pH and temperature of both muscles did not differ according to gender. Drip and thaw losses of the LM of barrows were lower ($P \leq 0.01$) than in the LM of the gilts.

Discussion

Sow performance in relation to average litter birth weight
A greater incidence of low birth weight piglets has been associated with increased litter size, the latter being the consequence of intensive breeding efforts in the last few decades with the objective to enhance the prolificacy of dams and ultimately improve reproduction efficiency (Milligan *et al.*, 2002; Quiniou *et al.*, 2002; Quesnel *et al.*, 2008).

In the current study, the difference in average litter birth weight between L- and H-litters was ~500 g. In contrast to other studies (e.g., Bérard *et al.*, 2008), this magnitude of

Table 5 Effect of birth weight class and gender on the relative weight of the internal organs, brain and semitendinosus muscle¹

Trait	Birth weight class				Gender			P-value		
	H _I	H _L	L _H	L _I	Barrows	Gilts	s.e.m.	B	G	B × G
Heart (%)	0.49	0.48	0.47	0.54	0.48	0.38	0.02	0.18	0.27	0.88
Liver (%)	2.12	2.18	2.06	2.15	2.13	2.13	0.07	0.46	0.99	0.33
Kidneys (%)	0.40	0.41	0.41	0.44	0.41	0.42	0.02	0.13	0.20	0.40
Spleen (%)	0.16	0.19	0.17	0.17	0.17	0.17	0.01	0.11	0.72	0.80
Lung (%)	0.72	0.74	0.79	0.75	0.72	0.78	0.03	0.34	0.04	0.34
Brain (%)	0.13	0.13	0.12	0.13	0.13	0.13	<0.01	0.55	0.40	0.11
Semitendinosus (%)	0.55 ^{xy}	0.54 ^y	0.50 ^x	0.51 ^{xy}	0.51	0.54	0.02	0.07	<0.01	0.37
Brain : semitendinosus weight ratio	0.24	0.24	0.25	0.26	0.25	0.24	0.02	0.34	0.27	0.60

¹Relative weights of the organs and semitendinosus muscle are expressed as percentage of hot carcass weight. H_I and H_L = littermates with the intermediate ($n = 16$) and lowest ($n = 14$) birth weight within their litter, born from sows giving birth to litters classified as having high (>1.7 kg)-average litter birth weight; L_H and L_I = littermates with the greatest ($n = 15$) and intermediate ($n = 16$) birth weight within their litter born from sows giving birth to litters classified as having low (<1.3 kg)-average litter birth weight; $n = 31$ for barrow; $n = 30$ for gilts; B = birth weight class, G = gender.

^{x,y,a,b}Least square means within a row for the main factor birth weight class without a common superscript differ ($P < 0.10$).

Table 6 Effect of birth weight class and gender on meat quality traits in slaughter pigs¹

Trait	Birth weight class				Gender			P-value		
	H _I	H _L	L _H	L _I	Barrows	Gilts	s.e.m.	B	G	B × G
Longissimus muscle (pH)										
30 min	6.61	6.55	6.56	6.57	6.58	6.56	0.04	0.34	0.31	0.57
24 h	5.61	5.62	5.62	5.62	5.62	5.61	0.03	0.99	0.57	0.44
Temperature (°C)										
30 min	39.0	39.3	38.9	38.8	39.0	39.0	0.2	0.26	0.51	0.17
24 h	2.0 ^b	1.8 ^{ab}	1.6 ^{ab}	1.5 ^a	1.8	1.7	0.1	0.03	0.19	0.86
Color										
L*	46.3	47.5	45.5	45.5	46.5	45.9	1.0	0.12	0.32	0.27
Drip loss [% (48 h)]	2.6	2.6	2.3	2.6	2.3	2.8	0.4	0.76	0.01	0.01
Thaw loss (%)	11.5	11.3	11.7	11.5	11.0	12.1	0.6	0.89	<0.01	0.16
Cooking loss (%)	19.6	20.4	19.9	19.9	19.8	20.0	0.7	0.36	0.56	0.49
Shear force (kg)	63.7	61.4	66.9	68.6	63.5	66.7	5.5	0.58	0.34	0.14
Semitendinosus muscle dark portion (pH)										
30 min	6.56	6.52	6.56	6.56	6.58	6.52	0.05	0.78	0.05	0.76
24 h	5.54	5.57	5.62	5.60	5.57	5.59	0.04	0.36	0.59	0.97
Temperature (°C)										
30 min	39.9	40.0	39.9	39.9	39.8	40.0	0.1	0.67	0.03	0.42
24 h	2.5	2.4	3.9	2.4	2.5	3.1	1.1	0.33	0.40	0.20
Color										
L*	40.6	40.8	39.5	40.0	40.4	40.0	0.8	0.45	0.27	0.89
Drip loss (48 h) (%)	1.7	1.8	1.5	1.9	1.8	1.7	0.3	0.29	0.27	0.36

¹H_I and H_L = littermates with the intermediate ($n = 16$) and lowest ($n = 14$) birth weight within their litter, born from sows giving birth to litters classified as having high (>1.7 kg)-average litter birth weight; L_H and L_I = littermates with the greatest ($n = 15$) and intermediate ($n = 16$) birth weight within their litter born from sows giving birth to litters classified as having low (<1.3 kg)-average litter birth weight; $n = 31$ for barrow; $n = 30$ for gilts; B = birth weight class, G = gender.

^{a,b}Least square means within a row for the main factor birth weight class without a common superscript differ ($P < 0.05$).

difference in birth weight was unrelated to the number of total born. Nevertheless, in the present study, the standard deviation of within-litter birth weight tended to be lower in L- than in H-litters. Foxcroft *et al.* (2007) explained this phenomenon as a consequence of prenatal losses of weaker and smaller embryos and fetuses, thereby reducing intra-litter variability in birth weight. Another striking difference between the two litter birth weight groups was the greater proportion of stillborn in L- (13%) compared with H-litters

(2%), which, except for the fact that in the present case it was unrelated to litter size, matches results of earlier studies (Leenhouders *et al.*, 1999; Milligan *et al.*, 2002; Quiniou *et al.*, 2002). As observed by Leenhouders *et al.* (1999), a low average birth weight is associated with a greater pre-weaning mortality, which was more than three times greater in L- than in H-litters (39% v. 11%, respectively). The 33% difference between L- and H-litters in total litter weight at birth increased to 38% at weaning. In accordance with

Bérard and Bee (2010), ADG during lactation was independent of birth weight. Our findings are in contrast to results reported by Quiniou *et al.* (2002) showing that during lactation heavier piglets grow faster than lighter piglets. These authors assumed that heavier piglets have a greater ability to occupy the best-performing teats, to stimulate and to drain them, thereby inducing a larger milk flow. However, because of the greater number of stillborn and greater mortality rate in L- than in H-litters, and as no cross-fostering was performed in the current study, competition among littermates for the teats was markedly reduced after the first week of lactation. Thus, growth of L-offspring was not hindered by lacking nutrient supply.

Growth performance as affected by birth weight class and gender

The pigs selected at weaning to evaluate growth performance, carcass and meat quality traits mirrored the selection criteria of the experimental design, as birth weights of L_I- and H_I-pigs were close to the average birth weight of the respective litters and L_H- and H_L-pigs had a similar birth weight. Therefore, based on the results of previous studies reporting a close relationship between birth weight and early postnatal growth (Gondret *et al.*, 2006; Bérard *et al.*, 2010), the ranking of BW at weaning could have been expected to remain similar for the four groups as at birth ($H_I > L_H = H_L > L_I$). In fact, this relationship was confirmed in the H-litters where growth rate and weaning weight depended on birth weight. By contrast in the L-litters, L_I-pigs reached comparable weaning weights as their heavier littermates (L_H). A likely explanation for the similar growth rate might be found in the greater mortality rate in L-litters that entailed less competition for teats and therefore nutrient supply from sow milk, and thus might not have limited growth of lighter and usually less vital pigs. Along the same line of explanation, it was unexpected to observe that L_H-pigs did not translate the greater nutrient availability into greater growth rates, as their ADG was comparable to the one of H_L-pigs.

Consistent with results of previous studies (Gondret *et al.*, 2005; Bérard *et al.*, 2008; Bérard *et al.*, 2010), pigs born with a lower birth weight displayed catch-up growth, because differences in BW between the four experimental groups gradually disappeared after weaning. To compensate for the 2.33 kg lower weaning weight, it took the lightest pigs (L_I) on average 6.3 days longer than the heaviest pigs (H_I) to reach the beginning of the grower period, which was defined to start at an average BW of 27 kg. This helps explaining the greater total intake of starter diet in L_I- compared with H_I-pigs, as neither ADG nor ADFI gains to feed ratio differed between the birth weight groups. In the grower–finisher period, feed efficiency was lower in L- than in H-offspring owing to the numerically lower ADG and greater ADFI. Consistent with the study by Bérard *et al.* (2010), this effect was more distinct in female pigs, implying that under *ad libitum* feed supply, especially female pigs, born from low average litter birth weight sows, will perform less efficiently.

Except for the relative ST weights, which tended to be lower in L_H- than in H_L-pigs and were lower in barrows than gilts, neither birth weight nor gender affected relative weights of internal organs and brain. This was surprising in view of the fact that using a similar experimental design, Pardo *et al.* (2010) reported a lower relative brain weight (expressed relative to the birth weight) in H_I- than in L_I-newborns and lower in H_L- than in L_H-newborns. This had let the authors conclude that those lighter piglets had some characteristics associated with growth retardation as supported by other studies (Bauer *et al.*, 2003; Town *et al.*, 2005; Vallet and Freking, 2006). Furthermore, it was shown that an impaired development of organs such as the kidneys, heart and liver can still be observed in low birth weight compared with high birth weight pigs when they reach the slaughter weight (Bérard *et al.*, 2008; Rehfeldt *et al.*, 2008; Nissen and Oksbjerg, 2011). Therefore, based on the findings on organ weight and carcass composition, it is difficult to assess whether, or to which extent, offspring from L-sows had been subjected to intrauterine growth retardation.

Carcass and meat quality as affected by birth weight class and gender

The data published on the impact of birth weight on carcass characteristics are inconsistent owing to differences in applied feeding strategies, rearing conditions or range of birth weight used in the various studies. For instance, carcasses of low birth weight compared with high birth weight pigs were fatter at slaughter if pigs were reared in individual pens and had *ad libitum* access to the diet (Gondret *et al.*, 2006). By contrast, carcass composition was similar when low birth weight and high birth weight pigs penned in groups had restricted access to the feed (Gondret *et al.*, 2005). Covering a wide birth weight range of up to 850 g, data of various studies reported no relationship between birth weight and lean meat percentage (Gondret *et al.*, 2005; Bérard *et al.*, 2008; Beaulieu *et al.*, 2010). By contrast, when differences in birth weight were ~900 g, Rehfeldt and Kuhn (2006) found a tendency toward leaner carcasses in low birth weight (0.94 kg) compared with high (1.80 kg) birth weight pigs slaughtered at the same age. However, in a similar study by the same authors, carcass leanness did not differ when birth weight differences were <400 g (1.39 v. 1.80 kg or 1.37 v. 1.67 kg) (Rehfeldt and Kuhn, 2006; Rehfeldt *et al.*, 2008). Thus, it appears that the ratio of adipose and muscle tissue deposition rate is affected only in littermates with very low birth weight, which might explain the lack of effect on carcass characteristics in the current experiment with a difference in average birth weight of up to 530 g (L_I- v. H_I-pigs). Furthermore, it has to be kept in mind that, in contrast to all of the aforementioned studies where pigs with extremely high or low birth weight were compared, L_I-pigs were in the intermediate birth weight range within their litters and were 320 and 180 g heavier than the piglets in the studies by Rehfeldt and Kuhn (2006) and Rehfeldt *et al.* (2008), respectively. Nevertheless, there was one exception. Belly percentage, which is a cut with a great amount of intermuscular fat, was greater in L_H- compared with H_L-pigs and H_I-pigs,

suggesting that the impaired feed efficiency of the L_H-pigs in the grower–finisher period affected this pork cut.

Usually when female pigs and barrows have *ad libitum* access to the grower–finisher diets, carcasses of female pigs are leaner at slaughter (Ellis *et al.*, 1996; Rehfeldt *et al.*, 2008; Bérard *et al.*, 2010). This difference was noticeable between gilts and barrows born from the H-litters, whereas percentages of lean meat, loin, ham, subcutaneous and back fat were similar in gilts and barrows of the L_H-group and L_I-group. In addition, these findings match the similar feed efficiency found in the grower–finisher period in the L-litters. Results of other studies also suggested that low birth weight has stronger effects on carcass composition in female pigs compared with male pigs (Poore and Fowden, 2004; Gondret *et al.*, 2006; Bérard *et al.*, 2010).

Effects of birth weight or gender on technological traits of the LM and dark portion of the ST were marginal, which matches results of previous studies (Nissen *et al.*, 2004; Bérard *et al.*, 2008; Rehfeldt *et al.*, 2008; Beaulieu *et al.*, 2010).

In conclusion, this study confirms the occurrence of negative effects on viability and pre-weaning performance of piglets born from litters with a low average weight probably as a result of low gestation efficiency. This gets obvious by the increased number of stillborns and a greater pre-weaning mortality of progeny. Therefore, the number of piglets weaned was reduced, which adversely affected sow performance. Contrary to expectations, low birth weight did not impair postnatal growth of their individuals and, thus fattening performance and the initial variability in weight observed at birth in these litters of normal size gradually disappeared after weaning. There was also no differences in the brain:muscle ratio at slaughter making assumptions about the presence or absence of intrauterine crowding as a reason for low average birth weight inconclusive. Within the litters, gender variability caused more effects on postnatal growth in the L-litters, particularly in the female pigs, than in the H-litters. Further studies have to identify the reasons for the differences in the efficiency in sows with apparently normal litter size and the gender-dependence adverse effects on performance of the offspring of low birth weight litters.

Acknowledgments

The authors thank Guy Maikoff and his team for taking care of the animals, and Dr Paolo Silacci and his team for the help during sample collection and muscle analysis. This study was supported by the Fondation Sur-la-Croix (Basel, Switzerland).

References

Agroscope Liebefeld Posieux Research Station (ALP) 2012. Fütterungsempfehlungen und Nährwerttabellen für Schweine [Feeding recommendations and nutrient tables for pigs]. Agroscope, Posieux, Switzerland.

Bauer R, Walter B, Brust P, Fuchtnner F and Zwiener U 2003. Impact of asymmetric intrauterine growth restriction on organ function in newborn piglets. *European Journal of Obstetrics Gynecology and Reproductive Biology* 110, S40–S49.

Beaulieu AD, Aalhus JL, Williams NH and Patience JF 2010. Impact of piglet birth weight, birth order, and litter size on subsequent growth performance, carcass

quality, muscle composition, and eating quality of pork. *Journal of Animal Science* 88, 2767–2778.

Bee G, Gebert S and Messikommer R 2002. Effect of dietary energy supply and fat source on the fatty acid pattern of adipose and lean tissues and lipogenesis in the pig. *Journal of Animal Science* 80, 1564–1574.

Bee G, Jacot S, Guex G and Herzog W 2004. Effects of two supplementation levels of linseed combined with CLA or tallow on meat quality traits and fatty acid profile of adipose tissue and longissimus muscle in pigs. *Journal of Animal Science* 82, 134–134.

Bérard J and Bee G 2010. Effects of dietary L-arginine supplementation to gilts during early gestation on foetal survival, growth and myofiber formation. *Animal* 4, 1680–1687.

Bérard J, Kreuzer M and Bee G 2008. Effect of litter size and birth weight on growth, carcass and pork quality, and their relationship to postmortem proteolysis. *Journal of Animal Science* 86, 2357–2368.

Bérard J, Kreuzer M and Bee G 2010. In large litters birth weight and gender is decisive for growth performance but less for carcass and pork quality traits. *Meat Science* 86, 845–851.

Dwyer CM, Fletcher JM and Stickland NC 1993. Muscle cellularity and postnatal growth in the pig. *Journal of Animal Science* 71, 3339–3343.

Ellis M, Webb AJ, Avery PJ and Brown I 1996. The influence of terminal sire genotype, sex, slaughter weight, feeding regime and slaughter-house on growth performance and carcass and meat quality in pigs and on the organoleptic properties of fresh pork. *Animal Science* 62, 521–530.

Fix JS, Cassidy JP, Holl JW, Herring WO, Culbertson MS and See MT 2010. Effect of piglet birth weight on survival and quality of commercial market swine. *Livestock Science* 132, 98–106.

Foxcroft GR, Bee G, Dixon W, Hahn M, Harding J, Patterson J, Putman T, Sarmento S, Smit M, Tse W-Y and Town SC 2007. Consequences of selection for litter size on piglet development. In *Paradigms in pig science* (ed. J Wiseman, MA Varley, S McOrist and B Kemp), pp. 207–229. Nottingham University Press, Nottingham.

Gondret F, Lefaucheur L, Juin H, Louveau I and Lebret B 2006. Low birth weight is associated with enlarged muscle fiber area and impaired meat tenderness of the longissimus muscle in pigs. *Journal of Animal Science* 84, 93–103.

Gondret F, Lefaucheur L, Louveau I, Lebret B, Pichodo X and Le Cozler Y 2005. Influence of piglet birth weight on postnatal growth performance, tissue lipogenic capacity and muscle histological traits at market weight. *Livestock Production Science* 93, 137–146.

Honikel KO 1998. Reference methods for the assessment of physical characteristics of meat. *Meat Science* 49, 447–457.

Leenhouders JJ, van der Lende T and Knol EF 1999. Analysis of stillbirth in different lines of pig. *Livestock Production Science* 57, 243–253.

Milligan BN, Fraser D and Kramer DL 2002. Within-litter birth weight variation in the domestic pig and its relation to pre-weaning survival, weight gain, and variation in weaning weights. *Livestock Production Science* 76, 181–191.

Nissen PM and Oksbjerg N 2011. Birth weight and postnatal dietary protein level affect performance, muscle metabolism and meat quality in pigs. *Animal* 5, 1382–1389.

Nissen PM, Jorgensen PF and Oksbjerg N 2004. Within-litter variation in muscle fiber characteristics, pig performance, and meat quality traits. *Journal of Animal Science* 82, 414–421.

Pardo C, Kreuzer M and Bee G 2010. Relationship between average litter weight and intralitter weight variability on myogenesis in newborn piglets. *Journal of Dairy Science* 93, 364.

Poore KR and Fowden AL 2004. The effects of birth weight and postnatal growth patterns on fat depth and plasma leptin concentrations in juvenile and adult pigs. *The Journal of Physiology Online* 558, 295–304.

Quesnel H, Brossard L, Valancogne A and Quiniou N 2008. Influence of some sow characteristics on within-litter variation of piglet birth weight. *Animal* 2, 1842–1849.

Quiniou N, Dagorn J and Gaudré D 2002. Variation of piglets' birth weight and consequences on subsequent performance. *Livestock Production Science* 78, 63–70.

Rehfeldt C and Kuhn G 2006. Consequences of birth weight for postnatal growth performance and carcass quality in pigs as related to myogenesis. *Journal of Animal Science* 84, E113–E123.

Rehfeldt C, Tuchscherer A, Hartung M and Kuhn G 2008. A second look at the influence of birth weight on carcass and meat quality in pigs. *Meat Science* 78, 170–175.

Suisag 2012. Zahlen und Projekte 2011. Retrieved January 23, 2013, from <http://www.suisag.ch/Dokumente/tabid/111/Default.aspx>.

Town SC, Patterson JL, Pereira CZ, Gourley G and Foxcroft GR 2005. Embryonic and fetal development in a commercial dam-line genotype. *Animal Reproduction Science* 85, 301–316.

Vallet JL and Freking BA 2006. Changes in fetal organ weights during gestation after selection for ovulation rate and uterine capacity in swine. *Journal of Animal Science* 84, 2338–2345.

van der Waaij EH, Hazeleger W, Soede NM, Laurensen BFA and Kemp B 2010. Effect of excessive, hormonally induced intrauterine crowding in the gilt on fetal development on day 40 of pregnancy. *Journal of Animal Science* 88, 2611–2619.

Wu G, Bazer FW, Wallace JM and Spencer TE 2006. Board-invited review: intrauterine growth retardation: implications for the animal sciences. *Journal of Animal Science* 84, 2316–2337.

Zhao X, Mo D, Li A, Gong W, Xiao S, Zhang Y, Qin L, Niu Y, Guo Y, Liu X, Cong P, He Z, Wang C, Li J and Chen Y 2011. Comparative analyses by sequencing of transcriptomes during skeletal muscle development between pig breeds differing in muscle growth rate and fatness. *PLoS One* 6, e19774.