

Blood Pressure and Physical Activity: Time to Move (On)

Grégoire Wuerzner^{1,2}, Murielle Bochud,³ and Michel Burnier¹

To the Editor: In their article, “Blood Pressure Circadian Pattern and Physical Exercise Assessment by Accelerometer and 7-Day Physical Activity Recall Scale,” Garcia-Ortiz *et al.* report that physical activity recorded as counts per minute with an accelerometer was associated with the dipper pattern in a cohort of healthy participants.¹ This study merits several comments. The interest in the relation of physical activity with 24-hour blood pressure is not new.² Using a similar design but a different accelerometer and an arm ambulatory blood pressure monitoring device instead of a wrist

device, we have previously reported similar association between step count and systolic dipping in a smaller cohort that included mainly (60%) hypertensive participants.³ In our study, however, most of the effect was driven by hypertensive participants. Although the authors have taken into account the use of antihypertensive drugs in their model, adjustments for the hypertensive status or the level of 24-hour blood pressure have not been reported. These variables are important confounders that might have affected the results. Additionally, the authors did not describe how daytime and nighttime ambulatory blood pressure monitoring were defined. Defining these periods (diary, activity, and fixed period) is recommended by guidelines and may influence the results.^{4,5} Finally, we believe that in the era of “quantified self” or “lifelogging” when technology outpaces medical evidence, physical activity should be reported in units, such as steps or kilocalories that are understandable for both the patients and the physician. This will possibly help to set understandable targets, which should help our patients increase their physical activity level.

DISCLOSURE

The authors declared no conflict of interest.

REFERENCES

1. Garcia-Ortiz L, Recio-Rodriguez JI, Puig-Ribera A, Lema-Bartolome J, Ibanez-Jalon E, Gonzalez-Viejo N, Guenaga-Saenz N, Agudo-Conde C, Patino-Alonso MC, Gomez-Marcos MA, Group E. Blood pressure circadian pattern and physical exercise assessment by accelerometer and 7-day physical activity recall scale. *Am J Hypertens* 2014; 27:665–673.
2. Kario K, Schwartz JE, Pickering TG. Ambulatory physical activity as a determinant of diurnal blood pressure variation. *Hypertension* 1999; 34:685–691.
3. Wuerzner G, Bochud M, Zweieracker C, Tremblay S, Pruijm M, Burnier M. Step count is associated with lower nighttime systolic blood pressure and increased dipping. *Am J Hypertens* 2013; 26:527–534.
4. O'Brien E, Asmar R, Beilin L, Imai Y, Mancia G, Mengden T, Myers M, Padfield P, Palatini P, Parati G, Pickering T, Redon J, Staessen J, Stergiou G, Verdecchia P. Practice guidelines of the European Society of Hypertension for clinic, ambulatory and self blood pressure measurement. *J Hypertens* 2005; 23:697–701.
5. Eissa MA, Yetman RJ, Poffenbarger T, Portman RJ. Comparison of arbitrary definitions of circadian time periods with those determined by wrist actigraphy in analysis of ABPM data. *J Hum Hypertens* 1999; 13:759–763.

Correspondence: Grégoire Wuerzner (gregoire.wuerzner@chuv.ch).

¹Service of Nephrology, Lausanne University Hospital, Lausanne, Switzerland; ²Clinical Research Center, Lausanne University Hospital, Lausanne, Switzerland; ³University Institute of Social and Preventive Medicine, Lausanne, Switzerland.

Initially submitted May 1, 2014; accepted for publication May 8, 2014.

doi:10.1093/ajh/hpu118

© American Journal of Hypertension, Ltd 2014. All rights reserved. For Permissions, please email: journals.permissions@oup.com