

ORIGINAL MANUSCRIPT

CCDC26, CDKN2BAS, RTEL1 and TERT Polymorphisms in pediatric brain tumor susceptibility

Maral Adel Fahmideh^{*}, Catharina Lavebratt¹, Joachim Schüz²,
Martin Rööslī^{3,4}, Tore Tynes^{5,6}, Michael A. Grotzer⁷, Christoffer Johansen^{8,9},
Claudia E. Kuehni¹⁰, Birgitta Lantering¹¹, Michaela Prochazka,
Lisbeth S. Schmidt¹² and Maria Feychting

Unit of Epidemiology, Institute of Environmental Medicine, Karolinska Institutet, Box 210, Stockholm SE-171 77, Sweden, ¹Neurogenetics Unit, Department of Molecular Medicine and Surgery, Karolinska Institutet, and Center for Molecular Medicine, Karolinska University Hospital, Stockholm SE-171 76, Sweden, ²Section of Environment and Radiation, International Agency for Research on Cancer (IARC), 69372 Lyon CEDEX 08, France, ³Department of Epidemiology and Public Health, Swiss Tropical and Public Health Institute, Basel 4002, Switzerland, ⁴University of Basel, Basel 4003, Switzerland, ⁵The Cancer Registry of Norway, Oslo N-0304, Norway, ⁶National Institute of Occupational Health, Oslo NO-0033, Norway, ⁷Department of Oncology, University Children's Hospital of Zurich, Zurich 8091, Switzerland, ⁸Unit of Survivorship, The Danish Cancer Society Research Centre, Copenhagen DK-2100, Denmark, ⁹5073 Oncology Clinic, Finsen Centre Rigshospitalet, University of Copenhagen, Copenhagen DK-2100, Denmark, ¹⁰Swiss Childhood Cancer Registry, Institute of Social and Preventive Medicine, University of Bern, Bern 3012, Switzerland, ¹¹Department of Clinical Sciences, Pediatric Oncology, University of Gothenburg, SE 416 85 Gothenburg, Sweden and ¹²Department of Pediatric Oncology, University Hospital Rigshospitalet, Copenhagen DK-2100, Denmark

^{*}To whom correspondence should be addressed. Tel. +46 8 524 839 54; Fax. +46 8 313961; Email: maral.adel.fahmideh@ki.se

Abstract

The role of genetic polymorphisms in pediatric brain tumor (PBT) etiology is poorly understood. We hypothesized that single nucleotide polymorphisms (SNPs) identified in genome-wide association studies (GWAS) on adult glioma would also be associated with PBT risk. The study is based on the Cefalo study, a population-based multicenter case-control study. Saliva DNA from 245 cases and 489 controls, aged 7–19 years at diagnosis/reference date, was extracted and genotyped for 29 SNPs reported by GWAS to be significantly associated with risk of adult glioma. Data were analyzed using unconditional logistic regression. Stratified analyses were performed for two histological subtypes: astrocytoma alone and the other tumor types combined. The results indicated that four SNPs, *CDKN2BAS* rs4977756 ($p = 0.036$), rs1412829 ($p = 0.037$), rs2157719 ($p = 0.018$) and rs1063192 ($p = 0.021$), were associated with an increased susceptibility to PBTs, whereas the *TERT* rs2736100 was associated with a decreased risk ($p = 0.018$). Moreover, the stratified analyses showed a decreased risk of astrocytoma associated with *RTEL1* rs6089953, rs6010620 and rs2297440 ($p_{\text{trend}} = 0.022$, $p_{\text{trend}} = 0.042$, $p_{\text{trend}} = 0.029$, respectively) as well as an increased risk of this subtype associated with *RTEL1* rs4809324 ($p_{\text{trend}} = 0.033$). In addition, SNPs rs10464870 and rs891835 in *CCDC26* were associated with an increased risk of non-astrocytoma tumor subtypes ($p_{\text{trend}} = 0.009$, $p_{\text{trend}} = 0.007$, respectively). Our findings indicate that SNPs in *CDKN2BAS*, *TERT*, *RTEL1* and *CCDC26* may be associated with the risk of PBTs. Therefore, we suggest that pediatric and adult brain tumors might share common genetic risk factors and similar etiological pathways.

Received: January 9, 2015; Revised: April 29, 2015; Accepted: May 20, 2015

© The Author 2015. Published by Oxford University Press. All rights reserved. For Permissions, please email: journals.permissions@oup.com.

Abbreviations

CI	confidence interval
GWAS	genome-wide association studies
OR	odds ratio
PBT	pediatric brain tumor
SNP	single nucleotide polymorphism

Introduction

Brain tumors are the second most common type of pediatric cancer and the leading cause of childhood cancer mortality. The etiology of pediatric brain tumors (PBTs) is poorly understood (1). As in adults (2), the only established risk factors for brain tumors in children are exposure to high doses of ionizing radiation and several inherited disorders, and these cause only a minority of cases. Therefore, it is likely that brain tumorigenesis results from complex interactions between genetic and epigenetic variations in concert with exposure to environmental factors (1).

Although large genetic studies on adult brain tumors have been conducted (3–7), very few and only small studies of brain tumors in children and adolescents have been reported (8–11). In the last few years, four genome-wide association studies (GWAS) on adult glioma identified seven susceptibility loci at 5p15.33 (TERT), 8q24.21 (CCDC26), 9p21.3 (CDKN2A-CDKN2B), 20q13.33 (RTEL1), 11q23.3 (PHLDB1) and 7p11.2 (EGFR) (4–7). However, limited data are available on the role of genetic polymorphisms in the etiology of PBTs, probably because of difficulties in collecting a sufficient number of DNA samples. Considering this lack of knowledge about genetic risk factors for brain tumors in children, it is important to identify germ-line DNA polymorphisms that might influence the susceptibility to PBTs.

The aim of this study, based on the largest series of PBT cases to date, was to test the hypothesis that the single nucleotide polymorphisms (SNPs) identified by GWAS on adult glioma are also associated with the risk of brain tumors in children.

Materials and methods

Study population and procedures

This study is based on the Cefalo study, a large, international, population-based, case-control study of brain tumors in children and adolescents conducted in centers in Sweden, Denmark, Norway and Switzerland. All centers followed a common protocol for data collection, as described in more detail elsewhere (12,13). Eligible cases were children aged 7–19 years during the period 1 April 2004 to 31 August 2008, diagnosed with a primary intracranial brain tumor defined according to the International Classification of Childhood Cancer, third edition (ICCC-3) (14), group III, restricted to the third edition of the International Classification of Diseases for Oncology (ICD-O-3), location C71 and subclassified according to the fourth edition of the World Health Organization (WHO) classification of tumors of the central nervous system (15). Medulloblastoma cases will be the subject of a separate study and therefore have been excluded from the present analysis. Two controls per case were randomly selected from the general population matched to the case by age, sex and geographical region. Interviews were conducted with 352 (82%) cases and 646 (71%) controls. Participants with neurofibromatosis or tuberous sclerosis were excluded from the analyses. The study was approved by the national data protection boards and ethical committees in all participating countries, and written informed consent was obtained from all participants and/or their parents.

The Oragene self-collection kit (DNA Genotek, Ottawa, ON, Canada) was used for saliva collection and DNA extraction, following the manufacturer's recommended protocol. DNA samples are stored at the Karolinska Institutet Biobank. The DNA yield was quantitated by using PicoGreen

(Invitrogen, Carlsbad, CA). Overall, saliva DNA from 245 cases and 489 controls was included in this study.

SNP selection and genotyping

A total of 29 SNPs reported by GWAS to be significantly associated with risk of adult glioma were selected for genotyping (4–7). Genotyping was performed at the Mutation Analysis core Facility, Clinical Research Centre, Huddinge University Hospital, Stockholm, Sweden, with staff blinded to sample status, using the Sequenom iPLEX Gold platform with matrix-assisted laser desorption/ionization-time-of-flight (MALDI-TOF) mass spectrometry. The average success rate was 97% and the concordance rate for duplicate genotyping was 100%.

Statistical analysis

The consistency of allele frequencies with Hardy-Weinberg equilibrium was assessed in the controls for all SNPs using the χ^2 goodness-of-fit test, and $p < 0.001$ was considered statistically significant. Analyses were performed based on the subset of Cefalo subjects who provided saliva sample. Unconditional logistic regression was used to estimate the association between SNPs and PBT susceptibility based on the Cochran-Armitage trend test of additivity (trend) as well as dominant (DOM) and recessive (REC) models, with adjustment for the matching variables (age, sex and country). The allelic frequencies of the genotyped SNPs were compared between cases and controls using the χ^2 test. Analyses were also conducted stratified by astrocytoma alone and the combination of other tumor types, including ependymoma, intracranial embryonal tumors (except medulloblastoma), other gliomas, other specified intracranial neoplasms and unspecified intracranial neoplasms. Country specific analyses were performed to assess consistency across countries. The Wald test was used to evaluate the significance of interactions between SNPs and demographic variables and D' , a measure of the linkage disequilibrium (LD) between the genotyped SNPs, was calculated. Haploblocks were defined based on the default linkage disequilibrium block parameters in Haploview v4.2. Haplotype analyses were performed for the haplotype blocks harboring the SNPs that were found to be associated with PBTs. Haplotypes with a frequency of >1% were considered in the analyses. The effects of specific haplotypes were analyzed if the distribution of all the haplotypes was suggestively different between cases and controls ($p < 0.05$ for all PBTs; $p < 0.1$ for subgroup analyses). Selection of SNPs for the analyses was based on a priori knowledge from GWAS on adults, and therefore odds ratios (ORs) are presented with 95% confidence intervals (CIs). The possibility of false-positive findings was, however, considered by also providing the reference p value for an experiment-wide significance with Bonferroni correction. The analyses were performed using PLINK v1.07 (16) and SAS statistical software version 9.3 (SAS Institute, Cary, NC).

Results

We successfully genotyped 29 SNPs in 245 cases and 489 controls. The distributions of allele frequencies in the controls were in agreement with the Hardy-Weinberg equilibrium. Table 1 shows demographic characteristics of cases and controls and the distributions of diagnostic subtypes. The age and sex distributions were similar in cases and controls. More than 50% of cases were diagnosed with astrocytoma. No significant interactions were detected between SNPs and confounders including age, sex and country (Table 2).

As shown in Table 2, TERT rs2736100 A allele was associated with a decreased risk of PBTs [OR_{DOM} 0.66 (95% CI 0.46–0.93), $p = 0.018$], whereas the SNPs rs4977756 G allele [OR_{DOM} 1.45 (95% CI 1.03–2.06), $p = 0.036$], rs1412829 G allele [OR_{DOM} 1.45 (95% CI 1.02–2.05), $p = 0.037$], rs2157719 C allele [OR_{DOM} 1.53 (95% CI 1.08–2.19), $p = 0.018$] and rs1063192 G allele [OR_{DOM} 1.53 (95% CI 1.07–2.19), $p = 0.021$] in CDKN2BAS were associated with increased susceptibility to these tumors.

The stratified analyses of two histological subtypes indicated that the risk effects of CDKN2BAS rs1063192, rs2157719, rs1412829 and rs4977756 remained significant in patients with astrocytoma

Table 1. Characteristics of cases and controls

Characteristics	Cases	Controls
No. of participants	245	489
Sex		
Males	136 (56%)	261 (53%)
Females	109 (44%)	228 (47%)
Age-group (at reference date)		
7–9 years old	48 (20%)	112 (23%)
10–14 years old	108 (44%)	219 (45%)
15–19 years old	89 (36%)	158 (32%)
Country		
Sweden	106 (43%)	174 (36%)
Norway	24 (10%)	62 (13%)
Denmark	62 (25%)	134 (27%)
Switzerland	53 (22%)	119(24%)
Type of tumor (International Classification of Childhood Cancer-3 group III) ^a		
Astrocytoma (IIIb)	134 (55%)	
Pilocytic astrocytoma	93	
Supependymal giant cell astrocytoma	5	
Pleomorphic xanthoastrocytoma	4	
Diffuse astrocytoma	13	
Anaplastic astrocytoma	11	
Fibrillary astrocytoma	2	
Glioblastoma	5	
Giant cell glioblastoma	1	
Other gliomas (IIIId)	20 (8%)	
Malignant glioma	11	
Oligoastrocytoma	2	
Oligodendroglioma	6	
Anaplastic oligodendroglioma	1	
Ependymoma (IIIa)	19 (8%)	
Subependymoma	2	
Choroid plexus papilloma	4	
Choroid plexus carcinoma	1	
Ependymoma	7	
Papillary ependymoma	1	
Anaplastic ependymoma	4	
Intracranial embryonal tumors (IIIc)	7 (3%)	
CNS primitive neuroectodermal tumor	6	
Neuroepithelioma	1	
Other specified intracranial neoplasms (IIIe)	49 (20%)	
Germinoma	7	
Yolk sac tumor	1	
Teratoma, mature	1	
Haemangioblastoma	1	
Desmoplastic infantile ganglioglioma	2	
Dysembryoplastic neuroepithelial tumor	6	
Ganglioglioma	26	
Anaplastic ganglioglioma	1	
Central neurocytoma	3	
Neurilemoma	1	
Unspecified intracranial neoplasm (IIIIf)	16 (6%)	

^aRestricted to ICD-O-3 location C71, subclassified according to World Health Organization histological subclassification, 2007; patients with neurofibromatosis and tuberous sclerosis were excluded.

($p_{\text{trend}} = 0.036$, $p_{\text{trend}} = 0.034$, $p_{\text{trend}} = 0.044$ and $p_{\text{trend}} = 0.023$, respectively), whereas the protective effect of *TERT* rs2736100 was more evident in patients with other brain tumor subtypes [OR_{DOM} 0.53 (95% CI 0.34–0.85), $p = 0.007$]. Moreover, the stratified analyses showed a decreased risk of astrocytoma associated with the

polymorphisms *RTEL1* rs6089953 A allele, rs6010620 A allele and rs2297440 T allele ($p_{\text{trend}} = 0.022$, $p_{\text{trend}} = 0.042$ and $p_{\text{trend}} = 0.029$, respectively), as well as an increased risk of this subtype associated with the C allele of *RTEL1* rs4809324 ($p_{\text{trend}} = 0.033$). In addition, an increased risk of non-astrocytoma tumor subtypes was associated with the SNPs rs10464870 C allele and rs891835 G allele in *CCDC26* ($p_{\text{trend}} = 0.009$ and $p_{\text{trend}} = 0.007$, respectively) (Table 3).

The non-significant findings, possibly resulting from the limited statistical power of the study, are shown in the online appendix [Supplementary Tables 1–3](#) (available at *Carcinogenesis Online*) and the raw data showing the number of cases and controls for each genotype of significant SNPs are reported in [Supplementary Table 4](#) (available at *Carcinogenesis Online*).

Strong linkage disequilibrium ($D' \geq 0.95$) was observed between three of the genotyped SNPs in *CDKN2BAS* (rs1412829, rs2157719 and rs1063192) and four SNPs in *RTEL1* (rs6089953, rs6010620, rs2297440 and rs4809324). For each of the two blocks, three haplotypes with frequency of >1% were found. The distribution of haplotypes was different for PBT patients compared with controls for the *CDKN2BAS* block ($\chi^2 = 7.0$, $df = 2$ and $p = 0.030$) and showed a tendency to be different for the *RTEL1* block ($\chi^2 = 5.9$, $df = 2$ and $p = 0.053$). The most common haplotype (ATA) of *CDKN2BAS* SNPs had a significant protective effect compared with the other haplotypes combined [OR 0.75 (95% CI 0.60–0.93), $p = 0.009$], whereas the second most common haplotype (GCG) had a significant risk effect [OR 1.32 (95% CI 1.06–1.64), $p = 0.012$]. The haplotype analyses suggested an increased risk of PBTs by increasing the number of risk alleles in *CDKN2BAS* and *RTEL1* SNPs. In the astrocytoma subgroup, the same haploblocks and haplotypes with frequencies of >1% were detected. However, in this subgroup, the distribution of haplotypes in the *RTEL1* block was significantly different between patients and controls ($\chi^2 = 9.0$, $df = 2$ and $p = 0.011$), whereas this difference was not statistically significant in the *CDKN2BAS* block ($\chi^2 = 5.7$, $df = 2$ and $p = 0.059$). In the astrocytoma subgroup, a significant protective effect was observed for the most common haplotype (ATA) of *CDKN2BAS* SNPs compared with the other haplotypes combined [OR 0.73 (95% CI 0.56–0.95), $p = 0.021$], whereas the second most common haplotype (GCG) showed a significant risk effect [OR 1.34 (95% CI 1.02–1.76), $p = 0.036$]. Moreover, in the *RTEL1* block, the second most common haplotype (AATT) had a significant protective effect compared with the other haplotypes combined [OR 0.67 (95% CI 0.47–0.95), $p = 0.023$], whereas the third most common haplotype (GGCC) had a significant risk effect [OR 1.57 (95% CI 1.06–2.34), $p = 0.026$] (Table 4).

Overall, we performed 116 testing procedures as described above. When the Bonferroni correction is applied, the reference p value is 0.0004 for an experiment-wide significance level of 0.05, and 0.0009 for a significance level of 0.10; none of the observed associations met these limits. The consistency of results across countries was investigated and the results of stratified analyses are reported in the online appendix [Supplementary Tables 5–8](#) (available at *Carcinogenesis Online*). No significant differences between countries were observed.

Discussion

The results of this study indicate that several SNPs associated with adult glioma risk are also associated with the risk of PBTs. Our findings suggest that SNPs rs4977756 G allele, rs1412829 G allele, rs2157719 C allele and rs1063192 G allele in *CDKN2BAS* may increase the risk of PBTs, whereas the A allele of *TERT* rs2736100 polymorphism may confer protection against PBTs. In addition, polymorphisms rs6089953 A allele, rs6010620 A allele

Table 2. Summary results for SNPs associated with pediatric brain tumors

SNP	Chr.	Gene	Location (bp)	Minor allele	MAF ^a in cases	MAF ^a in controls	Model	OR ^b	95% CI	P	CHISQ	Pinteraction ^c
rs2736100	5	TERT	1286516	A	0.47	0.50	Dominant Recessive Additive Allelic	0.66 1.19 0.89	0.46–0.93 0.82–1.71 0.72–1.12	0.018 0.359 0.351		0.702
rs1063192	9	CDKN2BAS CDKN2B	22003367	G	0.52	0.45	Dominant Recessive Additive Allelic	1.53 1.36 1.31	1.07–2.19 0.95–1.95 1.05–1.63	0.021 0.095 0.015	0.94	0.889
rs2157719	9	CDKN2BAS	22033366	C	0.51	0.44	Dominant Recessive Additive Allelic	1.53 1.36 1.32	1.08–2.19 0.94–1.97 1.06–1.64	0.009 0.018 0.099 0.014	6.64	0.825
rs1412829	9	CDKN2BAS	22043926	G	0.50	0.43	Dominant Recessive Additive Allelic	1.45 1.38 1.29	1.02–2.05 0.95–1.99 1.04–1.61	0.037 0.089 0.021	6.67	0.734
rs4977756	9	CDKN2BAS	22068652	G	0.49	0.42	Dominant Recessive Additive Allelic	1.45 1.30 1.27	1.03–2.06 0.89–1.91 1.02–1.59	0.014 0.036 0.176 0.032	6.04	0.954
										0.024	5.09	

^aMAF, minor allele frequency.^bOR adjusted for age, sex and country.^cP value for interactions between SNPs and demographic variables including age, sex and country.

Table 3. Summary results for SNPs associated with pediatric brain tumors stratified by histological subtypes

SNP	Chr.	Gene	Location (bp)	Minor allele	MAF ^a in cases	MAF ^a in controls	Model	OR ^b	95% CI	P	CHISQ
Astrocytoma											
rs1063192	9	CDKN2BAS CDKN2B	22003367	G	0.52	0.45	Dominant Recessive Additive Allelic	1.84 1.21 1.34	1.15-2.94 0.76-1.91 1.02-1.76	0.011 0.419 0.036	
rs2157719	9	CDKN2BAS	22033366	C	0.51	0.44	Dominant Recessive Additive Allelic	1.75 1.26 1.34	1.11-2.77 0.79-1.99 1.02-1.76	0.038 0.016 0.329	4.29
rs1412829	9	CDKN2BAS	22043926	G	0.50	0.43	Dominant Recessive Additive Allelic	1.64 1.28 1.32	1.05-2.57 0.81-2.03 1.01-1.74	0.029 0.294 0.034	4.43
rs4977756	9	CDKN2BAS	22068652	G	0.5	0.42	Dominant Recessive Additive Allelic	1.85 1.25 1.38	1.17-2.91 0.78-2.01 1.05-1.82	0.008 0.358 0.023	4.09
rs6089953	20	RTEL1	62291008	A	0.18	0.25	Dominant Recessive Additive Allelic	0.64 0.49 0.67	0.43-0.96 0.19-1.31 0.48-0.95	0.032 0.157 0.022	5.02
rs6010620	20	RTEL1	62309839	A	0.19	0.25	Dominant Recessive Additive Allelic	0.66 0.59 0.71	0.44-0.99 0.24-1.45 0.50-0.99	0.048 0.254 0.042	5.46
rs2297440	20	RTEL1	62312299	T	0.17	0.24	Dominant Recessive Additive Allelic	0.64 0.52 0.68	0.41-0.98 0.19-1.37 0.47-0.96	0.038 0.187 0.029	4.27
rs4809324	20	RTEL1	62318220	C	0.15	0.10	Dominant Recessive Additive Allelic	1.54 2.94 1.54	0.98-2.39 0.78-11.14 1.04-2.28	0.060 0.112 0.033	5.18
Other											
rs2736100	5	TERT	1286516	A	0.44	0.50	Dominant Recessive Additive Allelic	0.54 1.07 0.78	0.34-0.85 0.64-1.77 0.57-1.07	0.007 0.802 0.122	
rs10464870	8	CCDC26	130477823	C	0.32	0.23	Dominant Recessive Additive Allelic	1.70 1.78 1.53	1.11-2.60 0.87-3.66 1.11-2.11	0.014 0.115 0.009	2.19
rs891835	8	CCDC26	130491752	G	0.34	0.24	Dominant Recessive Additive Allelic	1.59 2.32 1.55	1.04-2.44 1.18-4.57 1.13-2.14	0.004 0.032 0.007	8.11
											8.65

^aMAF, minor allele frequency.^bOR adjusted for age, sex, and country.

Table 4. Haplotype analysis of SNPs in *CDKN2BAS* and *RTEL1*

SNPs	Haplotype	Frequency	OR ^a	95% CI	P
CDKN2BAS: rs1412829, rs2157719, rs1063192	ATA	0.52	0.75	0.60–0.93	0.009
	GCG	0.45	1.32	1.06–1.64	0.012
	ATG	0.02	0.89	0.56–1.43	0.814
RTEL1: rs6089953, rs6010620, rs2297440, rs4809324	GGCT	0.65	1.03	0.81–1.32	0.787
	AATT	0.23	0.79	0.61–1.02	0.074
	GGCC	0.11	1.39	0.99–1.95	0.055
Astrocytoma					
CDKN2BAS: rs1412829, rs2157719, rs1063192	ATA	0.53	0.73	0.56–0.95	0.021
	GCG	0.44	1.34	1.02–1.76	0.036
	ATG	0.02	0.79	0.21–2.94	0.725
RTEL1: rs6089953, rs6010620, rs2297440, rs4809324	GGCT	0.65	1.06	0.81–1.39	0.676
	AATT	0.23	0.67	0.47–0.95	0.023
	GGCC	0.11	1.57	1.06–2.34	0.026

^aORs for haplotype compared with all other haplotypes adjusted for age, sex and country.

and rs2297440 T allele in *RTEL1* were associated with decreased susceptibility to astrocytoma, whereas the C allele of *RTEL1* rs4809324 was associated with an increased risk of this subtype. Furthermore, an increased risk of non-astrocytoma tumor subtypes associated with polymorphisms *CCDC26* rs10464870 C allele and rs891835 G allele was detected. Our findings suggest that genetic risk profiles of PBTs differ by histology.

To our knowledge, this study represents the largest series of pediatric brain tumor cases assembled for genetic association testing to date. The association between the 29 SNPs investigated in this study and the risk of PBTs has not been examined in previous studies (8–11). The SNPs were selected a priori for analyses in this study based on findings in GWAS on adult glioma, and our significant findings of associations between SNPs in *CDKN2BAS*, *TERT*, *RTEL1* and *CCDC26*, and risk of PBTs were consistent with findings on adult brain tumors with respect to the direction of ORs for the minor alleles (4–7).

CDKN2BAS (*ANRIL*) encodes antisense non-coding RNA in the *INK4* locus which is a long non-coding RNA (ncRNA). The exact function of *CDKN2BAS* is unclear, but it is known to be involved in regulating the expression of *CDKN2A* and *CDKN2B* genes that encode cyclin-dependent kinase inhibitors and block cell cycle division during the G1/S phase. Therefore, *CDKN2BAS* has a regulatory role in the context of cellular proliferation, and its alterations result in abnormal self-renewing capabilities typical of cancer cells (17,18). Germ-line mutations in *CDKN2BAS* predispose to a wide variety of human cancers (19,20).

Telomerase reverse transcriptase (*TERT*) is a catalytic subunit of telomerase that maintains telomere by adding telomeric repeat sequences onto chromosome ends. Telomerase expression can prevent telomere erosion in most eukaryotic cells, and cancer cells can prevent telomere loss through the abnormal upregulation of telomerase (21). The mutant allele of *TERT* rs2736100, which is an intronic polymorphism, may downregulate telomerase expression and consequently decrease the risk of brain tumors.

RTEL1 produces regulator of telomere elongation helicase 1 which is essential for telomere maintenance and genome stability by preventing homologous recombination (22). Polymorphisms in *RTEL1* are correlated with high grade glioma in adults (4,5,7,23). In contrast, *CCDC26* variations are associated with low grade tumors (4,5,23). *CCDC26* encodes a retinoic acid-dependent regulator of cell differentiation and death. *CCDC26* increases apoptosis induced by death stimuli in neuroblastoma cells (24) and in glioblastoma cells with downregulation of telomerase activity (25).

The majority of genetic variations found in this study to be associated with the risk of PBTs are related to telomerase activity which has an important role in the initiation and progression of gliomas (26). Moreover, it has been shown that telomerase expression is related to high grade gliomas and poor survival (27,28). Thus, telomerase could be considered as a therapeutic target for brain tumors (29,30).

The aim of this study was to provide evidence of the associations between SNPs and PBT risk, and not to investigate the mechanisms behind such associations; nevertheless the fact that we did not consider the effect of environmental risk factors represents a limitation of this work. Therefore, additional studies are needed to examine potentially relevant gene-environment interactions and to explore the mechanisms through which these genetic polymorphisms influence cancer susceptibility.

The present study was conducted based on a specific hypothesis that may lead to detection of clinically meaningful risk and protective factors. Moreover, the selection of SNPs for analysis was based on a priori knowledge from GWAS on adults, and therefore Bonferroni correction may be overly conservative and may make researchers miss important findings (31). To be able to evaluate potential false-positive findings, reference p values with Bonferroni corrections have been presented and the consistency of results across four countries has been reported. No significant differences between countries were observed. Replication studies are necessary to confirm these results in larger sample sizes.

In conclusion, the present findings indicate that SNPs in *CDKN2BAS* are associated with increased susceptibility to PBTs, whereas *TERT* polymorphisms may decrease the risk of these tumors. Moreover, polymorphisms in *RTEL1* and *CCDC26* genes are associated with the risk of astrocytoma and non-astrocytoma subtypes, respectively. Thus, we suggest that pediatric and adult brain tumors might share common genetic risk factors and similar etiological pathways.

Supplementary material

Supplementary Tables 1–8 can be found at <http://carcin.oxfordjournals.org/>.

Funding

Swedish Council for Working Life and Social Research (2004-0504 and 2007-0224); Swedish Research Council (K2008-70X-15366-04-3); Swedish Cancer Society (09 0666); Swedish

Childhood Cancer Society (PROJ06/050 and PROJ09/086); Swedish Radiation Protection Authority (SSI P 1572); Danish Strategic Research Council (2103-05-0006 and 2064-04-0010); Swiss Federal Office of Public Health (05.001626); Swiss Research Foundation on Mobile Communication (A2006.18); Swiss National Science Foundation (PDFMP3_122873); Research Council of Norway (175163/V40).

Acknowledgements

We gratefully acknowledge collaboration with and support from clinicians and other hospital staff in all countries, as well as assistance from national and local cancer registers with identification of patients. We also acknowledge the skillful work of the research nurses, interviewers, and research assistants in all countries. Finally, we acknowledge all those with whom we have collaborated previously within the CEFALO study. Please see Aydin et al. for an exhaustive list (12).

Conflict of Interest Statement: None declared.

URLs: PLINK: <http://pngu.mgh.harvard.edu/~purcell/plink/>; SAS: <http://www.sas.com/>

References

- Pollack, I.F. et al. (2011) Childhood brain tumors: epidemiology, current management and future directions. *Nat. Rev. Neurol.*, 7, 495–506.
- Fisher, J.L. et al. (2007) Epidemiology of brain tumors. *Neurol. Clin.*, 25, 867–890.
- Adel Fahmideh, M. et al. (2014) Association between DNA repair gene polymorphisms and risk of glioma: a systematic review and meta-analysis. *Neuro Oncol.*, 16, 807–814.
- Rajaraman, P. et al. (2012) Genome-wide association study of glioma and meta-analysis. *Hum. Genet.*, 131, 1877–1888.
- Sanson, M. et al. (2011) Chromosome 7p11.2 (EGFR) variation influences glioma risk. *Hum. Mol. Genet.*, 20, 2897–2904.
- Shete, S. et al. (2009) Genome-wide association study identifies five susceptibility loci for glioma. *Nat. Genet.*, 41, 899–904.
- Wrensch, M. et al. (2009) Variants in the CDKN2B and RTEL1 regions are associated with high-grade glioma susceptibility. *Nat. Genet.*, 41, 905–908.
- Jeon, S. et al. (2013) Genetic variants of AICDA/CASP14 associated with childhood brain tumor. *Genet. Mol. Res.*, 12, 2024–2031.
- Salnikova, L.E. et al. (2010) Association study of xenobiotic detoxication and repair genes with malignant brain tumors in children. *Acta Naturae*, 2, 58–65.
- Searles Nielsen, S. et al. (2010) Childhood brain tumors, residential insecticide exposure, and pesticide metabolism genes. *Environ. Health Perspect.*, 118, 144–149.
- Sirachainan, N. et al. (2008) Folate pathway genetic polymorphisms and susceptibility of central nervous system tumors in Thai children. *Cancer Detect. Prev.*, 32, 72–78.
- Aydin, D. et al. (2011) Mobile phone use and brain tumors in children and adolescents: a multicenter case-control study. *J. Natl Cancer Inst.*, 103, 1264–1276.
- Christensen, J.S. et al. (2012) Brain tumors in children and adolescents and exposure to animals and farm life: a multicenter case-control study (CEFALO). *Cancer Causes Control*, 23, 1463–1473.
- Steliarova-Foucher, E. et al. (2005) International Classification of Childhood Cancer, third edition. *Cancer*, 103, 1457–1467.
- Louis, D.N. et al. (2007) The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol.*, 114, 97–109.
- Purcell, S. et al. (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet.*, 81, 559–575.
- Cunnington, M.S. et al. (2010) Chromosome 9p21 SNPs Associated with Multiple Disease Phenotypes Correlate with ANRIL Expression. *PLoS Genet.*, 6, e1000899.
- Wapinski, O. et al. (2011) Long noncoding RNAs and human disease. *Trends Cell Biol.*, 21, 354–361.
- Gu, F. et al. (2013) Common genetic variants in the 9p21 region and their associations with multiple tumours. *Br. J. Cancer.*, 108, 1378–1386.
- Turnbull, C. et al. (2010) Genome-wide association study identifies five new breast cancer susceptibility loci. *Nat. Genet.*, 42, 504–507.
- Baird, D.M. (2010) Variation at the TERT locus and predisposition for cancer. *Expert Rev. Mol. Med.*, 12, e16.
- Barber, L.J. et al. (2008) RTEL1 maintains genomic stability by suppressing homologous recombination. *Cell*, 135, 261–271.
- Simon, M. et al. (2010) Genetic risk profiles identify different molecular etiologies for glioma. *Clin. Cancer Res.*, 16, 5252–5259.
- Jiang, M. et al. (2008) Retinoic acid induces caspase-8 transcription via phospho-CREB and increases apoptotic responses to death stimuli in neuroblastoma cells. *Biochim. Biophys. Acta.*, 1783, 1055–1067.
- Das, A. et al. (2007) Differentiation decreased telomerase activity in rat glioblastoma C6 cells and increased sensitivity to IFN-gamma and taxol for apoptosis. *Neurochem. Res.*, 32, 2167–2183.
- Shervington, A. et al. (2006) Glioma: what is the role of c-Myc, hsp90 and telomerase? *Mol. Cell. Biochem.*, 283, 1–9.
- Tabori, U. et al. (2006) Human telomere reverse transcriptase expression predicts progression and survival in pediatric intracranial ependymoma. *J. Clin. Oncol.*, 24, 1522–1528.
- Tchirkov, A. et al. (2003) Clinical implications of quantitative real-time RT-PCR analysis of hTERT gene expression in human gliomas. *Br. J. Cancer*, 88, 516–520.
- Kim, J.H. et al. (2013) Ependymoma in children: molecular considerations and therapeutic insights. *Clin. Transl. Oncol.*, 15, 759–765.
- Wong, V.C. et al. (2010) Telomerase inhibition as a novel therapy for pediatric ependymoma. *Brain Pathol.*, 20, 780–786.
- Rothman, K.J. (1990) No adjustments are needed for multiple comparisons. *Epidemiology*, 1, 43–46.