

RESEARCH PAPER

# Duration and extension of anatomical changes in wood structure after cambial injury

Estelle Arbellay<sup>1,2,\*</sup>, Patrick Fonti<sup>3</sup> and Markus Stoffel<sup>1,2</sup>

<sup>1</sup> Laboratory of Dendrogeomorphology, Institute of Geological Sciences, University of Berne, Baltzerstrasse 1+3, 3012 Berne, Switzerland

<sup>2</sup> Institute for Environmental Sciences, University of Geneva, Chemin de Drize 7, 1227 Carouge-Geneva, Switzerland

<sup>3</sup> Landscape Dynamics, Swiss Federal Research Institute WSL, Zürcherstrasse 111, 8903 Birmensdorf, Switzerland

\* To whom correspondence should be addressed. E-mail: estelle.arbellay@dendrolab.ch

Received 26 November 2011; Revised 6 January 2012; Accepted 31 January 2012

## Abstract

Cambial injury has been reported to alter wood structure in broad-leaved trees. However, the duration and extension of associated anatomical changes have rarely been analysed thoroughly. A total of 18 young European ash (*Fraxinus excelsior* L.) trees injured on the stem by a spring flood were sampled with the aim of comparing earlywood vessels and rays formed prior to and after the scarring event. Anatomical and hydraulic parameters were measured in five successive rings over one-quarter of the stem circumference. The results demonstrate that mechanical damage induces a decrease in vessel lumen size (up to 77%) and an increase in vessel number (up to 475%) and ray number (up to 115%). The presence of more earlywood vessels and rays was observed over at least three years after stem scarring. By contrast, abnormally narrow earlywood vessels mainly developed in the first ring formed after the event, increasing the thickness-to-span ratio of vessels by 94% and reducing both xylem relative conductivity and the index for xylem vulnerability to cavitation by 54% and 32%, respectively. These vessels accumulated in radial groups in a 30° sector immediately adjacent to the wound, raising the vessel grouping index by 28%. The wound-induced anatomical changes in wood structure express the functional need of trees to improve xylem hydraulic safety and mechanical strength at the expense of water transport. Xylem hydraulic efficiency was restored in one year, while xylem mechanical reinforcement and resistance to cavitation and decay lasted over several years.

**Key words:** Cambial injury, earlywood vessel, European ash, *Fraxinus excelsior*, ray, ring-porous, wood anatomy.

## Introduction

Trees are constantly exposed to environmental stresses that may impair their ability to metabolize normally (Schweingruber, 2007). While both heredity and the environment influence the physiological processes that control tree growth (Kozlowski *et al.*, 1991), it is acknowledged that tree adaptability to the environment can be assessed by analysing xylem cells across series of annual rings (Denn and Dodd, 1981; Sass and Eckstein, 1995; Fonti *et al.*, 2010). When the cambium of trees is injured, normal cambial activity is locally disrupted, which is followed by a series of defence and wound healing processes including compartmentalization of decay and formation of callus tissue (Shigo, 1984; Neely, 1988; Blanchette, 1992; Larson,

1994; Fink, 1999). The healing proceeds from the wound margin inward in order to shield the exposed xylem with new healthy tissue.

Wood anatomical investigations of injured broad-leaved trees have mostly involved diffuse-porous species experimentally wounded by partial girdling, pinning or drilling of the stem (Rier and Shigo, 1972; Sharon, 1973; Bauch *et al.*, 1980; Rademacher *et al.*, 1984; Kuroda and Shimaji, 1985; Lev-Yadun, 1994; Stobbe *et al.*, 2002). Broad-leaved trees with naturally inflicted injuries have only rarely been studied at the cellular level, with the exception of recent research aiming at retrieving environmental information on natural hazards or forest fires from xylem cells (Arbellay

et al., 2010; Ballesteros et al., 2010; Bigio et al., 2010; Kames et al., 2011). It is well established that ecologically relevant information can be obtained retrospectively from vessels of broad-leaved trees. Earlywood vessels of ring-porous species, in particular, have yielded successful results when screened for signals induced by climate (Woodcock, 1989; Pumijumngong and Park, 1999; García-González and Eckstein, 2003; Fonti and García-González, 2004; Tardif and Conciatori, 2006; Fonti et al., 2007), drought (Corcuera et al., 2004; Eilmann et al., 2006; Galle et al., 2010), flooding (Yanosky, 1983; Astrade and Bégin, 1997; St George and Nielsen, 2003), and insect defoliation (Huber, 1993; Asshoff et al., 1998–1999). Nevertheless, comparative studies focusing on wounding of ring-porous species and its functional implications are scarce.

The aim of this research is to identify and quantify anatomical changes in the wood structure of European ash (*Fraxinus excelsior* L.) caused by cambial injury and crystallized in the newly formed xylem. Young trees injured on the stem by a spring flood were sampled to compare earlywood vessels and rays formed prior to and after the scarring event. Anatomical and hydraulic parameters were analysed in both radial and tangential directions so as to determine the duration and extension of wound effects.

## Materials and methods


### Field campaign and sample preparation

Trees were sampled in summer 2009 along the St-Barthélemy torrent (Valais, Swiss Alps, 46°11' N, 7°00' E, 570 m a.s.l.), located in the upper Rhone river valley. The riparian vegetation at the site was predominantly composed of grey alder (*Alnus incana* L.) Moench and further included European ash (*Fraxinus excelsior* L.), sycamore maple (*Acer pseudoplatanus* L.), and goat willow (*Salix caprea* L.). Stem cross-sections were collected from 18 young *F. excelsior* trees displaying one elongated scar (140×2.5 cm maximum) up to the xylem and oriented to the flow direction of the torrent. All trees were injured in spring 2007 before the start of the growing season on roughly half of the stem circumference. The average tree age was  $17.11 \pm 4.46$  years and the average stem circumference at breast height was  $8.03 \pm 2.32$  cm. One sample per tree was taken at the mid-length of the injury. The average disc radius was  $0.86 \pm 0.28$  cm.

Each sample was sectioned with a chisel to obtain a disc quarter (Fig. 1a). After dating the rings, 15  $\mu$ m thick transverse sections of the quarters were cut using a Reichert sliding microtome. The microsections were then stained with a 1% safranin and astrablue solution, rinsed with water, alcohols, and xylol, and permanently mounted on microscope slides using Canada balsam.

### Xylem anatomical and hydraulic parameters

Earlywood vessels and rays were studied over the whole disc quarter in five successive rings for each tree (90 rings in total): the injury ring (Ir 1) built during the growing season following wounding, two control rings (Cr 1 and Cr 2) laid down previously and two post-injury rings (Pr 1 and Pr 2) formed subsequently (Fig. 1b). Anatomical measurements of the cells were generally performed from images of the microsections captured at 25× magnification with a digital camera mounted on a light microscope. The software WinCELL Pro V 2004a (Régent Instruments Inc., 2004) was used to measure the number of earlywood vessels and, for each of them, the wall thickness and the lumen area, as



**Fig. 1.** (a) Location of the disc quarter extracted from each sample. The dashed line shows the position of the cambium at the time of wounding. A callus pad (CP) started to grow over the scar from the wound margin level (star). Care was taken to avoid the callus pad and the decayed area (DA) so as to obtain an undisrupted and sound time series of xylem cells. Average disc radius ( $r$ ) is given. (b) Schematic view of the disc quarter with the different rings and sectors used for analysis. The 5° sectors can be grouped in three 30° sectors (A, B, C). Sector A is the closest to the injury. Measurements of the arc length of sectors were made on the outer edge of samples and then averaged. Ring types: Cr, control ring; Ir, injury ring; Pr, post-injury ring.

well as the radial and tangential lumen diameter. The vessels were then converted into circular conduits following White's (2006) equation  $d = [32 (ab)^3 / (a^2 + b^2)]^{1/4}$ , where  $d$  is the circular lumen diameter, and  $a$  and  $b$  the radial and tangential lumen diameter, respectively. The number of rays was manually surveyed from the images by counting the rays crossing an ideal line in the middle of the rings. The position (coordinates) of cells was used to determine vessel and ray numbers as well as average values of vessel lumen size in 5° sectors with increasing tangential distance from the wound (Fig. 1b). To our knowledge, wood anatomical features in the context of mechanical damage have never been continuously studied in the tangential direction. They have only been locally analysed in the injured area or, at best, in different radial segments (Arbellay et al., 2010; Bigio et al., 2010; Delvaux et al., 2010).

In addition, to evaluate the impact of wounding on tree hydraulic architecture, xylem relative conductivity (Zimmermann, 1983) and three indicators of xylem vulnerability to cavitation were calculated. Xylem relative conductivity (REC), i.e. xylem hydraulic efficiency per unit area, was obtained using a Hagen-Poiseuille modified equation (Van den Oever et al., 1981):  $REC = VF (AVLD/2)^4$ , where  $AVLD$  is the average vessel (circular) lumen diameter and  $VF$  the vessel frequency, i.e. the total number of vessels per (earlywood) unit area. Xylem vulnerability to hydraulic failure was determined by calculating the xylem vulnerability index (VUL), the thickness-to-span ratio (THS) of vessels, and the vessel grouping index (VG). The xylem vulnerability index was considered to assess xylem safety from embolism, as proposed by Carlquist (1977):  $VUL = AVLD/VF$ . The thickness-to-span ratio of vessels, i.e. the intervessel wall thickness divided by the vessel (circular) lumen diameter, is an indicator of cell mechanical support against implosion (Hacke et al., 2001). THS was measured at 50× magnification on 30 vessels for each ring (150 vessels in total). Finally, the vessel grouping index, i.e. the total number of vessels divided by the total number of vessel groups (including solitary and grouped vessels), was used as an indicator of alternative water pathways in case of hydraulic failure (Carlquist, 2001).

## Results

Table 1 provides information on xylem anatomical and hydraulic parameters for the five rings investigated. Average vessel lumen area (*AVLA*) and average vessel lumen diameter (*AVLD*) were respectively greater than 1000  $\mu\text{m}^2$  and 150  $\mu\text{m}$  in all rings, except in the injury ring (Ir 1) where both values were lower. The three rings built after cambial injury (Ir 1, Pr 1, and Pr 2) counted more earlywood vessels and rays than the two control rings (Cr 1 and Cr 2) (Table 1). No significant anatomical and hydraulic changes were statistically detected between the two control rings (Table 2, ANOVA test,  $P < 0.05$ ). However, significant ( $P < 0.05$ ) to highly significant ( $P < 0.001$ ) differences were found between Cr 2 and the three rings of the callus tissue.

Ir 1 showed the most evident response to wounding, with lower values for *AVLA* (35%) and *AVLD* (22%) as well as greater values for vessel number (119%) and ray number (33%). The formation of more and narrower earlywood vessels substantially reduced xylem relative conductivity (*REC*) and the xylem vulnerability index (*VUL*) by 54% ( $P < 0.001$ ) and 32% ( $P < 0.01$ ), respectively (Table 2). The thickness-to-span ratio (*THS*) of vessels increased by 94% ( $P < 0.01$ ) and the vessel grouping index (*VG*) by 28%

( $P < 0.001$ ). The wound-induced anatomical changes in wood structure were stronger close to the wound margin and were approximately limited to the extent of sector A (Fig. 2). *AVLA* decreased up to 77% in this 30° sector, whereas vessel number increased up to 475% and ray number up to 115%. Earlywood vessels with a lumen diameter less than 80  $\mu\text{m}$  predominantly accumulated in sector A (Fig. 3).

Pr 1 and Pr 2, similarly to Ir 1, were composed of more earlywood vessels and rays in comparison with the control rings (Table 2). Vessel number was significantly larger in Pr 1 (50%) and Pr 2 (41%) though less than in Ir 1 (119%). In both post-injury rings, it displayed a tangential constant pattern as opposed to the sharp increase observed in sector A of the injury ring (Fig. 2). Ray number was also larger in Pr 1 (48%) and Pr 2 (46%) (Table 2), showing a rather steady increase toward the wound margin (Fig. 2). After careful examination of the callus tissue, most of the rays counted in the middle of Pr 1 were seen to originate in the late portion of Ir 1. Finally, it was noteworthy that narrower earlywood vessels almost exclusively developed in sector A of the injury ring (Fig. 2). The post-injury rings, by contrast, recovered from the diminution of vessel lumen size in sector A and were then built of wider conduits in sectors

**Table 1.** Xylem anatomical and hydraulic variables analysed over the whole disc quarter for the five rings investigated


	Cr 1 Mean $\pm$ SD	Cr 2 Mean $\pm$ SD	Ir 1 Mean $\pm$ SD	Pr 1 Mean $\pm$ SD	Pr 2 Mean $\pm$ SD
<i>AVLA</i> ( $\mu\text{m}^2$ )	1191.84 $\pm$ 229.42	1267.27 $\pm$ 193.43	825.74 $\pm$ 217.08	1350.51 $\pm$ 239.45	1377.75 $\pm$ 432.56
<i>AVLD</i> ( $\mu\text{m}$ )	173.44 $\pm$ 18.31	179.72 $\pm$ 15.54	140.20 $\pm$ 20.26	185.48 $\pm$ 16.58	184.29 $\pm$ 30.60
<i>VN</i>	82.17 $\pm$ 35.75	82.83 $\pm$ 39.42	181.11 $\pm$ 101.39	124.47 $\pm$ 58.70	116.50 $\pm$ 52.12
<i>VF</i> ( $\text{mm}^{-2}$ )	34.86 $\pm$ 6.73	33.13 $\pm$ 6.63	43.12 $\pm$ 16.47	35.71 $\pm$ 5.90	36.20 $\pm$ 8.25
<i>REC</i> ( $\text{mm}^2$ )	2.03E-03 $\pm$ 7.76E-04	2.18E-03 $\pm$ 6.10E-04	1.01E-03 $\pm$ 3.85E-04	2.68E-03 $\pm$ 8.80E-04	2.68E-03 $\pm$ 1.25E-03
<i>VUL</i> ( $\text{mm}^3$ )	5.22E-03 $\pm$ 1.04E-03	5.69E-03 $\pm$ 1.47E-03	3.85E-03 $\pm$ 1.88E-03	5.37E-03 $\pm$ 1.24E-03	5.51E-03 $\pm$ 2.16E-03
<i>THS</i> ( $\mu\text{m}$ )	0.06 $\pm$ 0.03	0.06 $\pm$ 0.01	0.11 $\pm$ 0.10	0.06 $\pm$ 0.03	0.06 $\pm$ 0.03
<i>VG</i>	1.20 $\pm$ 0.07	1.20 $\pm$ 0.08	1.53 $\pm$ 0.35	1.28 $\pm$ 0.10	1.25 $\pm$ 0.10
<i>RN</i>	117.50 $\pm$ 41.70	126.56 $\pm$ 46.95	167.89 $\pm$ 53.07	187.12 $\pm$ 52.57	184.50 $\pm$ 57.75

Variables: *AVLA*, average vessel lumen area; *AVLD*, average vessel lumen diameter; *VN*, vessel number; *VF*, vessel frequency; *REC*, xylem relative conductivity; *VUL*, xylem vulnerability index; *THS*, thickness-to-span ratio of vessels; *VG*, vessel grouping index; *RN*, ray number. Ring types: Cr, control ring; Ir, injury ring; Pr, post-injury ring. Values are averaged over 18 trees.

**Table 2.** One-way ANOVA results when comparing xylem anatomical and hydraulic variables between rings


	Cr 1–Cr 2		Cr 2–Ir 1		Cr 2–Pr 1		Cr 2–Pr 2	
	<i>P</i> -value	Change (%)	<i>P</i> -value	Change (%)	<i>P</i> -value	Change (%)	<i>P</i> -value	Change (%)
<i>AVLA</i> ( $\mu\text{m}^2$ )	0.294	+6	<0.001	–35	0.265	+7	0.347	+9
<i>AVLD</i> ( $\mu\text{m}$ )	0.275	+4	<0.001	–22	0.297	+3	0.593	+3
<i>VN</i>	0.958	+1	0.001	+119	0.019	+50	0.044	+41
<i>VF</i> ( $\text{mm}^{-2}$ )	0.702	–5	0.023	+30	0.234	+8	0.269	+9
<i>REC</i> ( $\text{mm}^2$ )	0.279	+7	<0.001	–54	0.057	+23	0.159	+23
<i>VUL</i> ( $\text{mm}^3$ )	0.534	+9	0.003	–32	0.495	–6	0.791	–3
<i>THS</i> ( $\mu\text{m}$ )	0.226	–11	0.006	+94	0.864	+2	0.709	–4
<i>VG</i>	0.946	+1	<0.001	+28	0.016	+7	0.125	+5
<i>RN</i>	0.545	+8	0.018	+33	0.001	+48	0.005	+46

Variables: *AVLA*, average vessel lumen area; *AVLD*, average vessel lumen diameter; *VN*, vessel number; *VF*, vessel frequency; *REC*, xylem relative conductivity; *VUL*, xylem vulnerability index; *THS*, thickness-to-span ratio of vessels; *VG*, vessel grouping index; *RN*, ray number. Ring types: Cr, control ring; Ir, injury ring; Pr, post-injury ring. Significant results appear in bold.



**Fig. 2.** Line plots displaying radial and tangential variations in xylem anatomical parameters. The five rings investigated were examined in distinct sectors with increasing tangential distance from the wound: in 5° sectors for vessel features (size and number) and in 10° sectors for ray number. Measurements of the arc length of sectors were made on the outer edge of samples and then averaged. Values are averaged over 18 trees. Variable: *AVLA*, average vessel lumen area. Ring types: Cr, control ring; Ir, injury ring; Pr, post-injury ring.

B and C. ANOVA results were not significant when comparing *AVLA* and *AVLD* between Cr 2 and the two post-injury rings (Table 2). They indicated a slight increase in vessel lumen size comparable with that between the two control rings. Moreover, the majority of conduits in both ring types had a lumen diameter ranging from 160–280 μm (Fig. 3). Earlywood vessels in Pr 1 and Pr 2 enhanced xylem relative conductivity (*REC*) and reduced the xylem



**Fig. 3.** Histograms of vessel size distribution for each ring type. Data were split into equal-sized classes of vessel lumen diameter. This variable was analysed in three 30° sectors (A, B, C). Sector A is the closest to the injury. Values are averaged over 18 trees. For the control rings and post-injury rings, values are averaged over the two rings that were considered in each ring type. Ring types: Cr, control ring; Ir, injury ring; Pr, post-injury ring.

vulnerability index (*VUL*) though not significantly. The thickness-to-span ratio (*THS*) of vessels reached pre-wounding values in both post-injury rings, while the vessel grouping index (*VG*) was significantly higher by 7% ( $P < 0.05$ ) in Pr 1, but was no longer different from Cr 1 and Cr 2 in Pr 2 (Table 2).


## Discussion

### Wound-induced anatomical changes in wood structure

The results of this study confirm that wounding induces the formation of narrower vessels (Aloni and Zimmermann, 1984; Rademacher et al., 1984; Kuroda and Shimaji, 1985; Lowerts et al., 1986; Lev-Yadun and Aloni, 1993; Arbellay et al., 2010; Ballesteros et al., 2010). Regenerative earlywood vessels were found to be much narrower in sector A of the injury ring (Fig. 4a). The wider conduits in sectors B and C of the post-injury rings presumably reflect juvenile tree growth. Helińska-Raczkowska and Fabisiak (1999) determined that, in juvenile wood, earlywood vessel lumen diameter increases with cambial age. Therefore, these wider conduits observed in our young trees possibly attest to the resumption of normal juvenile tree growth.

Abnormally narrow as well as more numerous earlywood vessels developed as a consequence of cambial injury, which is consistent with recent wood anatomical investigations of fire scars (Bigio et al., 2010; Kames et al., 2011). Vessel number increase was also prevalent in sector A of the injury ring, where radial groups of vessels were clearly visible (Fig. 4a). The presence of more vessels, however, was noted over at least three years after stem scarring. Several authors have stated that fewer to no vessels are formed following wounding when actually only describing the barrier zone, i.e. the callus tissue initially built at the wound surface (Sharon, 1973; Moore, 1978; Mulhern et al., 1979; Rademacher et al., 1984; Lowerts et al., 1986; Stobbe et al., 2002; Schwarze et al., 2007).

The initiation of more rays was also noted over at least three years after stem scarring. Ray number was larger in



**Fig. 4.** Microscopic views of *F. excelsior* wood anatomy prior to and after cambial injury. The dashed line shows the position of the cambium at the time of wounding. (a) Xylem cells from the control ring Cr 2 and from the injury ring (Ir 1) as seen close to the wound margin (sector A). In response to mechanical damage, more and narrower earlywood vessels were formed during the growing season following wounding. They accumulated in radial groups. (b) Wound xylem also differed from regular xylem through the enlargement of pre-existing rays.

the post-injury rings, increasing toward the wound margin. It should be emphasized, however, that most of the rays originated in the late portion of the injury ring. Moreover, despite the fact that ray size was not measured in this study, pre-existing rays were noticed to enlarge following wounding (Fig. 4b). These findings further highlight that broad-leaved trees adjust to mechanical damage through ray number increase (Carmi *et al.*, 1972; Rademacher *et al.*, 1984; Lowerts *et al.*, 1986; Lev-Yadun and Aloni, 1992) and ray size increase (Carmi *et al.*, 1972; Sharon, 1973; Mulhern *et al.*, 1979; Bauch *et al.*, 1980; Rademacher *et al.*, 1984; Lev-Yadun and Aloni, 1992, 1993; Lev-Yadun, 1994).

#### *Functional meaning and ecophysiological origin of the anatomical anomalies*

More and narrower earlywood vessels substantially reduced xylem relative conductivity and xylem vulnerability to cavitation in the injury ring, attesting to the functional need of trees to balance xylem hydraulic efficiency against xylem safety from embolism (Zimmermann, 1983; Tyree *et al.*, 1994; Sperry *et al.*, 2008). Moreover, wood composed of narrower vessels is deemed mechanically stronger because of increased cross-sectional area available for fibres, which may be construed as a trade-off between xylem hydraulic efficiency and xylem mechanical strength (Wagner *et al.*, 1998; McCulloh *et al.*, 2004). The diminution of vessel lumen size actually led to a greater thickness-to-span ratio of vessels, which translates into higher wood density, greater mechanical reinforcement, and greater resistance to cavitation (Hacke *et al.*, 2001; Lens *et al.*, 2011). In addition, vessel grouping further increased xylem safety from embolism due to a greater

redundancy of water pathways (Carlquist, 1984; Lens *et al.*, 2011). The ray anomalies (more and larger rays) detected in the callus tissue are also of great adaptive value because rays enhance compartmentalization of decay (Shigo, 1984) and strengthen wood to better adjust to radial mechanical stress (Mattheck and Kubler, 1995; Burgert and Eckstein, 2001). In fact, rays in ring-porous wood reinforce radial wood cohesion acting as radial pins preventing layers with different stiffness (i.e. earlywood and latewood) from slipping past each other (Reiterer *et al.*, 2002).

Stem scarring resulted in the occurrence of strong vessel anomalies in the injury ring, presumably because wounding locally obstructed the basipetal flow of auxin, consequently raising auxin concentration in the injured area and causing rapid vessel differentiation (Aloni and Zimmermann, 1983, 1984; Roberts *et al.*, 1988). Furthermore, ethylene is synthesized following wounding (Imaseki, 1985; Abeles *et al.*, 1992) and is believed to influence, along with radial and axial signal flows, the number and size of regenerative cells (Lev-Yadun and Aloni, 1995). Ethylene applied on the stem of unscarred *Ulmus americana* seedlings already stimulated the development of numerous abnormally narrow earlywood vessels as well as the enlargement of rays (Yamamoto *et al.*, 1987). It is possible that the observed wound-induced anatomical changes in wood structure may also be accounted for by ethylene interfering with the axial flow of auxin (Jaffe, 1980; Abeles *et al.*, 1992).

#### *Impact of wounding on tree metabolism*

As pointed out by Schweingruber (2007), in times of environmental stresses, tree metabolism is subject to economical principles. Tree priorities following wounding include the re-establishment of xylem mechanical strength and xylem safety from embolism, which were found to occur through an increase in the callus mass, but at the expense of water transport and hence future tree growth. The higher proportion of radial parenchyma in the callus tissue corresponds to a considerable effort for defence against pathogens (compartmentalization) and wound healing. Cambial injury stimulates the production of parenchyma around the wound in order to protect the living tissue (Schmitt and Leise, 1990).

The young trees examined demonstrated high resistance to mechanical damage. On the one hand, the negative wound effects due to the reduced xylem hydraulic efficiency lasted only one year and were restrained to a relatively small sector (30°) immediately adjacent to the wound. An equally prompt recovery in vessel lumen size has been observed in *Quercus pyrenaica* and *F. angustifolia* (Ballesteros *et al.*, 2010). On the other hand, the increased investment in fixing carbohydrates seems to last over a longer period of at least three years. It is probable that, similarly to the time required for wound closure, the strength and persistence of wound-induced anatomical anomalies strongly vary according to tree species, tree vigour, and wound size (Neely, 1988; Delvaux *et al.*, 2010).

In conclusion, this study presents wounding as an environmental force moulding the wood structure of broad-leaved

trees and provides detailed information on the duration and extension of associated anatomical changes, thus improving knowledge on temporal (radial) and spatial (tangential) cambial activity in response to mechanical damage.

## Acknowledgements

The authors are indebted to the Service des Forêts et du Paysage (SFP) and the Service des Routes et des Cours d'Eau (SRCE), Canton of Valais, for their financial support of this study as part of the RUFINE project. The authors also acknowledge the valuable suggestions raised by the reviewers and editors.

## References

- Abeles FB, Morgan PW, Saltveit Jr ME.** 1992. *Ethylene in plant biology*. San Diego, USA: Academic Press.
- Aloni R, Zimmermann MH.** 1983. The control of vessel size and density along the plant axis: a new hypothesis. *Differentiation* **24**, 203–208.
- Aloni R, Zimmermann MH.** 1984. Length, width and pattern of regenerative vessels along strips of vascular tissue. *Botanical Gazette* **145**, 50–54.
- Arbellay E, Stoffel M, Bollschweiler M.** 2010. Wood anatomical analysis of *Alnus incana* and *Betula pendula* injured by a debris-flow event. *Tree Physiology* **30**, 1290–1298.
- Asshoff R, Schweingruber FH, Wermelinger B.** 1998–1999. Influence of a gypsy moth (*Lymantria dispar* L.) outbreak on radial growth and wood anatomy of Spanish chestnut (*Castanea sativa* Mill.) in Ticino (Switzerland). *Dendrochronologia* **16–17**, 133–145.
- Astrade L, Bégin Y.** 1997. Tree-ring response of *Populus tremula* L. and *Quercus robur* L. to recent spring floods of the Saône River, France. *Ecoscience* **4**, 232–239.
- Ballesteros JA, Stoffel M, Bollschweiler M, Bodoque JM, Díez-Herrero A.** 2010. Flash-flood impacts cause changes in wood anatomy of *Alnus glutinosa*, *Fraxinus angustifolia* and *Quercus pyrenaica*. *Tree Physiology* **30**, 773–781.
- Bauch J, Shigo AL, Starck M.** 1980. Wound effects in the xylem of *Acer* and *Betula* species. *Holzforchung* **34**, 153–160.
- Bigio E, Gärtner H, Conedera M.** 2010. Fire-related features of wood anatomy in a sweet chestnut (*Castanea sativa*) coppice in southern Switzerland. *Trees* **24**, 643–655.
- Blanchette RA.** 1992. Anatomical responses of xylem to injury and invasion by fungi. In: Blanchette RA, Biggs AR, eds. *Defense mechanisms of woody plants against fungi*. Berlin, Germany: Springer, 76–95.
- Burgert I, Eckstein D.** 2001. The tensile strength of isolated wood rays of beech (*Fagus sylvatica* L.) and its significance for the biomechanics of living trees. *Trees* **15**, 168–170.
- Carlquist S.** 1977. Ecological factors in wood evolution: a floristic approach. *American Journal of Botany* **64**, 887–896.
- Carlquist S.** 1984. Vessel grouping in dicotyledon wood: significance and relationship to imperforate tracheary elements. *Aliso* **10**, 505–525.
- Carlquist S.** 2001. *Comparative wood anatomy: systematic, ecological, and evolutionary aspects of dicotyledon wood*. Berlin, Germany: Springer.
- Carmi A, Sachs T, Fahn A.** 1972. The relation of ray spacing to cambial growth. *New Phytologist* **71**, 349–353.
- Corcuera L, Camarero JJ, Gil-Pelegrín E.** 2004. Effects of a severe drought on *Quercus ilex* radial growth and xylem anatomy. *Trees* **16**, 83–92.
- Delvaux C, Sinsin B, Van Damme P, Beeckman H.** 2010. Wound reaction after bark harvesting: microscopic and macroscopic phenomena in ten medicinal tree species (Benin). *Trees* **24**, 941–951.
- Denn MP, Dodd RS.** 1981. The environmental control of xylem differentiation. In: Barnett JR, ed. *Xylem cell development*. Kent, UK: Castle House, 236–255.
- Eilmann B, Weber P, Rigling A, Eckstein D.** 2006. Growth reactions of *Pinus sylvestris* L. and *Quercus pubescens* Willd. to drought years at a xeric site in Valais, Switzerland. *Dendrochronologia* **23**, 121–132.
- Fink S.** 1999. *Pathological and regenerative plant anatomy. Encyclopedia of plant anatomy*, Vol. 14, Part 6. Berlin, Germany: Gebrüder Bornträger.
- Fonti P, García-González I.** 2004. Suitability of chestnut earlywood vessel chronologies for ecological studies. *New Phytologist* **163**, 77–86.
- Fonti P, Solomonoff N, García-González I.** 2007. Earlywood vessels of *Castanea sativa* record temperature before their formation. *New Phytologist* **173**, 562–570.
- Fonti P, von Arx G, García-González I, Eilmann B, Sass-Klaassen U, Gärtner H, Eckstein D.** 2010. Studying global change through investigation of the plastic responses of xylem anatomy in tree rings. *New Phytologist* **185**, 42–53.
- Galle A, Esper J, Feller U, Ribas-Carbo M, Fonti P.** 2010. Response of wood anatomy and carbon isotope composition of *Quercus pubescens* saplings subjected to two consecutive years of summer drought. *Annals of Forest Science* **67**, 809.
- García-González I, Eckstein D.** 2003. Climatic signal of earlywood vessels of oak on a maritime site. *Tree Physiology* **23**, 497–504.
- Hacke UG, Sperry JS, Pockman WT, Davis SD, McCulloh KA.** 2001. Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure. *Oecologia* **126**, 457–461.
- Helińska-Raczkowska L, Fabisiak E.** 1999. Radial variation of earlywood vessel lumen diameter as an indicator of the juvenile growth period in ash (*Fraxinus excelsior* L.). *Holz als Roh- und Werkstoff* **57**, 283–286.
- Huber F.** 1993. Déterminisme de la surface des vaisseaux du bois des chênes indigènes (*Quercus robur* L., *Quercus petraea* Liebl.): effet individuel, effet de l'appareil foliaire, des conditions climatiques et de l'âge de l'arbre. *Annals of Forest Science* **50**, 509–524.
- Imaseki H.** 1985. Hormonal control of wound-induced responses. In: Pharis RP, Reid DM, eds. *Encyclopedia of plant physiology*, New Series, Vol. 11, Part 3. Berlin, Germany: Springer, 485–512.
- Jaffe MJ.** 1980. Morphogenetic responses of plants to mechanical stimuli or stress. *BioScience* **30**, 239–243.

- Kames S, Tardif JC, Bergeron Y.** 2011. Anomalous earlywood vessel lumen area in black ash (*Fraxinus nigra* Marsh.) tree rings as a potential indicator of forest fires. *Dendrochronologia* **29**, 109–114.
- Kozłowski TT, Kramer PJ, Pallardy SG.** 1991. *The physiological ecology of woody plants*. San Diego, USA: Academic Press.
- Kuroda K, Shimaji K.** 1985. Wound effects on cytodifferentiation in hardwood xylem. *IAWA Bulletin* **6**, 107–118.
- Larson PR.** 1994. *The vascular cambium. Development and structure*. Berlin, Germany: Springer.
- Lens F, Sperry JS, Christman MA, Choat B, Rabaey D, Jansen S.** 2011. Testing hypotheses that link wood anatomy to cavitation resistance and hydraulic conductivity in the genus *Acer*. *New Phytologist* **190**, 709–723.
- Lev-Yadun S.** 1994. Experimental evidence for the autonomy of ray differentiation in *Ficus sycomorus* L. *New Phytologist* **126**, 499–504.
- Lev-Yadun S, Aloni R.** 1992. The role of wounding and partial girdling in differentiation of vascular rays. *International Journal of Plant Sciences* **153**, 348–357.
- Lev-Yadun S, Aloni R.** 1993. Effect of wounding on the relations between vascular rays and vessels in *Melia azedarach* L. *New Phytologist* **124**, 339–344.
- Lev-Yadun S, Aloni R.** 1995. Differentiation of the ray system in woody plants. *Botanical Review* **61**, 45–84.
- Lowerts G, Wheeler EA, Kellison RC.** 1986. Characteristics of wound-associated wood of yellow poplar (*Liriodendron tulipifera* L.). *Wood and Fiber Science* **18**, 537–552.
- Mattheck C, Kubler H.** 1995. *Wood: the internal optimization of trees*. Berlin, Germany: Springer.
- McCulloh KA, Sperry JS, Adler FR.** 2004. Murray's law and the hydraulic vs mechanical functioning of wood. *Functional Ecology* **18**, 931–938.
- Moore KE.** 1978. Barrier zone formation in wounded stems of sweetgum. *Canadian Journal of Forest Research* **8**, 389–397.
- Mulhern J, Shortle WC, Shigo AL.** 1979. Barrier zones in red maple: an optical and scanning microscope examination. *Forest Science* **25**, 311–316.
- Neely D.** 1988. Tree wound closure. *Journal of Arboriculture* **14**, 148–152.
- Pumijumngong N, Park WK.** 1999. Vessel chronologies from teak in northern Thailand and their climatic signal. *IAWA Journal* **20**, 285–294.
- Rademacher P, Bauch J, Shigo AL.** 1984. Characteristics of xylem formed after wounding in *Acer*, *Betula* and *Fagus*. *IAWA Bulletin* **5**, 141–151.
- Régent Instruments Inc.** 2004. WinCELL Pro V 2004a. [www.regentinstruments.com](http://www.regentinstruments.com).
- Reiterer A, Burgert I, Sinn G, Tschegg S.** 2002. The radial reinforcement of the wood structure and its implication on mechanical and fracture mechanical properties: a comparison between two tree species. *Journal of Materials Science* **37**, 935–940.
- Rier JP, Shigo AL.** 1972. Some changes in red maple, *Acer rubrum*, tissues within 34 days after wounding in July. *Canadian Journal of Botany* **50**, 1783–1784.
- Roberts LW, Gahan PB, Aloni R.** 1988. *Vascular differentiation and plant growth regulators*. Berlin, Germany: Springer.
- Sass U, Eckstein D.** 1995. The variability of vessel size of beech (*Fagus sylvatica* L.) and its ecophysiological interpretation. *Trees* **9**, 247–252.
- Schmitt U, Liese W.** 1990. Wound reaction of the parenchyma in *Betula*. *IAWA Bulletin* **11**, 413–420.
- Schwarze F, Grüner J, Schubert M, Fink S.** 2007. Defence reactions and fungal colonisation in *Fraxinus excelsior* and *Tilia platyphyllos* after stem wounding. *Arboricultural Journal* **30**, 1–22.
- Schweingruber FH.** 2007. *Wood structure and environment*. Berlin, Germany: Springer.
- Sharon EM.** 1973. Some histological features of *Acer saccharum* wood formed after wounding. *Canadian Journal of Forest Research* **3**, 83–89.
- Shigo AL.** 1984. Compartmentalization: a conceptual framework for understanding how trees grow and defend themselves. *Annual Review of Phytopathology* **22**, 189–214.
- Sperry JS, Meinzer FC, McCulloh KA.** 2008. Safety and efficiency conflicts in hydraulic architecture: scaling from tissues to trees. *Plant, Cell and Environment* **31**, 632–645.
- St George S, Nielsen E.** 2003. Palaeoflood records for the Red River, Manitoba, Canada, derived from anatomical tree-ring signatures. *Holocene* **13**, 547–555.
- Stobbe H, Schmitt U, Eckstein D, Dujesiefken D.** 2002. Developmental stages and fine structure of surface callus formed after debarking of living lime trees (*Tilia* sp.). *Annals of Botany* **89**, 773–782.
- Tardif JC, Conciatori F.** 2006. Influence of climate on tree rings and vessel features in red oak and white oak growing near their northern distribution limit, southwestern Quebec, Canada. *Canadian Journal of Forest Research* **36**, 2317–2330.
- Tyree MT, Davis SD, Cochard H.** 1994. Biophysical perspectives of xylem evolution: is there a tradeoff of hydraulic efficiency for vulnerability to dysfunction? *IAWA Journal* **15**, 335–360.
- Van den Oever L, Baas P, Zandee M.** 1981. Comparative wood anatomy of *Symplocos* and latitude and altitude of provenance. *IAWA Bulletin* **2**, 3–24.
- Wagner KR, Ewers FW, Davis SD.** 1998. Tradeoffs between hydraulic efficiency and mechanical strength in the stems of four co-occurring species of chaparral shrubs. *Oecologia* **117**, 53–62.
- White FM.** 2006. *Viscous fluid flow*. New York, USA: McGraw-Hill Book Company.
- Woodcock DW.** 1989. Climate sensitivity of wood anatomical features in a ring-porous oak (*Quercus macrocarpa*). *Canadian Journal of Forest Research* **19**, 639–644.
- Yamamoto F, Angeles G, Kozłowski TT.** 1987. Effect of ethrel on stem anatomy of *Ulmus americana* seedlings. *IAWA Bulletin* **8**, 3–10.
- Yanosky TM.** 1983. Evidence of floods on the Potomac River from anatomical abnormalities in the wood of flood-plain trees. *US Geological Survey Professional Paper* **1296**, 1–42.
- Zimmermann MH.** 1983. *Xylem structure and the ascent of sap*. Berlin, Germany: Springer.