

Cellular Calcium Regulation in Hypertension

K. Hermsmeyer and P. Erne

In vascular muscle cells, two distinct types of functionally important calcium (Ca^{2+}) channels, called transient (T) and sustained (L), are differentiated by dihydropyridine calcium antagonists (CaA). We studied the ratio of T/L Ca^{2+} channels in isolated, spontaneously contracting azygous venous cells of spontaneously hypertensive rats (SHR) and Wistar-Kyoto rats (WKY) by quantitating Ca^{2+} currents and intracellular Ca^{2+} release. While total transmembranous Ca^{2+} current was not different between the two strains, the proportion of Ca^{2+} currents carried by L-type channels was enhanced in vascular muscle cells from SHR. We have recently compared subcellular distribution of intracellular free Ca^{2+} concentration in the same cells, at rest and during stimulation, by quantitation

with a digital photon-counting camera. Fura-2 fluorescence intensity showed that Ca^{2+} release was principally from sarcoplasmic reticulum and that cells from SHR had higher levels of Ca^{2+} upon calcium channel stimulation, especially at the cell periphery. These findings suggest fundamental differences in SHR and WKY vascular muscle cells implicating the importance of changes in calcium channels, modulation of Ca^{2+} release, and Ca^{2+} uptake in SHR hypertension. *Am J Hypertens* 1989; 2:655-658

KEY WORDS: Vascular muscle, calcium channels, calcium concentration, fura-2, calcium release, SHR, WKY.

Hypertension in its established phase is hemodynamically characterized by an increase in systemic vascular resistance.¹⁻³ This derangement is due to structural differences between normotensive and hypertensive subjects,⁴ but also entails functional components.⁵

As the force-generating bridging between actin and myosin depends on changes of intracellular calcium activity (calcium free-ion concentration) and hence transmembranous calcium movements, modulations of transmembranous calcium influx and intracellular cal-

cium activity, $(\text{Ca}^{2+})_i$, have a pivotal role in determining vasoconstriction. Therefore, mechanisms that lead to an increase in $(\text{Ca}^{2+})_i$ through these processes cause vasoconstriction.

Several lines of evidence suggest that abnormalities of calcium metabolism are involved in the pathogenesis of established human hypertension. In particular, low serum concentrations of ionized Ca^{2+} have been observed in at least a fraction of patients with this disease.^{6,7} Furthermore, there is an enhanced vasodilation^{8,9} and antihypertensive effect^{10,11} of calcium antagonists in patients with essential hypertension.

Studies using the platelet as a model of contractile systems indicate that alterations of Ca^{2+} handling in human hypertensives may occur at the cellular level.^{12,13} However, these studies do not determine whether similar alterations are operative in vascular muscle cells and, if so, whether they are causally implicated or a consequence of elevated blood pressure.

To define vascular muscle mechanisms which are altered in an animal model of hypertension prior to the development of elevated blood pressure, studies were

From the Chiles Research Institute, Providence Medical Center and Departments of Medicine and Cell Biology, Oregon Health Science University, Portland, Oregon, and Department of Pharmacology and The Cardiovascular Center, University of Iowa, Iowa City, Iowa.

This research was supported by grants nos. HL38537 and HL38645 from the National Institutes of Health. P. E. was supported by the Swiss Foundation for Biological and Medical Fellowships.

Address correspondence and reprint requests to Kent Hermsmeyer, M.D., Chiles Research Institute, Providence Medical Center, 4805 NE Glisan Street, Portland, Oregon 97213. Present address for P. E., Departments of Medicine and Research, University Hospital, 4031 Basel, Switzerland.

carried out on isolated azygous vein cells from three-day-old spontaneously hypertensive rats (SHR) and genetically matched Wistar-Kyoto rats (WKY). The preparation and characterization of these cells from primary cultures were described in detail elsewhere.¹⁴ These cells are spontaneously contracting and exhibit repeated contraction/relaxation cycles with a high shortening velocity, high pharmacological sensitivity, and electrophysiological integrity.^{15,16}

METHODS

Experiments were carried out on vascular muscle cells isolated from azygous veins of neonatal rats.^{17,18} Whole cell recordings of Ca^{2+} currents, isolated by replacement of Na^+ and K^+ in internal and external solutions allowed separation of L and T types of current, where L are longer lasting (sustained) and T are transient.¹⁹

In cells from similarly prepared cultures, intracellular Ca^{2+} was quantitated by fura-2 fluorescence using the three wavelength protocols explained in detail elsewhere.²⁰ High resolution ($0.5 \mu\text{m}$) quantitation of Ca^{2+} was carried out with a two-stage microchannel plate camera (Hamamatsu VIM). Excitation of fluorescence at 340, 360, and 380 nm with emission at 510 nm (three wavelength analysis) provides point-by-point quantitation of intracellular free Ca^{2+} activity at each pixel.²⁰

RESULTS

Altered Calcium Currents in Vascular Muscle Cells of SHR

The rationale to study Ca^{2+} currents is based on the observation that alterations in Ca^{2+} entry or exit might well lead to changes in tension without intervention by other ions.²¹ Calcium which triggers excitation-contraction coupling presumably enters vascular muscle cells through voltage-dependent Ca^{2+} channels. These precise Ca^{2+} currents were recorded using tight-seal pipettes by patch-clamp techniques.^{17,18,22} These techniques allowed the identification and characterization of two types of Ca^{2+} channel currents. The low threshold, transient channel (T) is activated with small depolarizations and inactivates rapidly. The high threshold, long-lasting channel (L, sustained) is activated at more positive membrane potentials, inactivates slowly, and is sensitive to dihydropyridine calcium antagonists.¹⁸ Based on these characteristics, it has been suggested that the T-channel participates in the generation of Ca^{2+} -dependent electrical spiking and spontaneous activity in vascular muscle cells, whereas the L-channel is involved in the triggering and maintenance of vascular muscle contraction.

Peak amplitudes of T, L, and total (sum of T and L) Ca^{2+} currents were measured in 30 cells from WKY and 30 cells from SHR.¹⁹ While no difference in total peak amplitude of Ca^{2+} current was observed in these cells from neonatal rats, Ca^{2+} current carried by L (sustained) channels contributed significantly more ($62 \pm 5\%$) to




Figure 1. Ratios of sustained (L) to transient (T) calcium current were greater in SHR than WKY isolated vascular muscle cells measured under whole cell voltage-clamp conditions. Each dot represents one cell from which the ratios were measured in 30 SHR and 30 WKY, with means indicated by solid lines (significantly different) and means by dotted horizontal lines.¹⁹

total peak current in SHR than in WKY ($42 \pm 7\%$) based on 60 experiments ($P < 0.05$). When ratios (L/T) of Ca^{2+} current were calculated, the mean value for 30 WKY was 0.9 and 2.9 for 30 SHR cells (Figure 1). These results demonstrate that even at the prehypertensive stage, L channels are more prominent in SHR than WKY, suggesting a genetic membrane channel defect with the proper timing to be a cause of increased Ca^{2+} in vascular muscle cells.

Elevated Free Calcium Concentrations at Subsarcolemmal Sites

To study if these Ca^{2+} currents associated with L channels are linked to altered intracellular Ca^{2+} handling, the distribution of intracellular free Ca^{2+} activities, $(\text{Ca}^{2+})_i$, was measured in isolated vascular muscle cells with fura-2. Incorporation of the fluorescent dye, quantitation of fluorescent intensities, and $(\text{Ca}^{2+})_i$ determination after correction for inhomogeneous dye distribution are described in detail elsewhere.^{20,23}

In a study quantitating the distribution of intracellular free Ca^{2+} in neonatal vascular muscle cells from SHR and WKY,²⁴ we have measured the average intracellular dye concentration, and quantitated $(\text{Ca}^{2+})_i$ of the whole cytoplasm, as well as in defined regions. We divided the cell into a peripheral boundary, $0.5 \mu\text{m}$ distant from the

cell edges, and central areas, defined as all that remains. Quantitation of Ca^{2+} was according to the following equations which were applied to each of the 250,000 image elements

$$(\text{Ca}^{2+})_i = K_d (cR - 1.36)/(3.48 - cR),$$

where K_d is the dissociation constant established by calibration procedures (239 nM at 37°C), R is the ratio of 510 nm fluorescence at 340/380 nm wavelength excitation, and c corrects for the inhomogeneous dye distribution according to

$$c = 0.92 + 2.24ae^{-0.0028dF},$$

where F is the fura-2 fluorescence intensity count at 360 nm excitation wavelength, a corrects for day-to-day variability of the optical system as calibrated by phosphor beads, and d adjusts for accumulation times. We have performed these measurements in cells perfused with isotonic medium and following administration of 100 mM potassium and 1 μM SDZ 202-791 (+S), a stereoselective dihydropyridine with calcium entry stimulating properties.

Intracellular fura-2 concentrations were similar in cells of both SHR and WKY (62 and 85 μM , respectively, not significantly different). $(\text{Ca}^{2+})_i$ in central regions of nonstimulated cells perfused with isotonic medium was comparable in both strains (WKY: 119 v SHR: 126 nM). But $(\text{Ca}^{2+})_i$ in the peripheral boundary was slightly elevated in SHR (WKY: 100 v SHR: 130 nM, $P < 0.05$). Upon stimulation of the cells by 100 mM potassium and SDZ 202-971, the average myoplasmic Ca^{2+} activity in central regions of SHR tended to increase to a greater extent (WKY: 442 v SHR: 485 nM) accompanied by a more pronounced elevation of $(\text{Ca}^{2+})_i$ in the peripheral boundary (WKY: 539 v SHR: 866 nM, $P < 0.05$). These results point to the important role intracellular Ca^{2+} release from subsarcolemmal Ca^{2+} pools may play in vascular muscle during the prehypertensive stage, suggesting a defect of SHR Ca^{2+} uptake or release mechanisms in these cells.

DISCUSSION/CONCLUSION

These studies were directed towards the characterization of cellular Ca^{2+} handling abnormalities in SHR at a prehypertensive age. The results demonstrate that before development of elevated blood pressure in SHR, the sustained L Ca^{2+} channel is more dominantly expressed,¹⁹ and that Ca^{2+} release or uptake mechanisms from subsarcolemmal stores are already altered. The underlying factors leading to changes in the membrane structure and intracellular organelles could well be linked. For example, Ca^{2+} entering through L channels may have greater effects on the cytoplasmic sites which release intracellular Ca^{2+} . Or, Ca^{2+} entry through L channels may have smaller effects on Ca^{2+} uptake, as suggested by Table 1. Whichever is true, these results

TABLE 1. Ca^{2+} ACTIVITY (FREE ION CONCENTRATION) IN AZYGIOUS VEIN VASCULAR MUSCLE CELLS OF SHR v WKY (nmol/L)

	WKY	SHR
Peripheral rim		
Resting	100 \pm 2	130 \pm 2*
Stimulated	530 \pm 11	882 \pm 33*
Central region		
Resting	119 \pm 2	126 \pm 5
Stimulated	442 \pm 36	485 \pm 31

Resting condition was contractile cells between contractions in control solution. Stimulated condition was cells in 100 mM K^+ with 1 μM Ca^{2+} agonist (SDZ 202-791S). The peripheral rim is defined as the 0.5 μm from the cell edge. Central region was all parts of the cell at least 1 μm from the cell edge, and actually includes the top and bottom peripheral edge contributions to that fluorescence (ie, optical sectioning of fluorescence was not practical in these thin cells).

Values are means \pm SEM; n = 3 for WKY and 5 for SHR.

* $P < 0.05$.

underline the importance of Ca^{2+} regulation of submembranous Ca^{2+} stores.

That total myoplasmic Ca^{2+} activity in unstimulated cells from both strains is comparable tallies with the finding that elevated $(\text{Ca}^{2+})_i$ in platelets can only be observed in older SHR with elevated blood pressure.²⁵ Whether a similar elevation of peripheral myoplasmic Ca^{2+} activity in vascular cells from adult SHR (with elevated blood pressure) develops is unknown. However, it could be postulated that the enhanced responsiveness of SHR to stimuli giving rise to the observed Ca^{2+} increase could lead to a profound resetting of Ca^{2+} uptake and extrusion mechanisms over time. To further characterize the responsible mechanisms, more studies of SHR are needed to investigate these processes in different ages and blood vessels.

REFERENCES

- Cohn JN: Calcium, vascular smooth muscle, and calcium entry blockers in hypertension. *Ann Intern Med* 1983;98:806–809.
- Bohr DF, Webb RC: Vascular smooth muscle function and its changes in hypertension. *Am J Med* 1984;77 (suppl 4A):3–16.
- Hermesmeier RK: Vascular muscle membrane cation mechanisms and total peripheral resistance. *Hypertension* 1987;10 (suppl I):20–22.
- Folkow B: The hemodynamic consequences of adaptive structural changes of resistance vessels in hypertension. *Clin Sci* 1971;41:1–12.
- Bolli P, Erne P, Hulthen UL, et al: Parallel reduction of calcium-influx-dependent vasoconstriction and platelet free calcium concentration with calcium entry and β -adrenoceptor blockade. *J Cardiovasc Pharmacol* 1984;6 (suppl 7):996–1001.
- McCarron DA: Low serum concentrations of ionized cal-

- cium in patients with hypertension. *N Engl J Med* 1982;307:226–228.
7. Resnick LM, Laragh JH, Sealey JE, Alderman MA: Divalent cations in essential hypertension. Relations between serum ionized calcium, magnesium and plasma renin activity *N Engl J Med* 1984;309:888–891.
 8. Hulthen UL, Bolli P, Erne P, et al: Peripheral vasodilating effect of nitrendipine in man, *in* Scriabine A, Vanov S, Deck K (eds): Nitrendipine. Baltimore, Urban and Schwarzenberg, 1984, pp 463–468.
 9. Kiowski W, Bolli P, Erne P, et al: Mechanisms of action of calcium antagonists in hypertension. *J Cardiovasc Pharmacol* 1987;10 (suppl 10):23–27.
 10. Erne P, Bolli P, Bertel O, et al: Factors influencing the hypotensive effects of calcium antagonists. *Hypertension* 1983;5 (suppl II):97–102.
 11. Erne P, Conen D, Kiowski W, et al: Calcium antagonist induced vasodilation in peripheral, coronary and cerebral vasculature as important factors in the treatment of elderly hypertensives. *Eur Heart J* 1987;8 (suppl K):49–56.
 12. Erne P, Bolli P, Burgisser E, Buhler FR: Correlation of platelet calcium with blood pressure: effect of antihypertensive therapy. *N Engl J Med* 1984;310:1084–1088.
 13. Resink TJ, Tkachuk VA, Erne P, Buhler FR: Platelet membrane calmodulin-stimulated calcium-adenosine triphosphatase: altered activity in essential hypertension. *Hypertension* 1986;8:159–166.
 14. Marvin WJ, Robinson RB, Hermsmeyer K: Correlation of function and morphology of neonatal rat and embryonic chick cultured cardiac and vascular muscle cells. *Circ Res* 1979;45:528–540.
 15. Hermsmeyer K, Mason R: Norepinephrine sensitivity and desensitization of cultured single vascular muscle cells. *Circ Res* 1982;50:627–632.
 16. Hermsmeyer K: Excitation of vascular muscle by norepinephrine. *Ann Biomed Eng* 1983;11:567–577.
 17. Sturek M, Hermsmeyer K: Calcium and sodium channels in spontaneously contracting vascular muscle cells. *Science* 1986;233:475–478.
 18. Bean BP, Sturek M, Puga A, Hermsmeyer K: Calcium channels in vascular muscle cells: modulation by dihydropyridine drugs. *Circ Res* 1986;59:229–235.
 19. Rusch N, Hermsmeyer K: Calcium currents are altered in the vascular muscle cell membranes of spontaneously hypertensive rats. *Circ Res* 1988;63:997–1002.
 20. Erne P, Hermsmeyer K: Intracellular Ca^{2+} release in vascular muscle cells by caffeine, ryanodine, norepinephrine, and neuropeptide Y. *J Cardiovasc Pharmacol* 1988;12(suppl 5):S85–S91.
 21. Hermsmeyer K: Might nitrendipine enhance Ca^{2+} transport in vascular muscle? *in* Merrill GF, Weiss HR (eds): Ca^{2+} -Entry Blockers, Adenosine and Neurohumors. Baltimore, Urban and Schwarzenberg, 1983, pp 51–61.
 22. Hermsmeyer K, Rusch N: Felodipine actions on vascular muscle Ca^{2+} channels. *J Cardiovasc Pharmacol* 1987;10 (suppl 1) 40–43.
 23. Erne P, Hermsmeyer K: Desensitization of norepinephrine includes refractoriness of calcium release in myocardial cells. *Biochem Biophys Res Commun* 1988;151:333–338.
 24. Erne P, Hermsmeyer K: Intracellular vascular muscle Ca^{2+} modulation in SHR/WKY hypertension. *Hypertension* 1989;14 (in press).
 25. Bruschi G, Bruschi ME, Caroppo M, et al: Cytoplasmic free Ca^{2+} is increased in the platelets of spontaneously hypertensive rats and essential hypertensive patients. *Clin Sci* 1985;66:179–184.