

# Why $z > 1$ radio-loud galaxies are commonly located in protoclusters

N. A. Hatch,<sup>1</sup>★ D. Wylezalek,<sup>2</sup> J. D. Kurk,<sup>3</sup> D. Stern,<sup>4</sup> C. De Breuck,<sup>2</sup> M. J. Jarvis,<sup>5,6</sup>  
A. Galametz,<sup>3,7</sup> A. H. Gonzalez,<sup>8</sup> W. G. Hartley,<sup>9</sup> A. Mortlock,<sup>10</sup> N. Seymour<sup>11,12</sup>  
and J. A. Stevens<sup>13</sup>

<sup>1</sup>*School of Physics and Astronomy, University of Nottingham, University Park, Nottingham NG7 2RD, UK*

<sup>2</sup>*European Southern Observatory, Karl-Schwarzschild-Str. 2, D-85748 Garching, Germany*

<sup>3</sup>*Max-Planck-Institut fuer Extraterrestrische Physik, Giessenbachstrasse, D-85748 Garching, Germany*

<sup>4</sup>*Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Dr., Pasadena, CA 91109, USA*

<sup>5</sup>*Astrophysics, Department of Physics, Keble Road, Oxford OX1 3RH, UK*

<sup>6</sup>*Physics Department, University of the Western Cape, Bellville 7535, South Africa*

<sup>7</sup>*INAF – Osservatorio di Roma, Via Frascati 33, I-00040 Monteporzio, Italy*

<sup>8</sup>*Department of Astronomy, University of Florida, Gainesville, FL 32611, USA*

<sup>9</sup>*ETH Zürich, Institut für Astronomie, HIT J 11.3, Wolfgang-Pauli-Str. 27, CH-8093 Zürich, Switzerland*

<sup>10</sup>*Institute for Astronomy, University of Edinburgh, Royal Observatory, Edinburgh EH9 3HJ, UK*

<sup>11</sup>*CASS, PO Box 76, Epping, NSW 1710, Australia*

<sup>12</sup>*Curtin Institute of Radio Astronomy, Curtin University, GPO Box U1987, Perth WA 6845, Australia*

<sup>13</sup>*Centre for Astrophysics Research, STRI, University of Hertfordshire, Hatfield AL10 9AB, UK*

Accepted 2014 August 20. Received 2014 August 18; in original form 2014 April 21

## ABSTRACT

Distant powerful radio-loud active galactic nuclei (RLAGN) tend to reside in dense environments and are commonly found in protoclusters at  $z > 1.3$ . We examine whether this occurs because RLAGN are hosted by massive galaxies, which preferentially reside in rich environments. We compare the environments of powerful RLAGN at  $1.3 < z < 3.2$  from the Clusters Around Radio-Loud AGN survey to a sample of radio-quiet galaxies matched in mass and redshift. We find that the environments of RLAGN are significantly denser than those of radio-quiet galaxies, implying that not more than 50 per cent of massive galaxies in this epoch can host powerful radio-loud jets. This is not an observational selection effect as we find no evidence to suggest that it is easier to observe the radio emission when the galaxy resides in a dense environment. We therefore suggest that the dense Mpc-scale environment fosters the *formation* of a radio jet from an AGN. We show that the number density of potential RLAGN host galaxies is consistent with every  $> 10^{14} M_{\odot}$  cluster having experienced powerful radio-loud feedback of duration  $\sim 60$  Myr during  $1.3 < z < 3.2$ . This feedback could heat the intracluster medium to the extent of 0.5–1 keV per gas particle, which could limit the amount of gas available for further star formation in the protocluster galaxies.

**Key words:** galaxies: active – galaxies: high-redshift.

## 1 INTRODUCTION

Radio-loud active galactic nuclei (RLAGN) are typically located in dense environments (e.g. Yates, Miller & Peacock 1989; Hill & Lilly 1991; Best, Longair & Roettgering 1998; Roche, Eales & Hippelein 1998; Best 2000; Donoso et al. 2010). At  $z \gtrsim 1.5$ , many of these regions are dense enough that they will collapse into clusters

by today, so they are commonly referred to as protoclusters (e.g. Venemans et al. 2007; Hatch et al. 2011a).

The Clusters Around Radio-Loud AGN (CARLA) survey recently showed that approximately half of all powerful RLAGN ( $L_{500\text{MHz}} \geq 10^{27.5} \text{ W Hz}^{-1}$ ) at  $1.3 < z < 3.2$  reside in regions that are denser than average by more than  $2\sigma$ . Many of these dense regions are likely to be protoclusters (Wylezalek et al. 2013, 2014). Low-luminosity RLAGN ( $L_{1.4\text{GHz}} \sim 10^{25.5} \text{ W Hz}^{-1}$ ) also tend to reside in rich groups and clusters. Castignani et al. (2014) showed that  $\sim 70$  per cent of low-luminosity RLAGN at  $1 < z < 2$  are surrounded by Mpc-scale galaxy overdensities. They also demonstrated that

\* E-mail: [nina.hatch@nottingham.ac.uk](mailto:nina.hatch@nottingham.ac.uk)

the simple cluster-detection method used by the CARLA survey (Wylezalek et al. 2013, 2014) missed many of the clusters around the low-luminosity RLAGN, so it is likely that the fraction of powerful RLAGN that reside in protoclusters is higher than 50 per cent. These surveys prove that RLAGN efficiently trace galaxy protoclusters; the goal of this paper is to understand why RLAGN are such good beacons of protoclusters.

At  $z \leq 0.7$ , RLAGN occupy richer environments than similarly massive radio-quiet galaxies (e.g. Kauffmann, Heckman & Best 2008; Ramos Almeida et al. 2013). This implies that the presence of a radio jet depends on environment as well as galaxy mass. However, this may not be true at higher redshifts. Powerful RLAGN are hosted by galaxies with a stellar mass of  $> 10^{10.5} M_{\odot}$  (Seymour et al. 2007), and the bias of such massive galaxies implies that they reside in dark matter haloes of  $10^{12.5} M_{\odot}$  or greater at all redshifts (Hartley et al. 2013). In the local Universe, this means most massive galaxies reside in group environments, but such massive haloes at  $z > 1.5$  typically grow into cluster-mass structures by today (Chiang, Overzier & Gebhardt 2013). Thus, massive galaxies at high redshift are likely to trace the progenitors of rich cluster environments, and the spatial correspondence between RLAGN and protoclusters may simply occur because distant RLAGN are hosted by massive galaxies. In this paper, we compare the environments of  $1.3 < z < 3.2$  RLAGN from the CARLA survey to similarly massive galaxies without radio jets, to determine whether RLAGN reside in protoclusters simply because they are massive galaxies, or if the presence of a radio jet depends on environment at high redshift, as it does in the local Universe.

Another reason why powerful RLAGN may be such good tracers of protoclusters is that the radio emission could be amplified if the relativistic electrons are constricted by the dense ambient gas (Barthel & Arnaud 1996). An enhancement of the radio power would produce a selection bias, meaning it is easier to observe RLAGN when they reside in dense environments. Several studies have investigated whether this selection bias exists by searching for a correlation between radio power and environmental density, but the results are contradictory. For example, Karouzos, Jarvis & Bonfield (2014) found no trend, Donoso et al. (2010) found a negative trend and Falder et al. (2010) found a positive trend between radio power and environmental density. Wylezalek et al. (2013) showed that there is no correlation between the radio power of the CARLA AGN and their environment, suggesting that no selection bias exists at  $z > 1.3$ . Nevertheless, there is so much confusion in the literature on this topic that a more in-depth analysis is warranted. Therefore, we will perform a more detailed comparison of the radio properties of the CARLA AGN with environment to determine if the ambient gas affects the radio emission.

The purpose of this study is to explore why RLAGN are such good beacons for locating protoclusters. We test two hypotheses: (i) that RLAGN are in denser environments simply because they are hosted by massive galaxies and (ii) that the dense protocluster environment amplifies the radio emission causing a selection bias. In Section 2, we introduce our data and describe how we create a radio-quiet control sample that is matched in redshift and mass proxy to the CARLA RLAGN sample. In Section 3, we compare the environments of the RLAGN and radio-quiet massive galaxies, and then search for any correlation between the properties of the radio emission and the surrounding environment. We discuss the implications of our results in Section 4. We use AB magnitudes throughout and a  $\Lambda$  cold dark matter flat cosmology with  $\Omega_M = 0.3$ ,  $\Omega_{\Lambda} = 0.7$  and  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ .

## 2 METHOD

### 2.1 Data

#### 2.1.1 RLAGN sample: CARLA

CARLA is a 400 h Warm *Spitzer Space Telescope* programme designed to investigate the environments of powerful RLAGN. The CARLA sample consists of 419 very powerful RLAGN lying at  $1.3 < z < 3.2$  and having a 500 MHz luminosity  $\geq 10^{27.5} \text{ W Hz}^{-1}$ . The sample comprises 211 radio-loud quasars (RLQs) and 208 radio galaxies.

The radio galaxies at  $z > 2$  were selected from the compendium of Miley & De Breuck (2008); radio galaxies at  $z < 2$  were selected from flux-limited and ultra-steep-spectrum radio surveys to have the same distribution of radio power as the higher redshift sample. The RLQ sample comprises optically bright ( $M_B < -26.5$ ) quasars in SDSS (Schneider et al. 2010) and the 2dF QSO Redshift Survey (Croom et al. 2004) with NRAO VLA Sky Survey (NVSS) detections above the  $L_{500\text{MHz}} \geq 10^{27.5} \text{ W Hz}^{-1}$  threshold. More RLQs than radio galaxies matched these criteria, so we limited the RLQ sample to 211 sources which matched the redshift and radio-power distribution of the radio galaxy sample. Full details of the CARLA RLAGN sample can be found in Wylezalek et al. (2013). No information about the environments was taken into account when selecting targets, so the sample is representative of the entire powerful RLAGN population at these redshifts.

Deep *Spitzer* data covering  $5.2 \text{ arcmin} \times 5.2 \text{ arcmin}$  were obtained on each field with the Infrared Array Camera (IRAC; Fazio et al. 2004) on board the *Spitzer Space Telescope* at 3.6 and 4.5  $\mu\text{m}$  during cycles 7 and 8. The CARLA 95 per cent completeness limiting magnitudes are  $[3.6] = 22.6$  and  $[4.5] = 22.9$  mag. Details of the observations and data reduction can be found in Wylezalek et al. (2013). Using the well-tested *Spitzer* IRAC colour criterion  $[3.6] - [4.5] > -0.1$  to select galaxies at  $z > 1.3$ , Wylezalek et al. (2013) showed that 55 per cent of these RLAGN are surrounded by significant excesses of galaxies that are likely associated with the RLAGN.

#### 2.1.2 Control fields: UDS and SpUDS

To obtain a control field sample of radio-quiet galaxies with the same mass and redshift distribution as the RLAGN, we utilize the UKIDSS Ultra Deep Survey (UDS; Almaini et al., in preparation). The UDS is a deep  $0.8 \text{ deg}^2$  near-infrared (IR) survey overlapping part of the *SubaruXMM-Newton* Deep Survey (SXDS; Furusawa 2008). In the overlapping  $\sim 0.54 \text{ deg}^2$  area, the total comoving volume between  $z = 1.3$  and 3.2 is  $0.012 \text{ Gpc}^{-3}$ . In this volume, we expect approximately 50 protoclusters that will collapse to form clusters of  $M_{200} \geq 10^{14} M_{\odot}$  (assuming local galaxy cluster number counts from Vikhlinin et al. 2009).

A sample of massive galaxies are identified from the UDS using the photometric redshifts and stellar masses derived by Hartley et al. (2013) and Mortlock et al. (2013). These authors combine *U*-band data from the Canada–France–Hawaii Telescope (Foucaud et al., in preparation) with optical photometry from the SXDS, *JHK* photometry from the eighth data release (DR8) of the UDS, and *Spitzer* Ultra Deep Survey data (SpUDS; PI: J. Dunlop) to create a *K*-selected *UBVRizJHK*[3.6][4.5] catalogue.

Photometric redshifts were determined by fitting spectral energy distribution (SED) templates to the photometric data points.

Following Hartley et al. (2013), we remove objects with poorly determined photometric redshifts by removing the objects whose minimum  $\chi^2$  is greater than 11.35 from the photometric redshift-fitting procedure (15 per cent of the sample). Many of the removed objects are blended sources or optically bright AGN, so the control sample may be biased against galaxies containing optically bright AGN. These objects have average environments, so removing them from the UDS catalogues does not bias our results. The dispersion of the remaining photometric redshifts is  $\Delta z/(1+z) = 0.031$ . For full details regarding the methodology and resulting photometric redshifts of the UDS catalogue, see Hartley et al. (2013).

The stellar masses of the  $K$ -selected UDS galaxies were measured by fitting the photometry to a large grid of synthetic SEDs from Bruzual & Charlot (2003) stellar population models (created using a Chabrier initial mass function; Chabrier 2003). Full details of the fitting procedure are given in Mortlock et al. (2013). The catalogue is 95 per cent complete to  $\log(M/M_\odot) = 10.3$  at the highest redshift of the CARLA sample ( $z = 3.2$ ).

The UDS galaxies were classified as star forming or quiescent using the two-colour rest-frame  $UVJ$  selection introduced by Wuyts et al. (2007). Our division follows the boundaries defined by Williams et al. (2009) and extends their  $1 < z < 2$  values to higher redshift. However, quiescent galaxies selected by this method can be contaminated by AGN and dusty star-forming objects. Therefore, we reclassify galaxies as star forming if they are fitted by a template with specific star formation rate (sSFR)  $> 10^{-8} \text{ yr}^{-1}$ , or if the galaxy is associated with a *Spitzer* 24  $\mu\text{m}$  source, with a flux that would imply an sSFR  $> 7.43^{-11} \text{ yr}^{-1}$  (a stellar mass doubling time less than the  $z = 0$  Hubble time). As shown by Stern et al. (2006), the 24  $\mu\text{m}$  data reliably break the degeneracy between passive and dusty red galaxies.

The environment of the UDS massive galaxies was measured using SpUDS, a 1 deg<sup>2</sup> cycle 4 *Spitzer* Legacy programme which encompasses the UDS field. We used the SpUDS 3.6 and 4.5  $\mu\text{m}$  catalogues of Wylezalek et al. (2013), which were extracted from the public mosaics in the same way as for CARLA. The catalogues were extracted in dual-image mode with the 4.5  $\mu\text{m}$  image used as the detection image. The SpUDS data reach  $3\sigma$  sensitivities of  $[3.6] \sim [4.5] \sim 24 \text{ mag}$ , but in all following work the catalogues were cut to the shallower depth of the CARLA data.

## 2.2 Control sample

### 2.2.1 Creating a control sample matched in mass and redshift to CARLA RLAGN

In this section, we describe how a radio-quiet control sample of galaxies, with similar masses and redshifts as the CARLA sample, were selected from the UDS catalogues. We first removed all sources within 2 arcmin of bad regions of the SpUDS data, so that we do not include sources whose environments are affected by bright stars or field edges.

To form a radio-quiet sample, we removed all UDS galaxies from the catalogue that were detected in the 100  $\mu\text{Jy}$  *Subaru/XMM-Newton* Deep Field radio source sample of Simpson et al. (2006). There are 109 sources with radio fluxes greater than 100  $\mu\text{Jy}$  in the redshift, [3.6] magnitude and colour range of the CARLA sample, most of which lie at  $z < 1.8$ . Although we refer to the remaining galaxy sample as ‘radio quiet’, we note that the 100  $\mu\text{Jy}$  flux density limit of the Simpson et al. (2006) catalogue means that the sample may still include radio-emitting galaxies with a 500 MHz luminosity of  $10^{24.4} \text{ W Hz}^{-1}$  at  $z = 1.3$  and up to  $10^{25.2} \text{ W Hz}^{-1}$  at  $z = 3.2$

(assuming a spectral index  $\alpha = -0.9$ , see Fig. 5c). These radio luminosities are at least two orders of magnitude lower than the radio luminosities of the CARLA RLAGN.

The next step was to select galaxies which have a similar distribution of stellar mass as the CARLA RLAGN. Radio-loud galaxies are among the most massive galaxies at every redshift; Seymour et al. (2007) found that none of the 70 radio galaxies in the *Spitzer* high-redshift radio galaxy (SHzRG) programme had a stellar mass<sup>1</sup>  $< 10^{10.3} M_\odot$ , so all UDS galaxies with stellar masses  $< 10^{10.3} M_\odot$  were removed from the control sample. We tested the impact of this mass cut by including all control galaxies with  $< 10^{10.3} M_\odot$ , and find that our results do not change.


To derive the stellar masses of the CARLA RLAGN requires multicomponent SED fitting that takes into account light from the AGN as well as the stellar population. To do this requires mid-IR data at wavelengths  $> 5 \mu\text{m}$  to remove the hot dust component (e.g. De Breuck et al. 2010; Drouart et al. 2014), and high-resolution data to remove the direct AGN light (e.g. Hatch et al. 2013). Since these data are not available for the majority of the CARLA sample, we are unable to perform adequate fits to the SEDs to obtain stellar masses.

An alternative to deriving stellar masses is using the *Spitzer* IRAC fluxes and colours as mass proxies. The *Spitzer* [3.6] and [4.5] bands cover rest-frame near-IR emission (0.8–2  $\mu\text{m}$ ) for galaxies at  $1.3 < z < 3.2$ , and hence are relatively good tracers of stellar mass. We cannot use the *Spitzer* IRAC fluxes of the CARLA quasars as mass proxies, as these objects are dominated by the light from the central AGN. However, the IRAC fluxes of the 208 CARLA radio galaxies, whose central AGN light is mostly obscured, are likely to be good mass proxies.

To form the control sample, we divided the 208 radio galaxies in the CARLA sample into 10 redshift bins of width  $\Delta z = 0.19$ ; all CARLA RLAGN have spectroscopically measured redshifts. Each redshift bin was subdivided into twelve [3.6] magnitude bins of  $\Delta m = 0.33$ , which was further subdivided into four bins of [3.6]–[4.5] colour with  $\Delta \text{colour} = 0.3$ . Thus, the 208 CARLA radio galaxies were divided into 480 bins. The radio-quiet UDS galaxies were divided into the same 480 bins, and from each bin we randomly selected twice the number of UDS galaxies as CARLA radio galaxies. 18 bins contained fewer UDS galaxies than CARLA radio galaxies; therefore, we gave an extra weight to the 38 UDS galaxies in these bins so they had the equivalent weight of 114 galaxies. The final control sample consisted of 280 galaxies.

A subset of 68 CARLA radio galaxies fell in bins that were not occupied by any UDS source. These sources were generally bright and red galaxies at high redshift. It is likely that the IRAC fluxes from these galaxies are contaminated by direct AGN light or hot dust. The caveat to using IRAC magnitudes as a mass proxy is that approximately a third of radio galaxies at  $1.3 < z < 3.2$  are dominated, at rest-frame 1.6  $\mu\text{m}$ , by AGN-heated hot dust emission rather than stellar emission (De Breuck et al. 2010). These galaxies appear slightly brighter and redder than galaxies of similar mass without AGN. We thus removed the 68 galaxies from the CARLA radio galaxy sample to leave a subsample of 140 galaxies that matches the control sample perfectly in redshift, [3.6] and [3.6]–[4.5]. Removing these galaxies does not affect our results as the redshifts and environmental densities of the remaining CARLA

<sup>1</sup> The SHzRG galaxy masses were reduced by 13 per cent to account for the differences in assumed initial mass functions in the SED fitting [Kroupa (2001) for SHzRG versus Chabrier (2003) for the UDS].



**Figure 1.** The distributions of the redshifts (left), [3.6] magnitudes (middle) and [3.6]–[4.5] colours (right) of the CARLA reduced radio galaxy sample (red), the UDS control sample (blue) and the whole CARLA sample (black; shown in the left-hand panel only as the IRAC magnitudes and colours of the RLQs are dominated by the quasar light. See Section 2.2.1 for details). The control sample is closely matched to the CARLA radio galaxy sample; KS tests result in probabilities in the range of 0.3–0.97, so there are no significant differences between these distributions.

radio galaxies are very similar to the full CARLA sample (as shown in Figs 1 and 3). It is possible that some RLAGN in our remaining CARLA radio galaxy sample are still contaminated by low levels of hot dust emission in the IRAC bands; however, this means that our mass-matching will be on the conservative side, and the control galaxies selected may be slightly more massive than the CARLA radio galaxies.


In Fig. 1, we compare the redshifts, [3.6] and [3.6]–[4.5] colours of the CARLA radio galaxy subsample and the control sample. Kolmogorov–Smirnov (KS) tests result in probabilities of 0.77, 0.97 and 0.33, for the redshifts, magnitudes and colours, respectively. These tests suggest that there are no significant differences in these properties between the CARLA radio galaxy subsample and UDS control galaxies.

The fraction of control galaxies that host radio-quiet AGN was measured using deep *XMM–Newton* data (Ueda et al. 2008). The closest *K*-band selected galaxy within 5 arcsec of the X-ray point source was assumed to be the galaxy counterpart, which resulted in 191 X-ray point sources in the redshift range  $1.3 < z < 3.2$ . Approximately 7 per cent of the control galaxies have X-ray detections indicating that they host radio-quiet AGN. There is no significant difference in the IRAC luminosity, colour or environment between the X-ray-bright control galaxies and the rest of the control sample.

## 2.2.2 Stellar masses of the CARLA and control galaxies

One of the aims of this work is to test whether RLAGN reside in denser environments than similarly massive radio-quiet galaxies. Hence, the validity of our results hinges on our selection of a control sample that is well matched in both redshift and mass. The redshift selection is trivial since the CARLA RLAGN all have spectroscopic redshifts and the photometric redshifts of the UDS catalogue are of high quality (Hartley et al. 2013). Ensuring that our control sample matches the stellar masses of the CARLA RLAGN is harder because we must rely on mass proxies.

The near-IR luminosity of a galaxy is a good measure of its stellar mass (Kauffmann & Charlot 1998); therefore, we convert the IRAC [3.6] and [4.5] photometry of the RLAGN and control galaxies to rest-frame luminosities using the spectroscopic and photometric redshifts, respectively. We then linearly interpolate or extrapolate these two data points to measure the rest-frame 1.25  $\mu\text{m}$  luminosities of the galaxies. This wavelength is close to the *J*-band central wavelength and was chosen because it lies in between the observed



**Figure 2.** A comparison of the rest-frame 1.25  $\mu\text{m}$  absolute magnitudes (top panel) and stellar masses (bottom panel) between the CARLA RLAGN and the control galaxies. There is no significant difference in the rest-frame 1.25  $\mu\text{m}$  absolute magnitudes, and the stellar masses of the control galaxies are very similar to the stellar masses of the 25 CARLA RLAGN that were measured using SED fitting of 3.6–70  $\mu\text{m}$  photometry by Seymour et al. (2007). Hence, the control galaxies are well matched in stellar mass to the CARLA RLAGN.

[3.6] and [4.5] bands for the majority of the galaxies. As illustrated in the top panel of Fig. 2, the CARLA radio galaxies and control galaxies are so similar in rest-frame 1.25  $\mu\text{m}$  absolute magnitude that a KS test is unable to distinguish between these samples ( $P = 0.97$ ).

Approximately a fifth of our reduced CARLA radio galaxy subsample have stellar masses measured by Seymour et al. (2007) using SED fitting of 3.6–70  $\mu\text{m}$  photometry. These masses are very similar to those of the control galaxies (measured through SED fitting of the *U* band to 4.5  $\mu\text{m}$  photometry), as shown in the bottom panel of Fig. 2. A KS test results in  $P = 0.15$ , which means there is no significant difference between the masses of these 25 RLAGN and

the control galaxies. These two tests suggest that the control galaxy sample is well matched in mass to the CARLA RLAGN.

### 2.2.3 Galaxy type

Whilst forming a control sample, it is important to consider galaxy type, as passive galaxies are located in denser environments than their star-forming counterparts up to  $z \sim 1.8$  (Chuter et al. 2011; Quadri et al. 2012). RLAGN at  $1.3 < z < 3.2$  do not conform to a single galaxy type. RLAGN hosts at  $z \lesssim 1.8$  tend to contain old, quiescent stellar populations, and their light profiles are similar to the de Vaucouleurs profile (e.g. Best et al. 1998). In contrast, RLAGN hosts at  $z \gtrsim 2$  often have clumpy morphologies and high star formation rates (Pentericci et al. 1999; Drouart et al. 2014). It is likely that the CARLA RLAGN comprise a range of galaxy types. The control sample similarly contains a mix of galaxy types, with approximately 55 per cent defined as quiescent.

With only IRAC colour information for the CARLA RLAGN, it is not possible to form a control sample that matches galaxy type exactly. However, we checked our results whilst limiting the control sample to quiescent galaxies only, and then with star-forming galaxies only, and found no difference in the results (see Section 3.1 for more details).

### 2.3 Measuring environment

The environments of the CARLA RLAGN and control galaxies were measured by counting the number of *Spitzer* IRAC colour-selected galaxies within 1 arcmin radius circles centred on each galaxy in the samples. This radius was chosen because it corresponds to an angular size of  $\sim 0.5$  Mpc at  $1.3 < z < 3.2$ , which is the typical radius of a high-redshift cluster. The well-tested  $[3.6] - [4.5] > -0.1$  colour cut was used to remove galaxies with  $z < 1.3$  (Papovich 2008; Muzzin et al. 2013a; Rettura et al. 2014, see discussion in Galametz et al. 2012). Using this criterion, we expect to obtain a  $z > 1.3$  galaxy sample with only 10–20 per cent contamination by low-redshift interlopers (Muzzin et al. 2013a). This colour selection criterion identifies a homogeneous sample of galaxies out to  $z \sim 3.2$  because  $[4.5]$  is almost constant for galaxies at  $z > 0.7$  due to a negative  $k$ -correction, and these wavelengths cover the  $1.6 \mu\text{m}$  stellar bump for galaxies at  $1.3 < z < 3.2$ . This stellar bump is a prominent feature regardless of a galaxy's star formation history. For the rest of this work, we refer to the objects matching the  $[3.6] - [4.5] > -0.1$  colour selection as 'IRAC-selected sources'.

An object is defined as an IRAC-selected source if it is detected above the  $4.5 \mu\text{m}$  95 per cent completeness limit of the CARLA data ( $[4.5] = 22.9$  mag) and has a colour of  $[3.6] - [4.5] > -0.1$ . If the source is not detected at  $3.6 \mu\text{m}$ , an upper limit of the  $[3.6] - [4.5]$  colour is determined using the  $3.5\sigma$  detection limit of the  $3.6 \mu\text{m}$  CARLA data ( $[3.6] = 22.8$  mag). This is the same criteria used as in Wylezalek et al. (2014), although it differs from that used in Wylezalek et al. (2013) in terms of the  $3.6 \mu\text{m}$  depth.

Most galaxies associated with the RLAGN will have magnitudes fainter than  $m^* - 1$ , where  $m^*$  is the characteristic apparent magnitude of the protoclusters surrounding the RLAGN (McLure & Dunlop 2001). Wylezalek et al. (2014) measured  $m^*$  at  $[4.5]$  for all protoclusters associated with the CARLA RLAGN in six redshift bins spanning  $1.3 < z < 3.2$ . They found  $m^*$  to be in the range of  $19.85$  to  $20.41 \pm 0.20$  mag, which is consistent with the protocluster galaxies forming at  $z \sim 3$  and passively evolving thereafter.  $m^*$  at

$4.5 \mu\text{m}$  is approximately constant across  $1.3 < z < 3.2$  because of a negative  $k$ -correction caused by the  $1.6 \mu\text{m}$  stellar bump that enters the *Spitzer* IRAC bands in this redshift range. Therefore, we assume that the average  $m^*$  of  $\sim 20.1$  mag is valid for all the CARLA protoclusters,<sup>2</sup> and any IRAC-selected source with  $[4.5] < 19.1$  mag was not included in our measurement of the environment for both the RLAGN fields and the control field. This selection should also remove many of the low-redshift interlopers.

### 2.4 Properties of CARLA RLAGN

If RLAGN are good tracers of protoclusters because they are easier to detect when they reside in dense environments, we expect to see correlations between their environment and their radio properties. We therefore measured the spectral index,  $\alpha$  (where  $S_\nu \propto \nu^\alpha$ ), bolometric luminosity, radio emission extent and supermassive black hole (SMBH) mass of the CARLA RLAGN to investigate possible trends with our environmental measurement.

#### 2.4.1 Black hole masses and bolometric quasar luminosity

The SMBH masses and the bolometric luminosities of the 211 RLQs in the CARLA sample were obtained from the SDSS (Shen et al. 2011). The SMBH masses are virial masses, i.e. it is assumed that the broad-line emission region is virialized and the continuum luminosity and broad emission line width are good proxies for the broad-line region radius and velocity, respectively. Due to the spectral coverage of the SDSS data, the Mg II line is used to calculate the virial mass of the SMBH for RLAGN at  $z < 1.9$ , whilst for all higher redshift RLAGN the C IV line is used (McLure & Jarvis 2002; Vestergaard 2002). We note that there are large uncertainties and significant systematic biases associated with these SMBH mass estimates, which are described in detail in Shen et al. (2011).

#### 2.4.2 Extent of the radio emission

The sizes of the radio emission of CARLA sources were measured from the Very Large Array (VLA) Faint Images of the Radio Sky at Twenty-cm (FIRST) survey (White et al. 1997) at  $\sim 1.4$  GHz which has a resolution of 5 arcsec. We matched the 2013 June 05 version of the FIRST catalogue with the CARLA sources and identified 284 sources covered by FIRST. All sources in the catalogue that have a  $>5$  per cent probability of being a spurious source were removed; the RLAGN are very bright so it is common that nearby sources are sidelobes.

We identified RLAGN which had multiple radio components within 1 arcmin, and defined the radio extent as the largest distance between the radio sources detected within 1 arcmin of the RLAGN host galaxy. All sources were visually inspected for a classical radio component distribution. For AGN with only single radio components, we identified the closest radio counterpart within 5 arcsec and the size was defined as the full width at half-maximum (FWHM) of the major axis, which had been deconvolved to remove blurring by the elliptical Gaussian point spread function, down to a major axis FWHM  $< 2$  arcsec. Sources with major axis FWHM  $< 2$  arcsec were classified as unresolved. Of the 284 CARLA RLAGN with FIRST coverage, 127 are extended sources (74 with multiple radio components) and 157 are unresolved sources.

<sup>2</sup> We found no difference in our results if we used the full range of  $m^*$  found by Wylezalek et al. (2014) as opposed to the average  $m^*$ .

### 2.4.3 Spectral indices


The spectral index of the radio emission was measured by cross-correlating two radio surveys with similar spatial resolution (45–80 arcsec): the 1.4 GHz NVSS (Condon et al. 1998) and the 74 MHz VLA Low-Frequency Sky Survey (VLSS; Cohen et al. 2007). RLQs may emit time-variable Doppler beamed emission so it is not reliable to measure their spectral indices from surveys taken several years apart. Therefore, we only measure the spectral indices for the 158 radio-loud galaxies in the CARLA sample that are covered by both the NVSS and VLSS radio surveys.

## 3 RESULTS

### 3.1 Do RLAGN trace protoclusters simply because they are hosted by massive galaxies?

If RLAGN reside in dense environments simply because they are hosted by massive galaxies, then we expect massive radio-quiet galaxies to occupy similarly dense environments. In Fig. 3 we compare the environments of CARLA RLAGN to the control sample. The environments of all 419 RLAGN in the CARLA sample are very similar to the radio galaxy subsample (a KS test gives  $P = 0.95$ ). The environments of the control galaxies are on average less dense than the surroundings of the RLAGN in both CARLA samples. A KS test results in a probability of  $< 10^{-4}$  ( $\sim 4\sigma$  significance) that the radio-loud and radio-quiet galaxies reside in similar environments. This means the high mass of the RLAGN hosts is not the only reason why they trace rich environments.

Galaxy type may also influence our results as quiescent galaxies are located in denser environments than actively star-forming




**Figure 3.** A comparison of the environment surrounding the CARLA RLAGN (red and black) and the UDS control sample (blue). The environment is assessed using the density of IRAC-selected sources within an arcmin radius. The black dashed histogram shows the full CARLA sample. The solid red histogram comprises the radio galaxy CARLA subsample described in Section 2.2.1. The environments of the subsample and full CARLA sample are statistically indistinguishable (KS test results in  $P = 0.95$ ). However, the environment of the control sample differs significantly from both CARLA samples (KS,  $P \sim 10^{-4}$ ).

galaxies, at fixed stellar mass, even up to  $z \sim 1.8$  (Quadri et al. 2012). We therefore checked the influence of galaxy type on our results by limiting the control sample to quiescent galaxies only, and then with star-forming galaxies only. We found no difference in the results, with both quiescent and star-forming control samples differing from the CARLA sample with  $4\sigma$  significance. This means that RLAGN reside in richer environments than all types of similarly massive galaxies.

In Fig. 4 we show the difference between the environments of the RLAGN and the control galaxies in more detail by comparing their average radial density profiles. The top panel of Fig. 4 shows that the environments of RLAGN are denser than the control galaxies at all scales, having both more nearby neighbours and a larger excess beyond 0.5 Mpc.

We then compare the radial profiles surrounding RLAGN and control galaxies that have similar densities within a 1 arcmin radius, choosing both high-density regions with  $14 < \Sigma < 18$  IRAC-selected galaxies per arcmin<sup>2</sup> (middle panel of Fig. 4) and under-dense regions with  $7 < \Sigma < 11$  IRAC-selected galaxies per arcmin<sup>2</sup> (bottom panel of Fig. 4). The central densities are comparable by construction, but beyond 1 arcmin ( $\sim 0.5$  Mpc) the profiles diverge: RLAGN reside in denser large-scale environments than the control galaxies.



**Figure 4.** The radial profile of the IRAC-selected galaxy density surrounding CARLA RLAGN and control galaxies. The top panel displays all CARLA and control galaxies, the middle panel shows galaxies which have central arcmin densities between 14 and 18 IRAC-selected galaxies per arcmin<sup>2</sup> and the bottom panel shows galaxies with central arcmin densities in the range of 7–11 IRAC-selected galaxies per arcmin<sup>2</sup>. The central RLAGN and massive control galaxy are not included, and uncertainties are  $\sqrt{N}$ . The RLAGN are in denser environments on all scales. In the bottom two panels, the profiles of the central arcmin are similar (by construction), but at larger radii the RLAGN are denser than the control galaxies indicating that larger and more massive structures surround the RLAGN. The radius is shown both in arcmin and proper angular distance for galaxies at  $z = 2.2$  (which is the median redshift of the CARLA sample).

This excess at large radii is not an artefact of the data reduction as the raw number counts of the SpUDS control field and the CARLA fields are in good agreement (shown in fig. 2 of Wylezalek et al. 2013). These radial profiles imply that the structures surrounding the RLAGN are more extended than those around massive radio-quiet galaxies. It is possible that the structures surrounding RLAGN are more massive or, alternatively, the extended structure may be the signature of merging galaxy groups or clusters. Simpson & Rawlings (2002) suggested that powerful RLAGN, like the CARLA galaxies, may pinpoint merging clusters as they may be triggered by galaxy–galaxy interactions that occur during mergers. Regardless of the origin of the extended structure, Fig. 4 shows that RLAGN are more likely to reside in high-mass groups and clusters than the average radio-quiet massive galaxy, even if their  $<1$  arcmin environment (0.5 Mpc) appears average or underdense.


Overall, these results mean that the probability that a radio jet is launched from an AGN depends on the Mpc-scale environment, as well as the galaxy mass. These results are in agreement with studies of  $z < 0.7$  RLAGN, such as Kauffmann et al. (2008) and Ramos Almeida et al. (2013), who showed that RLAGN are located in significantly denser environments than similarly massive samples of quiescent galaxies. They also suggest that the high-density environment promotes the launch of radio jets from SMBHs. Our results extend these studies to the high-redshift Universe, and show that environment continues to have a strong influence on AGN properties at  $1.3 < z < 3.2$ .

The overdensities surrounding the CARLA RLAGN extend beyond 2 arcmin, i.e. 2.3–4.2 cMpc (comoving Mpc). According to the models of Chiang et al. (2013), this implies that the typical masses of the overdensities are greater than  $10^{14} M_{\odot}$ . This mass is spread across a few hundred cMpc<sup>3</sup>, and many of these regions are likely to be diffuse protoclusters rather than rich clusters. In time, these regions will collapse to become rich clusters, but at the epoch of observation, the local environment of the high-redshift RLAGN may be similar to that of low- and intermediate-redshift RLAGN, which are generally located in galaxy groups and poor clusters (Fisher et al. 1996; Best 2000, 2004; McLure & Dunlop 2001). It is possible that this particular environment, where galaxy mergers and harassment are likely to be frequent, is very efficient at creating radio jets.

### 3.2 Observational selection biases

In the previous section, we showed that RLAGN are generally found in denser environments than radio-quiet galaxies. In this section, we examine whether there is an observational selection bias that makes it easier for us to observe RLAGN if they are located in dense environments.

Barthel & Arnaud (1996) suggested that radio luminosity is enhanced in rich environments because the relativistic plasma is confined by dense intracluster medium (ICM). This confinement prevents the radio lobes from expanding adiabatically, and a larger fraction of the electron’s energy is lost through synchrotron emission rather than expansion losses. Therefore, these confined sources are more efficient in transferring AGN power into radio luminosity. In a luminosity-limited sample of radio-loud galaxies, such as the CARLA sample, this effect may create a selection bias. If the relativistic plasma is confined, we also expect the source to appear smaller and the spectral index of their radio emission to steepen due to Fermi acceleration across the shock fronts as the hotspots decelerate (Athreya & Kapahi 1998; Klammer et al. 2006).



**Figure 5.** Examining relationships between the large-scale environment of the CARLA RLAGN and properties of the radio emission and SMBH. From top to bottom: (a) the extent of the radio emission for resolved CARLA sources, (b) ratio of radio to bolometric AGN power (for the RLQ only), (c) spectral index  $\alpha$  (where  $S_{\nu} \propto \nu^{\alpha}$ ) and (d) SMBH mass for the 211 CARLA AGN matched with the Shen et al. (2011) SDSS catalogue. Filled circles are SMBH masses derived from Mg II whilst open circles are derived from C IV. Open diamonds are the mean in each density bin. There is no correlation between RLAGN environment and any radio property, although there is a low-significance ( $2.3\sigma$ ) weak correlation with SMBH mass.

We test whether the properties of the radio emission are affected by the ambient gas in rich environments in Fig. 5 by investigating correlations between the environments of the CARLA galaxies and their extent, spectral index and the ratio of their radio to bolometric luminosity (for the quasars). There are two caveats to our method. First, our measure of environment is not robust for individual objects as it is strongly influenced by cosmic variance along the line of sight. Nevertheless, our large sample means this variance should balance out and we are able to measure statistical properties. Secondly, complex trends exist between radio extent, spectral index, luminosity and redshift (Blundell & Rawlings 1999). We try to minimize these issues by using the CARLA sample which is comprised of only the most powerful RLAGN which reside in a narrow 3 Gyr window ( $1.3 < z < 3.2$ ), yet these intrinsic relations may make it difficult to find independent correlations with environmental galaxy density.

As shown in Fig. 5, there is no significant correlation between the radio properties of the RLAGN and its environment. The mean radio extent of resolved sources is  $\sim 125$  kpc (median  $\sim 80$  kpc) and is approximately constant with density. A Spearman rank correlation test shows that neither the ratio of radio to bolometric luminosity nor

**Table 1.** Results from Spearman rank correlation test between the listed parameters and the surrounding galaxy density of RLAGN.  $P$  is the probability of obtaining the rank coefficient by chance;  $P$  values greater than 0.05 mean the correlation is not significant. The only significant correlation is between the environmental galaxy density and the SMBH mass of the RLAGN.

	Number	Spearman rank coefficient	$P$
Extent of radio emission	127	0.03	0.77
$L_{\text{Radio}}/L_{\text{Bolometric}}$	211	-0.05	0.40
Spectral index	158	0.08	0.29
SMBH mass	211	0.16	0.02

the spectral index is correlated with the surrounding galaxy density (see Table 1). There is also no significant difference between the environments of radio sources that are resolved in the FIRST survey and those that are not (KS,  $P = 0.16$ ). Taking into account the caveats discussed above, this indicates that the size of the radio emission is not affected by the Mpc-scale environment of the AGN host galaxy.

There is no evidence that the sizes, spectral indices or radio luminosities of the AGN are affected by their environment, but since all three properties depend on other properties, such as age, orientation and redshift, we are unable to draw strong conclusions from these null results alone. However, the literature also suggests that powerful radio emission is not enhanced in dense environments, nor is a dense environment a prerequisite for visible radio emission. Wylezalek et al. (2013) found no correlation between the radio power of the CARLA AGN and their environment over two orders of magnitude in radio power, covering  $10^{27.5-29.5}$  W Hz $^{-1}$ . Similarly, Karouzos et al. (2014) found no trend between radio power and environmental density of lower luminosity RLAGN, though at lower redshifts, and Donoso et al. (2010) found a negative trend between radio power and galaxy clustering for radio galaxies. In contrast to these works, Falder et al. (2010) found a trend of increasing galaxy overdensity with increasing AGN radio luminosity for  $z \sim 1$  RLAGN. These works all look at different ranges of radio power, which may explain some of the differences in the results. Whilst there is not unanimous agreement amongst the literature, the majority of studies suggest that the radio luminosity of an RLAGN is not amplified in dense environments.

Powerful radio jets can also be observed outside dense environments: Venemans et al. (2007) searched for protoclusters around eight RLAGN and found that two were not in dense environments. Additionally, almost a quarter (24 per cent) of CARLA RLAGN reside in average environments: the galaxy densities within 1 arcmin of these sources are less than  $1\sigma$  above the control field density. Whilst these RLAGN exhibit richer-than-average environments at radii greater than 1 arcmin (Fig. 4), the radio emission from all CARLA RLAGN is limited to within 1 arcmin of the host galaxy and therefore is unlikely to be greatly affected by the gas associated with these more distant structures. This implies that dense ambient gas is not a prerequisite for high-redshift radio jets to be detected.

We conclude that radio emission from the CARLA sample is not significantly or systematically amplified when a CARLA source lies within a dense environment, and therefore there is no observational selection bias that makes it easier for us to observe RLAGN when they are located in richer environments. Since we cannot invoke an observational selection bias to explain the results from Section 3.1 (that RLAGN reside in denser environments than radio-quiet galaxies), we resort to the alternative explanation: the probability of the

central SMBH launching a radio-emitting jet depends on the host galaxy's environment.

If the large-scale environment influences the properties of the SMBH directly, we may expect a correlation between the SMBH mass and its environment. Although there is a great deal of scatter in Fig. 5(d), we find a mildly significant, but small tendency for more massive SMBHs to lie in denser environments: the Spearman correlation coefficient is 0.16 at 98 per cent confidence. This correlation may simply result from the relation between stellar and SMBH mass (Ferrarese & Merritt 2000) and the tendency for high-mass galaxies to reside in denser environments than low-mass galaxies. Therefore further investigation of this trend requires a study of the environment and SMBH mass of RLAGN at fixed stellar mass. Because the significance of the correlation is so low, we do not discuss the result further; however, even if this correlation is real, it cannot explain why RLAGN tend to reside in dense environments. Powerful radio emission from AGN requires the SMBHs to have masses greater than  $10^8 M_{\odot}$ , but there is no strong correlation between black hole mass and radio luminosity (McLure & Jarvis 2004).

## 4 DISCUSSION

### 4.1 Why do RLAGN tend to reside in protoclusters?

The main goal of this study is to examine why RLAGN are commonly found in protoclusters. We showed that RLAGN reside in denser environments than similarly massive radio-quiet galaxies, which suggests a connection between the Mpc-scale environment and radio loudness. We also showed that this connection is not due to the ICM environment amplifying the radio emission. Our results therefore suggest that the *launching* of powerful radio jets depends in some way on the environment of the host galaxy. We can, however, only speculate on ways the Mpc-scale environment can influence the properties of the AGN on sub-pc scales.

The semi-analytic galaxy formation models of Fanidakis et al. (2011) suggest that galaxy mergers are very important for determining whether an AGN becomes radio-loud. Fanidakis et al. (2011) were able to reproduce the radio luminosity function as long as the black holes accreted gas in a chaotic manner (which keeps the spin of most black holes low; King & Pringle 2006), but the spin of the most massive black holes was driven by black hole mergers (which themselves are driven by the galaxy merger rate). These black hole mergers result in rapidly spinning black holes, which may be more likely to launch powerful jets (e.g. Blandford & Znajek 1977). We expect high galaxy merger rates in protoclusters because of the high galaxy densities and the relatively low velocity dispersions compared to virialized clusters (Venemans et al. 2007; Shimakawa et al. 2014). Indeed, Lotz et al. (2011) found that the merger rate in a  $z = 1.62$  protocluster was  $\sim 10$  times higher than the coeval field rate. High galaxy merger rates imply high black hole merger rates, and therefore high black hole spins, which may explain why protoclusters commonly host RLAGN.

However, the radio loudness of an AGN depends on more than just the spin of the black hole; the accretion mode of the AGN is also important (Tchekhovskoy, Narayan & McKinney 2010). Fernandes et al. (2011) found a correlation between jet power and accretion rate, which implies that high accretion rates are required to generate the powerful jets of the CARLA RLAGN. Frequent galaxy mergers in the protocluster environment will dump cold gas into the RLAGN host galaxy. Some fraction of this gas is likely to have sufficiently low angular momentum to fall into the nucleus, leading to high accretion rates on to the black hole. Thus, the protocluster



environment may promote high accretion rates on sub-pc scales, which are required to generate powerful RLAGN.

#### 4.2 Fraction of massive galaxies that become powerful RLAGN

The environments of radio-quiet massive galaxies differ from those of RLAGN, which means only a subset of massive galaxies undergo radio-loud feedback as powerful as  $L_{500\text{MHz}} = 10^{27.5} \text{ W Hz}^{-1}$ . We estimate this fraction by scaling the CARLA probability density histogram in Fig. 3, using a least-squares fitting algorithm, so it matches the height of the control histogram at high densities (14–20 galaxies per arcmin<sup>-2</sup>). We find that 50 per cent of the control galaxies have similar environments to the CARLA RLAGN.<sup>3</sup> Therefore only 50 per cent of the massive control galaxies can go through an RLAGN phase as powerful as  $L_{500\text{MHz}} = 10^{27.5} \text{ W Hz}^{-1}$  during  $1.3 < z < 3.2$ .

A major limitation of this result is that we only discuss the most powerful radio-loud sources at  $1.3 < z < 3.2$ , rather than all RLAGN. We cannot constrain the fraction of massive galaxies that have RLAGN feedback, only the fraction that undergo the most powerful feedback usually associated with Fanaroff–Riley type II radio galaxies.

#### 4.3 Total lifetime of the radio-emitting phase

The proportion of time an RLAGN host galaxy is observed as radio loud can be estimated through the ratio of RLAGN to potential host galaxies:

$$f = \frac{n(\text{RLAGN})}{n(\text{host}) \times f_{\text{RL}}}, \quad (1)$$

where  $f_{\text{RL}}$  is the fraction of galaxies that can become radio-loud ( $< 0.5$ ),  $n(\text{host})$  is the number density of galaxies that have similar masses as the RLAGN galaxies and  $n(\text{RLAGN})$  is the number density of RLAGN with  $L_{500\text{MHz}} > 10^{27.5} \text{ W Hz}^{-1}$ , which is  $5\text{--}10 \times 10^{-8} \text{ Mpc}^{-3}$  comoving at  $1.3 < z < 3.2$  (Rigby et al. 2011).

A large uncertainty in this calculation comes from  $n(\text{host})$ . The stellar mass function is so steep at the high-mass end that  $n(\text{host})$  is completely dependent on the number density of the most massive hosts (i.e. galaxies with  $\gtrsim 10^{11.5} \text{ M}_{\odot}$ ). The number density of  $> 10^{11.5} \text{ M}_{\odot}$  galaxies is  $\sim 1\text{--}7 \times 10^{-6} \text{ Mpc}^{-3}$  (Santini et al. 2012; Muzzin et al. 2013b), and does not appear to evolve rapidly across the  $1.3 < z < 3.2$  redshift interval. However, the uncertainties on this number density are large because cosmic variance has a strong impact on measuring the density of such rare sources.

Using equation (1), we estimate that a powerful RLAGN is radio bright for at least 2 per cent of the time between  $z = 3.2$  and 1.3, i.e.  $\gtrsim 60$  Myr. The radio-bright lifetime of RLAGN is limited to a few  $\times 10$  Myr (Blundell & Rawlings 1999), so this result is consistent with galaxies undergoing a single powerful RLAGN during this period. However, since we derive lower limits of the total radio-emitting lifetime, we cannot rule out that powerful radio-loud AGN feedback is a reoccurring phenomenon with each RLAGN host galaxy undergoing multiple powerful radio-loud episodes between  $z = 3.2$  and 1.3.

<sup>3</sup> We obtain similar results when we run the algorithm using different density bin sizes, and when we use the full CARLA sample or the CARLA radio galaxy subsample.

#### 4.4 Heating the ICM

Approximately half of CARLA RLAGN reside in rich environments containing IRAC-selected overdensities of  $> 2\sigma$  significance (Wylezalek et al. 2014). These environments are highly likely to be protoclusters or clusters, several of which have already been confirmed (Venemans et al. 2007; Galametz et al. 2010, 2013; Hatch et al. 2011b). However, Fig. 4 shows that even the RLAGN which have low densities in the central arcmin have richer environments at larger radii ( $> 0.5$  Mpc). Therefore it is possible that the majority of RLAGN reside in clusters and protoclusters.

The maximum number density of galaxies that become radio-loud is a few  $\times 10^{-6} \text{ Mpc}^{-3}$  and is therefore similar to that of  $> 10^{14} \text{ M}_{\odot}$  galaxy clusters in the present day (Vikhlinin et al. 2009). It is therefore possible that every galaxy cluster hosted a powerful RLAGN at some point between  $1.3 < z < 3.2$ .

The minimum radio-loud lifetime allows us to estimate the total energy deposited into the forming ICM. The energy output by powerful RLAGN at  $z \geq 2$  is a few  $\times 10^{59}$  erg in 30 Myr, considering only the electron population, and up to  $10^{62}$  erg when including the protons in the jets (Erlund et al. 2006; Johnson et al. 2007; Erlund, Fabian & Blundell 2008).

It is well known that extra heating of the ICM on top of gravitational heating, of about 0.5–1 keV per particle, is required to reproduce the observed excess of ICM entropy (Kravtsov & Borgani 2012). In a cluster of gas mass  $1 \times 10^{14} \text{ M}_{\odot}$ , this would require  $1\text{--}2 \times 10^{62}$  erg. So the energy deposited by the RLAGN across their active lifetime is sufficient to provide a large fraction of this energy to the ICM. If jets are able to couple efficiently with the surrounding gas, then these powerful early feedback events can have a dramatic and long-lasting impact on the intracluster gas. Heating the intracluster gas may choke the protocluster galaxies by cutting off the gas supply and limiting further stellar and SMBH growth (Rawlings & Jarvis 2004).

## 5 CONCLUSIONS

We have shown that the environments of high-redshift RLAGN and similarly massive radio-quiet galaxies differ so significantly that less than half of the massive galaxy population (with masses in the range  $10^{11\text{--}11.5} \text{ M}_{\odot}$ ) at  $1.3 < z < 3.2$  may host a powerful RLAGN. Even when we compare the subset of RLAGN and control galaxies that have the same density within  $\sim 0.5$  Mpc, we find that the RLAGN have richer environments beyond this radius, implying that they reside in larger structures. Thus, having a high stellar mass does not ensure that a galaxy will host an RLAGN, and we suggest that the presence of a radio-loud jet may be influenced by the Mpc-scale environment of the host galaxy.

The correlation between radio-loud AGN activity and high-density environments is not the result of an observational selection bias. We examined whether dense environments amplify the radio emission from an RLAGN by confining the radio-emitting plasma or transforming AGN power into radio power. We found no correlation between the environment and the radio size, spectral index or fraction of AGN power emitted at radio frequencies. In addition, approximately a quarter of RLAGN appear to reside in average environments on 0.5 Mpc scales (which is typically the size of the radio emission). We therefore conclude that dense environments do not greatly amplify the radio emission from an RLAGN at this redshift, but the probability that a jet is launched from the host galaxy is likely to depend on the Mpc-scale environment.

We estimate the maximum space density of galaxies that experience a radio-loud episode in the epoch at  $1.3 < z < 3.2$  to be a few  $\times 10^{-6} \text{ Mpc}^{-3}$ , which is similar to the space density of protoclusters: objects that can collapse to form a  $> 10^{14} M_{\odot}$  galaxy cluster by the present day. Distant clusters and protoclusters are reliably found near powerful radio-loud AGN (Venemans et al. 2007; Galametz et al. 2010, 2013; Hatch et al. 2011a,b; Wylezalek et al. 2013) so it is possible that every cluster progenitor experienced a powerful feedback episode during  $1.3 < z < 3.2$ . We estimate that each powerful radio-loud galaxy is active for at least 60 Myr, and if the jets contain protons, then this feedback could provide enough energy per gas particle to pre-heat the forming ICM.

## ACKNOWLEDGEMENTS

We sincerely thank the referee for providing useful and constructive comments which improved this paper. NAH acknowledges support from STFC through an Ernest Rutherford Fellowship. NS is the recipient of an ARC Future Fellowship. MJJ thanks the South African SKA for support. This work is based on observations made with the *Spitzer Space Telescope*, which is operated by the Jet Propulsion Laboratory, California Institute of Technology.

## REFERENCES

- Athreya R. M., Kapahi V. K., 1998, *J. Astrophys. Astron.*, 19, 63  
 Barthel P. D., Arnau K. A., 1996, *MNRAS*, 283, L45  
 Best P. N., 2000, *MNRAS*, 317, 720  
 Best P. N., 2004, *MNRAS*, 351, 70  
 Best P. N., Longair M. S., Roettgering H. J. A., 1998, *MNRAS*, 295, 549  
 Blandford R. D., Znajek R. L., 1977, *MNRAS*, 179, 433  
 Blundell K. M., Rawlings S., 1999, *Nature*, 399, 330  
 Bruzual G., Charlot S., 2003, *MNRAS*, 344, 1000  
 Castignani G., Chiaberge M., Celotti A., Norman C., De Zotti G., 2014, *ApJ*, 792, 114  
 Chabrier G., 2003, *PASP*, 115, 763  
 Chiang Y.-K., Overzier R., Gebhardt K., 2013, *ApJ*, 779, 127  
 Chuter R. W. et al., 2011, *MNRAS*, 413, 1678  
 Cohen A. S., Lane W. M., Cotton W. D., Kassim N. E., Lazio T. J. W., Perley R. A., Condon J. J., Erickson W. C., 2007, *AJ*, 134, 1245  
 Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J., 1998, *AJ*, 115, 1693  
 Croom S. M., Smith R. J., Boyle B. J., Shanks T., Miller L., Outram P. J., Loaring N. S., 2004, *MNRAS*, 349, 1397  
 De Breuck C. et al., 2010, *ApJ*, 725, 36  
 Donoso E., Li C., Kauffmann G., Best P. N., Heckman T. M., 2010, *MNRAS*, 407, 1078  
 Drouart G. et al., 2014, *A&A*, 566, A53  
 Erlund M. C., Fabian A. C., Blundell K. M., Celotti A., Crawford C. S., 2006, *MNRAS*, 371, 29  
 Erlund M. C., Fabian A. C., Blundell K. M., 2008, *MNRAS*, 386, 1774  
 Falder J. T. et al., 2010, *MNRAS*, 405, 347  
 Fanidakis N., Baugh C. M., Benson A. J., Bower R. G., Cole S., Done C., Frenk C. S., 2011, *MNRAS*, 410, 53  
 Fazio G. G. et al., 2004, *ApJS*, 154, 10  
 Fernandes C. A. C. et al., 2011, *MNRAS*, 411, 1909  
 Ferrarese L., Merritt D., 2000, *ApJ*, 539, L9  
 Fisher K. B., Bahcall J. N., Kirhakos S., Schneider D. P., 1996, *ApJ*, 468, 469  
 Furusawa H., 2008, *ApJS*, 176, 1  
 Galametz A., Stern D., Stanford S. A., De Breuck C., Vernet J., Griffith R. L., Harrison F. A., 2010, *A&A*, 516, A101  
 Galametz A. et al., 2012, *ApJ*, 749, 169  
 Galametz A. et al., 2013, *A&A*, 559, A2  
 Hartley W. G. et al., 2013, *MNRAS*, 431, 3045  
 Hatch N. A. et al., 2011a, *MNRAS*, 410, 1537  
 Hatch N. A., Kurk J. D., Pentericci L., Venemans B. P., Kuiper E., Miley G. K., Röttgering H. J. A., 2011b, *MNRAS*, 415, 2993  
 Hatch N. A. et al., 2013, *MNRAS*, 436, 2244  
 Hill G. J., Lilly S. J., 1991, *ApJ*, 367, 1  
 Johnson O., Almaini O., Best P. N., Dunlop J., 2007, *MNRAS*, 376, 151  
 Karouzos M., Jarvis M. J., Bonfield D., 2014, *MNRAS*, 439, 861  
 Kauffmann G., Charlot S., 1998, *MNRAS*, 297, L23  
 Kauffmann G., Heckman T. M., Best P. N., 2008, *MNRAS*, 384, 953  
 King A. R., Pringle J. E., 2006, *MNRAS*, 373, L90  
 Klamer I. J., Ekers R. D., Bryant J. J., Hunstead R. W., Sadler E. M., De Breuck C., 2006, *MNRAS*, 371, 852  
 Kravtsov A. V., Borgani S., 2012, *ARA&A*, 50, 353  
 Kroupa P., 2001, *MNRAS*, 322, 231  
 Lotz J. M., Jonsson P., Cox T. J., Croton D., Primack J. R., Somerville R. S., Stewart K., 2011, *ApJ*, 742, 103  
 McLure R. J., Dunlop J. S., 2001, *MNRAS*, 321, 515  
 McLure R. J., Jarvis M. J., 2002, *MNRAS*, 337, 109  
 McLure R. J., Jarvis M. J., 2004, *MNRAS*, 353, L45  
 Miley G., De Breuck C., 2008, *A&AR*, 15, 67  
 Mortlock A. et al., 2013, *MNRAS*, 433, 1185  
 Muzzin A., Wilson G., Demarco R., Lidman C., Nantais J., Hoekstra H., Yee H. K. C., Rettura A., 2013a, *ApJ*, 767, 39  
 Muzzin A. et al., 2013b, *ApJ*, 777, 18  
 Papovich C., 2008, *ApJ*, 676, 206  
 Pentericci L., Röttgering H. J. A., Miley G. K., McCarthy P., Spinrad H., van Breugel W. J. M., Macchetto F., 1999, *A&A*, 341, 329  
 Quadri R. F., Williams R. J., Franx M., Hildebrandt H., 2012, *ApJ*, 744, 88  
 Ramos Almeida C., Bessiere P. S., Tadhunter C. N., Inskip K. J., Morganti R., Dicken D., González-Serrano J. I., Holt J., 2013, *MNRAS*, 436, 997  
 Rawlings S., Jarvis M. J., 2004, *MNRAS*, 355, L9  
 Rettura A. et al., 2014, preprint ([arXiv:1404.0023](https://arxiv.org/abs/1404.0023))  
 Rigby E. E., Best P. N., Brookes M. H., Peacock J. A., Dunlop J. S., Röttgering H. J. A., Wall J. V., Ker L., 2011, *MNRAS*, 416, 1900  
 Roche N., Eales S., Hippelein H., 1998, *MNRAS*, 295, 946  
 Santini P. et al., 2012, *A&A*, 538, A33  
 Schneider D. P. et al., 2010, *AJ*, 139, 2360  
 Seymour N. et al., 2007, *ApJS*, 171, 353  
 Shen Y. et al., 2011, *ApJS*, 194, 45  
 Shimakawa R., Kodama T., Tadaki K.-i., Tanaka I., Hayashi M., Koyama Y., 2014, *MNRAS*, 441, L1  
 Simpson C., Rawlings S., 2002, *MNRAS*, 334, 511  
 Simpson C. et al., 2006, *MNRAS*, 372, 741  
 Stern D., Chary R.-R., Eisenhardt P. R. M., Moustakas L. A., 2006, *AJ*, 132, 1405  
 Tchekhovskoy A., Narayan R., McKinney J. C., 2010, *ApJ*, 711, 50  
 Ueda Y. et al., 2008, *ApJS*, 179, 124  
 Venemans B. P. et al., 2007, *A&A*, 461, 823  
 Vestergaard M., 2002, *ApJ*, 571, 733  
 Vikhlinin A. et al., 2009, *ApJ*, 692, 1060  
 White R. L., Becker R. H., Helfand D. J., Gregg M. D., 1997, *ApJ*, 475, 479  
 Williams R. J., Quadri R. F., Franx M., van Dokkum P., Labbé I., 2009, *ApJ*, 691, 1879  
 Wuyts S. et al., 2007, *ApJ*, 655, 51  
 Wylezalek D. et al., 2013, *ApJ*, 769, 79  
 Wylezalek D. et al., 2014, *ApJ*, 786, 17  
 Yates M. G., Miller L., Peacock J. A., 1989, *MNRAS*, 240, 129

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.