

- ⁹⁹Ghana VAST. Vitamin A supplementation in northern Ghana: effects on clinic attendances, hospital admissions, and child mortality. *Lancet* 1993;**342**:7–12.
- ¹⁰⁰Binka FN, Morris SS, Ross DA, Arthur P, Aryeetey ME. Patterns of malaria morbidity and mortality in children in northern Ghana. *Trans R Soc Trop Med Hyg* 1994;**88**:381–85.
- ¹⁰¹Binka F, Kubaje A, Adjuik M, Williams LA, Lengeler C, Maude GH *et al.* Impact of permethrin impregnated bednets on child mortality in Kassena-Nankana district, Ghana: a randomized controlled trial. *Trop Med Int Health* 1996;**1**:147–54.
- ¹⁰²Unpublished report to TDR/WHO by F. Binka, 1994. (A copy of this report could not be obtained; results from this report were based on a personal written communication from R.W. Snow on December 20, 2002).
- ¹⁰³Snow RW, Mung'ala VO. Unpublished data from Bahari Division, Kilifi, Kenya, May 1991–April 1993.
- ¹⁰⁴Snow RW, Mung'ala VO, Foster D, Marsh K. The role of the district hospital in child survival at the Kenyan Coast. *Afr J Health Sci* 1994;**1**:71–75.
- ¹⁰⁵Snow RW, Omumbo JA, Lowe B, Molyneux CS, Obiero JO, Palmer A *et al.* Relation between severe malaria morbidity in children and level of *Plasmodium falciparum* transmission in Africa. *Lancet* 1997;**349**:1650–54.
- ¹⁰⁶Barnish G, Maude G, Bockaire M, Eggelte T, Greenwood B, Ceesay S. Malaria in a rural area of Sierra Leone. I. Initial results. *Ann Trop Med Parasitol* 1993;**87**:125–36.
- ¹⁰⁷Barnish G, Maude GH, Bockarie MJ, Erunkulu OA, Dumbuya MS, Greenwood BM. Malaria in a rural area of Sierra Leone. II. Parasitological and related results from pre- and post-rains clinical surveys. *Ann Trop Med Parasitol* 1993;**87**:137–48.
- ¹⁰⁸Premji Z, Ndayanga P, Shiff C, Minjas J, Lubega P, MacLeod J. Community based studies on childhood mortality in a malaria holoendemic area on the Tanzanian coast. *Acta Trop* 1997;**63**:101–09.
- ¹⁰⁹Premji Z, Hamisi Y, Shiff C, Minjas J, Lubega P, Makwaya C. Anaemia and *Plasmodium falciparum* infections among young children in an holoendemic area, Bagamoyo, Tanzania. *Acta Trop* 1995;**59**:55–64.
- ¹¹⁰Malaria Burden of Disease in Africa data collection proforma (provided by R.W. Snow) for the reference: "Unpublished data from Brookebond Tea Estate, Kenya, January 1997–December 1998 by Shanks GD, Biomonde K, Snow RW; 1999" Section 6.2 of this proforma cited the following as the source of the parasite prevalence data: Dennis Shanks, personal communication, 1999.
- ¹¹¹Pison G, Trape JF, Lefebvre M, Enel C. Rapid decline in child mortality in a rural area of Senegal. *Int J Epidemiol* 1993;**22**:72–80.
- ¹¹²Omumbo JA, Snow RW. *Plasmodium falciparum* prevalence in East Africa: a review. *East Afr Med J* 2004;**81**:649–56.
- ¹¹³Jelliffe EFP, Jelliffe DB. *Plasmodium malariae* in Ugandan children. I. Prevalence in young children in rural communities. *Am J Trop Med Hyg* 1963;**12**:296–97.
- ¹¹⁴Mirza N, Macharia W, Wafula E, Agwanda R, Onyango F. Mortality patterns in a rural Kenyan community. *East Afr Med J* 1990;**67**:823–29.
- ¹¹⁵Njoroge, EP. *Community parasite survey monthly report*. Kenya: Division of Vector Borne Diseases, MOH, 1987.
- ¹¹⁶Yamagata Y. Malaria control in the United Republic of Tanzania (1993–1996). Report by Japanese International Cooperation Agency to Ministry of Health, Tanzania, October 1996.
- ¹¹⁷Lopez AD, Ahmad OB, Guillot M *et al.* *World Mortality in 2000: Life Tables for 191 Countries*. Geneva: World Health Organization, 2002.

Commentary: Malaria death rates remain highly pertinent

Tom Smith

Scientific papers and media coverage of malaria alike expose us so frequently to figures for the number of child deaths from *Plasmodium falciparum* in Africa that we become insensitive to the magnitude of the carnage. The numbers are so large that they overwhelm the imagination.

Until recently it was hard to work out where these estimates came from, since there was rarely any explanation of the primary sources or the way in which the data were analysed. However several papers in recent years^{1–3} have provided good documentation of their methods, which have steadily become more convincing as more data and appropriate methodology become available. The paper by Rowe *et al.*⁴ in this issue of the

International Journal of Epidemiology presents the best substantiated set of estimates to date in this series, using the latest population estimates for denominators and stratified analyses of an extensive database of verbal autopsy (VA)-based rates. Importantly, they include analyses that show that the estimated number of just over 800 000 deaths in children <5 years of age, for the year 2000, is not very sensitive to their main assumptions.

This paper is not the last word on the subject. The assembled estimates of malaria-specific mortality rates from different sites in Africa vary enormously. We do not know the reasons for this variation. Most of it does not relate to transmission intensity. If we knew the main factors that lead to the diversity in quoted malaria mortality rates in endemic areas it would help to focus control efforts on what works. It would also

give us more confidence in the estimates of average disease burden.

It seems likely that a major reason for the variation in the data is heterogeneity in the way in which malaria is diagnosed in VA. Rowe *et al.*⁴ point out that standard International Classification of Diseases (ICD) rules are not always used to determine causes of death, but in any case these rules were not designed with VA for malaria in mind. Malaria is one of the hardest diagnoses to get right in VA because its symptoms are so non-specific. A recent comparison of cause-specific mortality rates in demographic surveillance sites across Africa that mapped the various local coding systems onto the ICD10 codes managed to classify deaths consistently only by aggregating all fevers of unknown origin with explicit diagnoses of malaria.⁵ This leads to estimates of proportionate mortality substantially greater than the 18% average quoted by Rowe *et al.*⁴

It is possible to make adjustments for this diagnostic inaccuracy,⁶ but even though the authors of these adjustment techniques also collaborated on the Rowe *et al.*⁴ paper, they expressed scepticism of the value of using their own methods.

This is not the only potential interpretational pitfall that they point out. As methods improve, the best estimates change, irrespective of changes in disease burden. Some uncritical users may need to be reminded that the differences between the different estimates reflect mainly methodological developments, not real trends over time. An important real reason for changes over time in number of deaths is changes in the denominator. Rowe *et al.*'s⁴ estimate will certainly be projected to indicate greater number of deaths in the future on the basis of population growth rates. Will this be interpreted as meaning that things are getting worse?

The reader is also reminded that the number of deaths with malaria as the underlying cause is not the same as the number that would be averted by eliminating the disease. Half a century ago, it was shown that elimination of malaria, notably in the sugar plantations of Guyana⁷ and in Sri Lanka,⁸ led to much larger decreases in all-cause mortality than expected from the number of deaths diagnosed pre-intervention as malaria. Rowe *et al.*⁴ cite similar results from malaria control in Africa. In trial settings, insecticide treated nets reduced child mortality by 17%⁹ without coming close to eliminating malaria. We still do not really understand the biology of this indirect mortality, but should we not be estimating the total deaths that might be averted? Who needs to know the number of malaria-specific deaths, apart from those who suggest that funding allocations for diseases should be proportional to their frequency as underlying cause? It would certainly be at least as

useful to have estimates of how many deaths might be averted by specific interventions.

In fact we also need good estimates of malaria-specific death rates because the trend in this index is a key indicator of progress in malaria control. In particular, we need good estimates for more time points. 1990 was the baseline year for the Millennium Development Goals and estimates for that year would be especially useful, as would estimates for more recent years. At first sight the data presented by Rowe *et al.*⁴ seem to indicate a disturbing deterioration in the 1990s, possibly related to increasing drug resistance or failure to invest in health systems. They interpret this cautiously pointing out that study sites change and that the best way to investigate changes over time is time-series within sites. Longitudinal within-site studies will certainly make a major contribution to understanding how the burden of malaria in Africa changes in the years to come, but we still need to continue to improve the methods for estimating the overall picture. The paper by Rowe *et al.*⁴ is a useful step to achieving this.

References

- 1 Snow R, Craig M, Deichmann U, Marsh K. Estimating mortality, morbidity and disability due to malaria among Africa's non-pregnant population [see comments]. *Bull World Health Organ* 1999;**77**:624–40.
- 2 Snow RW, Craig M, Newton C, Steketee RW. *The public health burden of Plasmodium falciparum malaria in Africa: deriving the numbers. Working Paper No. 11. Disease Control Priorities Project.* Bethesda, MD: Fogarty International Center, National Institutes of Health, 2003.
- 3 Hay SI, Guerra CA, Tatem AJ, Atkinson PM, Snow RW. Urbanization, malaria transmission and disease burden in Africa. *Nat Rev Microbiol* 2005;**3**:81–90.
- 4 Rowe A, Rowe SY, Snow RW *et al.* The burden of malaria mortality among African children in the year 2000. *Int J Epidemiol* 2006;2006;**35**:691–704.
- 5 Adjuik M, Smith T, Clark S *et al.* 84, Number 3, March 2006, 161–256. Cause Specific Mortality Rates in developing countries. *Bull World Health Organ* 2006;**84**:181–92.
- 6 Korenromp EL, Williams BG, Gouws E, Dye C, Snow R. Measurement of trends in childhood malaria mortality in Africa: an assessment of progress toward targets based on verbal autopsy. *Lancet Infect Dis* 2003;**3**:349–58.
- 7 Giglioli G. Changes in the pattern of mortality following the eradication of hyperendemic malaria from a highly susceptible community. *Bull World Health Organ* 1972;**46**:181–202.
- 8 Newman P. Malaria and mortality. *J Am Stat Assoc* 1977;**72**:257–63.
- 9 Lengeler C. Insecticide-treated bednets and curtains for preventing malaria. *Cochrane Database Syst Rev* 2004;(2):CD000363.