

The effect of colostrum period management on BW and immune system in lambs: from birth to weaning

L. E. Hernández-Castellano^{1,2}, A. Suárez-Trujillo¹, D. Martell-Jaizme¹, G. Cugno¹, A. Argüello¹ and N. Castro^{1†}

¹Department of Animal Science, Universidad de Las Palmas de Gran Canaria, 35413 Arucas, Gran Canaria, Spain; ²Veterinary Physiology, Vetsuisse Faculty, University of Bern, Bremgartenstrasse 109a, CH-3001 Bern, Switzerland

(Received 28 May 2014; Accepted 10 May 2015; First published online 7 July 2015)

The aim of this study was to investigate the BW and immune status of lambs reared under natural conditions or under artificial conditions fed two different colostrum amounts. In this study, 60 lambs were randomly divided into groups according to treatment. Twenty lambs remained with their dams (natural rearing (NR) group). Forty lambs were removed from their dams at birth. Lambs were bottle-fed with a pool of sheep colostrum, receiving either 4 g of IgG/kg of BW at birth (C4 group) or 8 g of IgG/kg of BW at birth (C8 group). The total colostrum amount was equally divided into three meals at 2, 14 and 24 h after birth. After this period, lambs were bottle-fed a commercial milk replacer. Blood plasma sample analysis and BW recordings were carried out before feeding at birth and then at 1, 2, 3, 4, 5 and 20 days after birth. Another blood sample analysis and BW recording was carried out when animals reached 10 kg of BW. During weaning (30 days), sampling was carried out every 5 days. Blood plasma was used to determine the concentrations of IgG and IgM and the complement system activity – total and alternative pathways. The NR group showed greater BW than the C4 and C8 groups during milk feeding period, whereas the C4 and C8 groups had greater BW than the NR group at the end of weaning period. The C8 and NR groups had greater plasma IgG and IgM concentrations than the C4 group during milk feeding period. In addition, C4 and C8 groups showed similar IgG concentrations and greater IgM concentrations than the NR group at the end of the weaning period. Complement system activity was greater in the NR group than in the C4 and C8 groups during the first 3 days after birth. In conclusion, lambs fed amounts of colostrum equivalent to 8 g of IgG/kg of BW showed similar immune variables compared to lambs reared under natural conditions, obtaining a greater BW at the end of the weaning period. Nevertheless, this study shows that not only the colostrum amount but also the management during the milk feeding and weaning period, such as stress produced by dam separation, milk quality and suckling frequency, can affect the final immune status of lambs.

Keywords: immune, lamb, colostrum, weaning, artificial rearing

Implications

Colostrum plays a fundamental role in the immunization of newborn ruminants. Artificial rearing is a common method used in dairy farms, increasing the amount of milk for processing. The aim of this study was to feed newborn lambs colostrum with two different IgG concentrations and to compare these animals to natural dam fed lambs. This study showed that administration of 8 g of immunoglobulin g/kg BW at birth, equally divided into three colostrum meals (2, 14 and 26 h), achieved higher circulating immunoglobulin G concentration when compared to the natural rearing treatment, although both artificial rearing groups decreased BW after birth.

Introduction

The relationship between colostrum and the survival of newborn ruminants has been long characterized (Argüello *et al.*, 2004; Castro *et al.*, 2011; Abdou *et al.*, 2014). In fact, colostrum contains a complex mixture of proteins that actively participate in the protection of the neonate (passive immune transfer (PIT)) against pathogens and other *postpartum* environmental challenges (Bendixen *et al.*, 2011). Moreover, colostrum has diverse components such as fat, lactose, vitamins or minerals that have high nutritional importance (Ontsouka *et al.*, 2003; Lérias *et al.*, 2014; Hernández-Castellano *et al.*, 2014b).

In particular, colostrum plays an important role in newborn lambs, because they are born hypogammaglobulinemic,

† E-mail: noemi.castro@ulpgc.es

due to the complexity of the synepitheliochorial ruminant placenta, which does not allow sufficient transfer of immunoglobulins from the dam to the fetus (Hernández-Castellano *et al.*, 2014a). In addition, it has been described that lambs not fed colostrum in the first hours of life are more susceptible to diseases and mortality (Ahmad *et al.*, 2000; da Nobrega *et al.*, 2005; Nowak and Poindron, 2006).

Nowadays, there is an increase in the number of high-production dairy farms, where lambs are reared under an artificial feeding system. In such cases, lambs are bottle-fed colostrum and milk replacer in order to increase the amount of milk available for processing (Demiroren *et al.*, 1995; Napolitano *et al.*, 2008; David *et al.*, 2014), thus simplifying animal management (Emsen *et al.*, 2004).

As it has been described by several authors, goat kids need to be fed an amount of colostrum equivalent to 4 g of IgG/kg of BW, divided into three meals before 48 h after birth, in order to make an appropriate PIT (Castro *et al.*, 2005; Morales-delaNuez *et al.*, 2009; Moreno-Indias *et al.*, 2012). No references about the amount of colostrum required by newborn lambs reared under artificial conditions have been found. As it is essential to obtain adequate immunoglobulin concentrations in the blood (O'Doherty and Crosby, 1997; Quigley *et al.*, 2000; Christley *et al.*, 2003), this knowledge has high relevance for the survival of the lambs.

Although the transfer of immunoglobulins (IgG and IgM, mainly) is a key factor for host defense against pathogens, a wider range of colostrum components, such as the complement system proteins, have been suggested to contribute to the early protection of the neonate (Smith and Foster, 2007; Hernández-Castellano *et al.*, 2015a). The complement system activity – comprising the total (TCA) and alternative (ACA) pathways – plays an important role in host defense mechanisms against infectious microbes, because it is involved in specific and non-specific immunity (Rodríguez *et al.*, 2009; Hernández-Castellano *et al.*, 2015b). There are few studies about the complement system activity in sheep and lambs, with the exception of the study published by Hernández-Castellano *et al.* (2015b) and Hernández-Castellano *et al.* (2015c), as these studies are based on an old determination technique (Oswald *et al.*, 1990).

The aim of this study was to investigate the BW and immune status of lambs reared under natural conditions and lambs reared under artificial conditions fed two different colostrum amounts (4 g IgG/kg of BW at birth and 8 g IgG/kg of BW at birth) and milk replacer.

Material and methods

The present study was performed in the Department of Animal Science of the Universidad de Las Palmas de Gran Canaria in the Canary Islands (Spain) on 60 single partum lambs of the Canary breed (30 males and 30 females). As dams were estrous-synchronized, all lambs were born in the same period. Animal procedures were approved by the Ethics Committee of the University.

Animal health status was monitored during the experimental period (for diarrhea, parasites or fever), and the animals were found to be healthy throughout the experimental period.

Milk feeding period

At birth, animals were equally divided by sex and then randomly divided into three different groups. The natural rearing (NR) group was composed of 20 lambs, which suckled colostrum directly from their dams. The other 40 lambs were divided into two groups (20 lambs each) and were housed in artificial rearing rooms. Each room had central heating (20°C) and 0.3 m² floor space per lamb. Lambs were bottle-fed a sheep colostrum pool that was previously pasteurized at 63°C for 30 min according to Trujillo *et al.* (2007). Artificially reared lambs received colostrum amount equivalent to 4 g of IgG/kg of BW at birth (C4 group) or 8 g of IgG/kg of BW at birth (C8 group) divided into three meals at 2, 14 and 24 h after birth. Colostrum meals were always thawed and heated at 37°C according to Argüello *et al.* (2003). Colostrum IgG and IgM concentrations (64.37 and 5.61 mg/ml, respectively) were determined using commercial ELISA kits (Bethyl laboratories, Montgomery, TX, USA), using a purified sheep IgG or a purified sheep IgM as the standard curve.

During the milk feeding period (from birth to 10 kg of BW), NR lambs were raised with dams and had free access to dam milk until the weaning period. Artificially reared lambs (C4 and C8) received a commercial milk replacer at 16% (w/w; MR group, Bacilactol Corderos y Cabritos, Saprogal, La Coruña, Spain; 95.5% dry matter, 23.6% CP and 22.7% ether extract, air-dry powder basis). These groups were fed *ad libitum* (37°C) using nipple buckets twice a day (0800 and 1700 h).

Weaning period

Animals from the three studied groups started the weaning period when they reached 10 kg BW (30.70, 34.45, 34.20 days after birth in NR, C4 and C8 groups, respectively). The weaning period lasted for 30 days in the studied groups. During this period, animals had free access to starter feed (18% CP and 3.4% ether extract), alfalfa hay and water. Lambs from the NR group were separated from dams and placed in a pen for 30 days. Ewes were milked once a day (0900 h). From day 0 to 8 of weaning, lambs had access to ewes twice daily (1000 h after milking and 1700 h), reducing this access to once a day (1700 h) from day 9 to 15. After this, lambs had no more access to the ewes. Artificial rearing groups (C4 and C8) were fed half liter of milk replacer twice a day from day 0 to 8 (1000 and 1700 h), reducing the intake to once daily from day 9 to 15 (1700 h). During the rest of the weaning period, animals did not receive milk replacer.

BW recording and sample collection

Animals were weighed before each blood extraction, and the results are expressed in kg (MOBBA, Barcelona, Spain; accuracy, 5 g). During the milk feeding period, blood samples

were collected at 2 h after birth (day 0), and then blood samples were collected at day 1, 2, 3, 4, 5 and 20 after birth. Another sample was taken when each animal reached 10 kg BW. During the weaning period, samples were obtained every 5 days until the end of the experimental period. Blood samples were collected before morning feeding from the jugular vein in 2.5-ml tubes with K3EDTA. Blood was centrifuged at $2190 \times g$ for 5 min at 4°C (Hettich-Zentrifugen, Universal 32 R, Tuttlingen, Germany), and the plasma was stored at -80°C until analysis.

To determine blood plasma IgG and IgM concentrations, commercial ELISA kits (Bethyl Laboratories) were used, using purified sheep IgG and IgM as the standard curves. Results are expressed in mg of immunoglobulin/ml of plasma.

Complement system activity (total complement activity, TCA and alternative complement activity, ACA) was measured by the hemolytic rate according to a novel technique described by Moreno-Indias *et al.* (2012) and Hernández-Castellano *et al.* (2015b) in goat kid blood serum and lamb blood plasma, respectively. In this technique, a DGHB++ buffer (Hepes Gelatin Veronal Buffer with Ca^{++} and Mg^{++} : 5 mM HEPES, 71 mM NaCl, 0.15 mM CaCl_2 , 0.5 mM MgCl_2 , 2.5% (w/v) glucose, 0.1% (w/v) gelatin, pH 7.4) is used to measure total complement system activity, and DGHB-Mg-EGTA buffer (4.2 mM Hepes, 59 mM NaCl, 7.0 mM MgCl_2 , 2.08% (w/v) glucose, 0.08% (w/v) gelatin, 10 mM EGTA, pH 7.4) is used to measure the alternative pathway. For total complement system activity, rabbit red blood cells and lamb plasma were both diluted to 5% in DGHB++; 100 μl of each was then mixed in a microtiter plate and incubated at 37°C for 1 h. Cells were removed by centrifugation ($2500 \times g$, 5 min, 4°C), and supernatant absorbance was measured at 405 nm using a microplate reader (Model 680; BioRad microplate reader, Hercules, CA, USA). Complete hemolysis was achieved by mixing the cells with distilled water (100 μl), and spontaneous lysis was produced by mixing the diluted rabbit red blood cells with DGHB++. Complement-induced hemolysis of rabbit red blood cells by the test sera was calculated using the following formula: $[(A_{405} \text{ sample} - A_{405} \text{ spontaneous lysis}) / (A_{405} \text{ complete hemolysis} - A_{405} \text{ spontaneous lysis})] \times 100$. The same protocol was performed with DGHB-Mg-EGTA buffer to measure the alternative pathway.

Statistical analyses

Statistical analyses were performed using SAS Version 9.00 (SAS Institute Inc., Cary, NC, USA). As it was described by Brujeni *et al.* (2010) and Ciupersescu (1977), sex was not taken into consideration to simplify the analysis, because no influence was detected. The SAS PROC MIXED procedure for repeated measures was used to evaluate the fixed effect of rearing system, time and the interaction between both effects (rearing system \times time) on plasma IgG and IgM concentrations, complement system activity and BW evolution of lambs from birth to weaning. When significant by ANOVA ($P < 0.05$), differences between means were identified using the Tukey–Kramer test.

Results and discussion

Milk feeding period

Figure 1 shows the BW, IgG and IgM concentrations and complement system activity in the NR, C4 and C8 groups during the colostrum and milk feeding period. Focusing on BW development, NR group values increased from birth BW by day 3 and continued to increase until 10 kg BW was achieved. Nevertheless, BW in C4 and C8 did not increase during the first 5 days after birth ($P > 0.05$), increasing at day 20 and at the end of the milk feeding period ($P < 0.05$). No differences among groups were observed at birth ($P > 0.05$); however, the NR group showed greater BW than the C4 and C8 groups during the rest of the experimental period ($P < 0.05$). Differences observed among groups could be related to the amount of colostrum and milk consumed, as well as the differences between milk and milk replacer composition. As has been described in lambs (Murphy *et al.*, 1994; Napolitano *et al.*, 2002), food restriction could reduce the average daily gain of animals from the C4 and C8 groups, resulting in lower BW.

All the groups (NR, C4 and C8) had a very low IgG concentration in blood plasma at birth, probably from maternal origin through the placenta, as has been referenced by Castro *et al.* (2011). After the first colostrum intake, this concentration rapidly increased in the NR, C4 and C8 groups at 1 to 2 days after birth (16.79, 5.81 and 17.74 mg of IgG/ml, in NR, C4 and C8 groups at day 1 after birth, respectively, $P < 0.05$). After this, IgG concentration decreased until the end of the milk feeding period (5.22, 2.18 and 5.62 mg of IgG/ml, in NR, C4 and C8 groups, respectively, $P < 0.05$). No differences were observed among groups at birth. During the rest of the milk feeding period, IgG concentration was lower in lambs from the C4 group compared to the NR and C8 groups ($P < 0.05$). Moreover, no differences were observed in IgG concentrations between NR and C8 groups during this period (with the exception of day 4, $P = 0.049$). In contrast to these findings, Firat *et al.* (2003) appreciated differences in the milk feeding period between lambs raised under natural and artificial conditions, describing that lambs raised under artificial rearing systems had generally lower plasma IgG concentrations than lambs raised under natural conditions. As these authors have described, the use of a commercial colostrum supplement to feed lambs reared under artificial conditions could affect the immune concentration in blood. In addition, when only artificial rearing groups (C4 and C8) were compared, it was noted that plasma IgG concentrations did not increase according to the provided colostrum IgG concentration, showing that lambs from the C8 group had approximately three times higher plasma IgG concentrations compared to the C4 group. The higher colostrum IgG provided to C8 lambs compared to C4 lambs could reduce the degradation rate of immunoglobulins produced during the digestion process, allowing the absorption of a higher percentage of IgG in its native structure by C8 lambs. A similar absorption pattern has been observed in lambs (Halliday and Williams, 1979),




Figure 1 BW, IgG and IgM concentrations, TCA and ACA evolution in the NR ($n = 20$), C4 ($n = 20$) and C8 ($n = 20$) groups during colostrum and milk feeding periods. NR = natural rearing; C4 = 4 g of IgG/kg of BW at birth; C8 = 8 g of IgG/kg of BW at birth. TCA = total complement activity; ACA = alternative complement activity. *End means the end of milk feeding period and the start of weaning period when animals reached 10 kg of BW. NR, C4 and C8 finished the feeding period and started the weaning period at 30.70 ± 6.05 , 34.45 ± 5.75 , 34.20 ± 6.62 days, respectively. Data shown in this figure can be viewed in the Supplementary Table S1.

goat kids (Castro *et al.*, 2005; Rodríguez *et al.*, 2009) and calves (Muller and Ellinger, 1981; Quigley *et al.*, 2002). In contrast to these results, Moretti *et al.* (2010), using a fixed colostrum volume (250 ml), fed lambs either bovine colostrum (115.7 mg of IgG/ml) or ovine colostrum (48.1 mg of IgG/ml). Because of the differences in both colostrum IgG

concentrations, Moretti *et al.* (2010) concluded that IgG absorption by the enterocytes is limited and, therefore, is non-proportional to the IgG present in the colostrum. Nevertheless, these authors used different colostrum conditions compared to the ones used in the present study (fixed volume with two different IgG concentrations *v.* fixed IgG

concentration and fixed total IgG provided to the lambs). This fact could explain the greater IgG concentrations seen in the blood plasma of lambs from the C8 group compared to lambs from the C4 group. Finally, it has been described that lambs fed bovine colostrum run the risk of developing anemia (Winter, 2011), and the use of caprine colostrum as an alternative way to feed lambs in early life has been recommended (Hernández-Castellano *et al.*, 2015b).

A similar evolution was observed in the case of IgM concentration in blood plasma, although this immunoglobulin was not detectable at birth in any of the studied groups. A rapid increase of IgM in blood plasma was seen in NR, C4 and C8 lambs 1 to 2 days after birth (1.63, 0.35 and 2.84 mg of IgM/ml, in NR, C4 and C8 groups at day 1 after birth, respectively, $P < 0.05$). Despite double colostrum feeding in the C8 group (based on the provided IgG amount) compared to the C4 group, the IgM concentration was much higher in the C8 group than in the C4 group during this period. After that, IgM concentrations in blood plasma decreased until the end of the milk feeding period in the NR and C8 groups. Nevertheless, the C4 group had increased IgM concentrations at day 20 ($P < 0.05$) and at the end of the milk feeding period ($P < 0.05$). According to previous findings, Nonnecke *et al.* (2012) observed a rapid increase in plasma IgG and IgM concentrations after colostrum intake in calves, with a decrease in these concentrations with age. The colostrum-deprived calves, in contrast, had very low or undetectable plasma IgG and IgM concentrations, followed by an age-related increase of IgG and IgM concentrations, suggesting endogenous production of these immunoglobulins. During this period, a similar evolution was only observed in the IgM concentration, probably because this is the first immunoglobulin produced by the organism (Ehrenstein and Notley, 2010). When differences among groups were analyzed, it was noticed that the C8 group reached a greater IgM concentration at 1 to 3 days compared to the NR group. During these days, the lowest IgM concentration was obtained by the C4 group. No references have been found about the IgM absorption rate in lambs. As described for IgG absorption, a higher IgM amount provided to C8 lambs may produce a proportional lower IgM degradation compared to C4 lambs, and therefore a higher absorption of intact IgM by the enterocytes. In addition, no differences among groups were observed at day 20 and at the end of the milk feeding period. As described above, the endogenous production of IgM by the C4 group, related with the deficient colostrum intake, could produce an increase in IgM concentrations in this group. As a consequence, the C4 group obtained similar results to the NR and C8 groups at that time.

No evidence of any detectable complement system activity was observed at birth in the three studied groups (NR, C8 and C4). Conversely, this activity increased in all the groups during the entire study period, although a different rate was observed. In the case of the NR group, lambs had earlier complement system activation, which was greater compared to the C4 and C8 groups at 1 to 3 days. Despite the fact that colostrum freezing for storage does not affect

immunoglobulin concentration in newborn ruminants (Holloway *et al.*, 2002; Argüello *et al.*, 2003), other colostrum components could be affected by the freezing process (Ramírez-Santana *et al.*, 2012). As was described before, the C4 and C8 groups were fed colostrum from a frozen pool that may cause reduction of some complement components present in colostrum. Moreover, dam milk contains some complement system components that are not present in milk replacer. In a similar way, Castro *et al.* (2008) showed the importance of dam milk in earlier complement system activation compared to milk replacer feeding. Nevertheless, no differences were observed between the C8 and NR groups at day 4, showing that the C8 group had an earlier complement system activation than the C4 group, perhaps because of the greater amount of colostrum received by the C8 group. This fact may produce an earlier activation, because colostrum contains high amounts of complement system components (Wheeler *et al.*, 2007). Despite the fact that lambs from the C4 group showed lower TCA and ACA during most of this period, no differences among groups were observed at the end of the milk feeding period. In addition, ACA seems to be the most important pathway in the three groups, as was described by Hernández-Castellano *et al.* (2015b) and Castro *et al.* (2008) in newborn lambs and newborn goat kids, respectively.

Weaning period

Figure 2 shows the BW, IgG and IgM concentrations and complement system activity (TCA and ACA) in the NR, C4 and C8 groups during the weaning period. With reference to BW development, lambs from the NR, C4 and C8 groups increased their BW during this period. No differences were observed among groups, although lambs from the C4 and C8 groups showed a greater BW than lambs from the NR group at day 25 (15.72, 15.63 and 14.62 kg, in C4, C8 and NR, respectively, $P < 0.05$) and day 30 (16.89, 16.95 and 15.28 kg, in C4, C8 and NR, respectively, $P < 0.05$). It has been described by Napolitano *et al.* (2002) that lambs raised with dams suffer greater stress by separation during the weaning period than animals from artificial rearing systems, resulting in a decrease in the final BW. In addition, al-Sabbagh *et al.* (1995) showed that lambs born with low BW showed compensatory growth before weaning compared to lambs with high BW. This fact explains the negative correlation between the weight of the lamb at birth and the final weaned lamb weight, as described by those authors. Despite the lower weight of the NR group at the end of the weaning period, no differences in the BW at birth were detected between the three studied groups (NR, C4 and C8).

Regarding IgG concentration, lambs from the NR and C8 groups showed slight decreases ($P < 0.05$) in plasma IgG concentration in some specific time points of weaning. Nevertheless, both groups showed an increase in these values at the end of this period (7.11 and 5.62 IgG/ml in NR and C8, respectively, $P < 0.05$). In contrast, the values from the C4 group never decreased, reaching the maximum value at 30 days. Regarding differences among groups, lambs from




Figure 2 BW, IgG and IgM concentrations, TCA and ACA evolution in the NR ($n = 20$), C4 ($n = 20$) and C8 ($n = 20$) groups during the weaning period. NR = natural rearing; C4 = 4 g of IgG/kg of BW at birth; C8 = 8 g of IgG/kg of BW at birth. TCA = total complement activity; ACA = alternative complement activity. Data shown in this figure can be viewed in the Supplementary Table S2.

the NR and C8 group exhibited greater IgG concentrations than animals from the C4 group at the beginning of weaning period (5.22, 2.18, 5.62 mg/ml in NR, C4 and C8 groups, respectively, $P < 0.05$). The consequences of a low colostrum administration, in the case of C4, may have probably persisted at that time. In spite of this fact, no differences among groups were observed during the rest of the weaning

period. As described above, the reduction of colostrum intake in C4 could produce an earlier IgG synthesis compared to the NR and C8 groups, as was described by Nonnecke *et al.* (2012). This fact could explain the increase of IgG in lambs from C4 group during this period.

Conversely to the evolution observed in IgG concentration, IgM concentrations increased in the NR, C4 and C8 groups

during the entire period ($P < 0.05$). At the beginning of weaning (0 and 5 days), no differences were observed among groups, probably because at that time lambs from all the studied groups (not only C4 group) were able to produce IgM, the earliest immunoglobulin synthesized by the immune system (Ehrenstein and Notley, 2010). In addition, and in spite of no differences being observed among groups at 15 days, C4 and C8 groups showed greater values than the NR group during the rest of the period. It has been demonstrated that stress produces a decrease in IgM concentrations in the blood (Matos-Gomes *et al.*, 2010). The stress produced by dam separation in the NR group probably caused the differences observed among groups during the end of the weaning period.

During weaning, all the groups showed an increase in both complement system pathways (TCA and ACA), although different evolutions were observed. In general, the C8 group showed greater complement system activity (TCA and ACA) than the NR and C4 groups during the entire period ($P < 0.05$). Nevertheless, all the studied groups showed a similar TCA (65.06%, 60.05% and 78.10% of hemolysis in NR, C4 and C8 groups, respectively, $P < 0.05$) and ACA (55.40%, 50.77% and 64.65% of hemolysis in NR, C4 and C8 groups, respectively, $P < 0.05$) at the end of the weaning period. Unfortunately, the complement system activity during weaning has not been deeply described in ruminants (cow, sheep and goat). Nevertheless, Oswald *et al.* (1990) described the complement system activity (TCA and ACA) in lambs, showing a similar evolution to present results, although different values were described, probably due to differences in the analytical techniques used.

In conclusion, lambs that received an amount of colostrum equivalent to 8 g of IgG/kg of BW equally divided into three meals at 2, 14 and 24 h after birth were able to reach similar IgG and greater IgM concentrations in blood than lambs reared under natural conditions during the first few days after birth. Lambs from the NR system obtained greater complement system activity than those from the artificial rearing system during the same period. At the end of weaning (60.70, 65.20, 65.45 days after birth in NR, C4 and C8 groups, respectively), lambs from the artificial rearing system showed greater BW and IgM concentrations and similar IgG concentrations and complement system activity compared to lambs reared under natural conditions. However, this study shows that not only the colostrum amount but also the management during the milk feeding and weaning periods, such as stress produced by dam separation or milk quality and suckling frequency, can affect the final immune status of lambs.

This study reveals important information about the amount of colostrum required by newborn lambs reared under artificial conditions in order to reach similar immune values to those reared under natural conditions.

Acknowledgments

Author L.E. Hernández-Castellano acknowledges the financial support from the *Formación del Profesorado Universitario* (FPU)

program (Ministry of Education, Madrid, Spain). The authors also acknowledge Heather Robertson for her help in the grammar revision of this manuscript.

Supplementary material

For supplementary material referred to in this article, please visit <http://dx.doi.org/10.1017/S175173111500110X>

References

- Abdou H, Marichatou H, Beckers J-F, Dufrasne I and Hornick J-L 2014. Effect of bovine colostrum administration on plasma protein profile, growth, and survival in Red kid. *Small Ruminant Research* 117, 158–164.
- Ahmad R, Khan A, Javed MT and Hussain I 2000. The level of immunoglobulins in relation to neonatal lamb mortality in Pak-Karakul sheep. *Veterinarski Arhiv* 70, 129–139.
- al-Sabbagh TA, Swanson LV and Thompson JM 1995. The effect of ewe body condition at lambing on colostrum immunoglobulin G concentration and lamb performance. *Journal of Animal Science* 73, 2860–2864.
- Argüello A, Castro N, Capote J, Tyler JW and Holloway NM 2004. Effect of colostrum administration practices on serum IgG in goat kids. *Livestock Production Science* 90, 235–239.
- Argüello A, Castro N, Capote J, Gines R, Acosta F and Lopez JL 2003. Effects of refrigeration, freezing-thawing and pasteurization on IgG goat colostrum preservation. *Small Ruminant Research* 48, 135–139.
- Bendixen E, Danielsen M, Hollung K, Gianazza E and Miller I 2011. Farm animal proteomics – a review. *Journal of Proteomics* 74, 282–293.
- Brujeni GN, Jani SS, Alidadi N, Tabatabaei S, Sharifi H and Mohri M 2010. Passive immune transfer in fat-tailed sheep: evaluation with different methods. *Small Ruminant Research* 90, 146–149.
- Castro N, Capote J, Alvarez S and Argüello A 2005. Effects of lyophilized colostrum and different colostrum feeding regimens on passive transfer of immunoglobulin G in Majorera goat kids. *Journal of Dairy Science* 88, 3650–3654.
- Castro N, Capote J, Bruckmaier RM and Argüello A 2011. Management effects on colostrum transfer in small ruminants: a review. *Journal of Applied Animal Research* 39, 85–93.
- Castro N, Acosta F, Nino T, Vivas J, Quesada E, Capote J and Argüello A 2008. The effects of diet and age on serum complement system activity in goat kids. *Livestock Science* 119, 102–106.
- Christley RM, Morgan KL, Parkin TDH and French NP 2003. Factors related to the risk of neonatal mortality, birth-weight and serum immunoglobulin concentration in lambs in the UK. *Preventive Veterinary Medicine* 57, 209–226.
- Ciupersescu DD 1977. Dynamics of serum immunoglobulin concentrations in sheep during pregnancy and lactation. *Research in Veterinary Science* 22, 23–27.
- da Nobrega JE, Riet-Correa F, Nobrega RS, de Medeiros JM, de Vasconcelos JS, Simoes SVD and Tabosa IM 2005. Perinatal mortality of lambs in the semi-arid region of Paraíba, Brazil. *Pesquisa Veterinária Brasileira* 25, 171–178.
- David I, Bouvier F, Ricard E, Ruesche J and Weisbecker J-L 2014. Feeding behaviour of artificially reared Romane lambs. *Animal* 8, 982–990.
- Demirenen E, Shrestha JNB and Boylan WJ 1995. Breed and environmental effects on components of ewe productivity in terms of multiple births, artificial rearing and 8-month breeding cycles. *Small Ruminant Research* 16, 239–249.
- Ehrenstein MR and Notley CA 2010. The importance of natural IgM: scavenger, protector and regulator. *Nature Reviews Immunology* 10, 778–786.
- Emsen E, Yaprak M, Bilgin OC, Emsen B and Ockerman HW 2004. Growth performance of Awassi lambs fed calf milk replacer. *Small Ruminant Research* 53, 99–102.
- Firat A, Ozpınar A and Erhard MH 2003. Serum immunoglobulin G levels in lambs fed colostrum and dam milk or cow milk after birth. In *The 11th International Symposium of the World Association of Veterinary Laboratory Diagnosticians and OIE Seminars on Biotechnology* (ed. R. Thanawongnuwech and P. Ingkaninun), pp 24–25. The Thai Association of Veterinary Laboratory Diagnosticians, Bangkok, Thailand.

- Halliday R and Williams MR 1979. Absorption of immunoglobulin from colostrum by bottle-fed lambs. *Annales de Recherches Veterinaires* 10, 549–556.
- Hernández-Castellano LE, Almeida AM, Castro N and Argüello A 2014a. The colostrum proteome, ruminant nutrition and immunity: a review. *Current Protein and Peptide Science* 15, 64–74.
- Hernández-Castellano LE, Argüello A, Almeida AM, Castro N and Bendixen E 2015a. Colostrum protein uptake in neonatal lambs examined by descriptive and quantitative liquid chromatography-tandem mass spectrometry. *Journal of Dairy Science* 98, 135–147.
- Hernández-Castellano LE, Almeida AM, Ventosa M, Coelho AV, Castro N and Argüello A 2014b. The effect of colostrum intake on blood plasma proteome profile in newborn lambs: low abundance proteins. *BMC Veterinary Research* 10, 85.
- Hernández-Castellano LE, Morales-delaNuez A, Sánchez-Macias D, Moreno-Indias I, Torres A, Capote J, Argüello A and Castro N 2015b. The effect of colostrum source (goat vs. sheep) and timing of the first colostrum feeding (2 h vs. 14 h after birth) on body weight and immune status of artificially reared newborn lambs. *Journal of Dairy Science* 98, 204–210.
- Hernández-Castellano LE, Moreno-Indias I, Morales-delaNuez A, Sánchez-Macias D, Torres A, Capote J, Argüello A and Castro N 2015c. The effect of milk source on body weight and immune status of lambs. *Livestock Science* 175, 70–76.
- Holloway NM, Tyler JW, Lakritz J, Carlson SL, Tessman RK and Holle J 2002. Serum immunoglobulin G concentrations in calves fed fresh colostrum or a colostrum supplement. *Journal of Veterinary Internal Medicine* 16, 187–191.
- Lérias JR, Hernández-Castellano LE, Suárez-Trujillo A, Castro N, Poulis A and Almeida AM 2014. The mammary gland in small ruminants: major morphological and functional events underlying milk production – a review. *Journal of Dairy Research* 81, 304–318.
- Matos-Gomes N, Katsurayama M, Makimoto FH, Santana LLO, Paredes-Garcia E, Becker MAD and Dos-Santos MC 2010. Psychological stress and its influence on salivary flow rate, total protein concentration and IgA, IgG and IgM titers. *Neuroimmunomodulation* 17, 396–404.
- Morales-delaNuez A, Castro N, Moreno-Indias I, Juste MC, Sánchez-Macias D, Briggs H, Capote J and Argüello A 2009. Effects of a reputed immunostimulant on the innate immune system of goat kids. *Small Ruminant Research* 85, 23–26.
- Moreno-Indias I, Morales-delaNuez A, Hernandez-Castellano LE, Sanchez-Macias D, Capote J, Castro N and Arguello A 2012. Docosahexaenoic acid in the goat kid diet: effects on immune system and meat quality. *Journal of Animal Science* 90, 3729–3738.
- Moretti DB, Kindlein L, Pauletti P and Machado-Neto R 2010. IgG absorption by Santa Ines lambs fed Holstein bovine colostrum or Santa Ines ovine colostrum. *Animal* 4, 933–937.
- Muller LD and Ellinger DK 1981. Colostral immunoglobulin concentrations among breeds of dairy-cattle. *Journal of Dairy Science* 64, 1727–1730.
- Murphy TA, Loerch SC, McClure KE and Solomon MB 1994. Effects of restricted feeding on growth performance and carcass composition of lambs. *Journal of Animal Science* 72, 3131–3137.
- Napolitano F, Pacelli C, Girolami A and Braghieri A 2008. Effect of information about animal welfare on consumer willingness to pay for yogurt. *Journal of Dairy Science* 91, 910–917.
- Napolitano F, Cifuni GF, Pacelli C, Riviezi AM and Girolami A 2002. Effect of artificial rearing on lamb welfare and meat quality. *Meat Science* 60, 307–315.
- Nonnecke BJ, Waters WR, Goff JP and Foote MR 2012. Adaptive immunity in the colostrum-deprived calf: response to early vaccination with *Mycobacterium bovis* strain bacille Calmette Guerin and ovalbumin. *Journal of Dairy Science* 95, 221–239.
- Nowak R and Poindron P 2006. From birth to colostrum: early steps leading to lamb survival. *Reproduction Nutrition Development* 46, 431–446.
- O'Doherty JV and Crosby TF 1997. The effect of diet in late pregnancy on colostrum production and immunoglobulin absorption in sheep. *Journal of Animal Science* 64, 87–96.
- Ontsouka CE, Bruckmaier RM and Blum JW 2003. Fractionized milk composition during removal of colostrum and mature milk. *Journal of Dairy Science* 86, 2005–2011.
- Oswald IP, Lantier F and Bourgy G 1990. Classical and alternative pathway hemolytic activities of ovine complement – variations with age and sex. *Veterinary Immunology and Immunopathology* 24, 259–266.
- Quigley JD, French P and James RE 2000. Effect of pH on absorption of immunoglobulin G in neonatal calves. *Journal of Dairy Science* 83, 1853–1855.
- Quigley JD 3rd, Kost CJ and Wolfe TM 2002. Absorption of protein and IgG in calves fed a colostrum supplement or replacer. *Journal of Dairy Science* 85, 1243–1248.
- Ramírez-Santana C, Pérez-Cano FJ, Audí C, Castell M, Moretones MG, López-Sabater MC, Castellote C and Franch A 2012. Effects of cooling and freezing storage on the stability of bioactive factors in human colostrum. *Journal of Dairy Science* 95, 2319–2325.
- Rodríguez C, Castro N, Capote J, Morales-delaNuez A, Moreno-Indias I, Sanchez-Macias D and Arguello A 2009. Effect of colostrum immunoglobulin concentration on immunity in Majorera goat kids. *Journal of Dairy Science* 92, 1696–1701.
- Smith GW and Foster DM 2007. Short communication: absorption of protein and immunoglobulin G in calves fed a colostrum replacer. *Journal of Dairy Science* 90, 2905–2908.
- Trujillo AJ, Castro N, Quevedo JM, Arguello A, Capote J and Guamis B 2007. Effect of heat and high-pressure treatments on microbiological quality and immunoglobulin G stability of caprine colostrum. *Journal of Dairy Science* 90, 833–839.
- Wheeler TT, Hodgkinson AJ, Prosser CG and Davis SR 2007. Immune components of colostrum and milk – a historical perspective. *Journal of Mammary Gland Biology and Neoplasia* 12, 237–247.
- Winter A 2011. Bovine neonatal pancytopenia, and anaemia in lambs caused by feeding cow colostrum. *Veterinary Record* 168, 84–84.