



## Effects of EDTA-Induced Hypocalcaemia and Stress on Plasma TNF- $\alpha$ , IL-1-ra, G-CSF, GM-CSF and S-100 in Dairy Cows

J.-L. Riond<sup>1\*</sup>, A. Liesegang<sup>1</sup>, M. Wanner<sup>1</sup>, C. Kaiser<sup>2</sup>, M. Döbeli<sup>2</sup> and H.I. Joller-Jemelka<sup>3</sup>

<sup>1</sup>*Institute of Animal Nutrition and* <sup>2</sup>*Clinic for Andrology and Gynecology, Veterinary Hospital, University of Zürich;* <sup>3</sup>*Division of Clinical Immunology, Department of Internal Medicine, University Hospital of Zürich, Switzerland*

\*Correspondence: *Institute of Animal Nutrition, University of Zürich, Winterthurerstrasse 260, CH-8057 Zürich, Switzerland*

Riond, J.-L., Liesegang, A., Wanner, M., Kaiser, C., Döbeli, M. and Joller-Jemelka, H.I., 1999. Effects of EDTA-induced hypocalcaemia and stress on plasma TNF- $\alpha$ , IL-1-ra, G-CSF, GM-CSF and S-100 in dairy cows. *Veterinary Research Communications*, **23**(5), 299–306

### ABSTRACT

The pathophysiology of postparturient paresis is still not completely understood. Knowledge recently acquired in immunology, endocrinology and cell physiology has still to be integrated in order to elucidate the aetiopathogenesis of the disease. For that purpose, the effect of the EDTA infusion model on the plasma concentrations of selected cytokines and growth factors, and of a calcium binding protein was examined in dairy cows. Six 6- to 11-year-old Brown Swiss cows in mid lactation were infused with a 5% solution of Na<sub>2</sub>EDTA in one jugular vein over a period of 5 h. Blood samples were collected from the contralateral side daily two days before, and then hourly for five hours during the infusion, hourly for five hours after the end of the infusion, and once daily for 10 days thereafter. The plasma concentrations of cortisol, tumour necrosis factor- $\alpha$ , interleukin-1 receptor antagonist, granulocyte colony-stimulating factor, granulocyte and macrophage colony-stimulating factor, and the calcium binding protein S-100 were determined. Before the EDTA infusion, during the infusion and for two days thereafter, the mean plasma concentrations of cortisol were significantly higher than those from days 4 to 10 after the infusion. The plasma concentrations of tumour necrosis factor- $\alpha$  and interleukin-1 receptor antagonist followed a similar profile. At the end of EDTA infusion, low concentrations of granulocyte colony-stimulating factor were detected in one cow only. On days 3 and 4, the mean plasma concentrations of granulocyte colony-stimulating factor were significantly higher than the pre-infusion values, but this was followed by a significant decrease on post-infusion day 5. From day 4 to 7, the plasma concentrations of S-100 were significantly lower than the pre-infusion values. The importance of these findings in the pathophysiology of postparturient paresis remains to be established.

**Keywords:** calcium binding protein; colony-stimulating factor, cortisol, cow, cytokines, ethylenediaminetetraacetate, interleukin receptor antagonist, stress, tumour necrosis factor

**Abbreviations:** S-100, calcium binding protein S-100; EDTA, ethylenediaminetetraacetate; GM-CSF, granulocyte and macrophage colony-stimulating factor; G-CSF, granulocyte colony-stimulating factor; IL-1-ra, interleukin-1 receptor antagonist; TNF- $\alpha$ , tumour necrosis factor- $\alpha$

## INTRODUCTION

Postparturient hypocalcaemia of dairy cows is a commonly encountered disease with a complex pathogenesis (Goff *et al.*, 1991; Horst *et al.*, 1997). Many aspects of its pathophysiology have been investigated with the EDTA infusion model (van de Braak *et al.*, 1984; Desmecht *et al.*, 1995, 1996; Riond *et al.*, 1997). However, knowledge recently acquired in immunology, endocrinology and cell physiology has still to be integrated to fully elucidate the aetiopathogenesis of the disease. Furthermore, the immune system is suppressed during the periparturient period (Kehrli and Goff, 1989). Many cytokines and growth factors are important modulators of the immune responses and are also regulators of a variety of organ functions that are pivotal for calcium homeostasis, for example, bone remodelling (Mundy, 1993). A possible involvement of cytokines, growth factors and calcium binding proteins in the pathophysiology of postparturient hypocalcaemia has not been examined so far. Extensive reviews exist of the biological regulatory activities of tumour necrosis factor- $\alpha$  (Aggarwal and Natarajan, 1996; Bazzoni and Beutler, 1996; Hill and Lunec, 1996), interleukin-1 receptor antagonist (Nikolic-Paterson *et al.*, 1996; Mantovani *et al.*, 1996), granulocyte colony-stimulating factor (Hansson and Söderström, 1993; Metcalf, 1997), granulocyte and macrophage colony-stimulating factor, and the calcium binding protein S-100 (Zimmer *et al.*, 1995; Schäfer and Heizmann, 1996). The purpose of this study was to investigate the effects of EDTA-induced hypocalcaemia on the serum concentrations of TNF- $\alpha$ , IL-1-ra, G-CSF, GM-CSF and S-100.

## MATERIALS AND METHODS

### *Animals and treatments*

Six Brown Swiss cows, aged 6 to 11 years, in mid-lactation, weighing  $610 \pm 22$  (SD) kg and with an average daily milk production of  $22.5 \pm 2.0$  L, were used. The animals were housed in a closed barn and were fed 30 kg of a mixture of 2/3 grass silage, 1/6 corn silage, 1/6 beet chips, and 3 kg of concentrate in two daily portions at 06:00 and 17:00. The daily ration provided 100.1 MJ net energy for lactation, 1359 g absorbable protein in the intestine, 114 g calcium, 55 g phosphorus and 28 g magnesium. The cows were milked between 06:00 and 06:30 and between 17:00 and 17:30. After 2 days of acclimatization to a new place within the barn, a 5% solution of disodium ethylenediaminetetraacetate ( $\text{Na}_2\text{EDTA}$ ) was infused via a catheter in the right jugular vein at 0.90 mmol/min for 5 h. Blood was collected from the contralateral side via a catheter into blood tubes that contained potassium oxalate as an anticoagulant and sodium fluoride as a glycolytic inhibitor, daily 2 days before, shortly after and then hourly for 5 h during the infusion and hourly for 5 h after the end of the infusion. The catheters were implanted 1 day before the infusion. Further blood samples were collected twice daily on days 1 and 2 and then daily for 8 days thereafter. The blood was subsequently centrifuged at 1580g for 10 min at 4°C and the plasma was aspirated. The samples were then stored at 4°C and analysed within 1 week.

### *Biochemical analyses*

The concentration of cortisol in plasma was determined by a radioimmunoassay (RIA) as previously described (Bahr *et al.*, 1998). The sample (100  $\mu$ l) was extracted in duplicate with ethanol (2.0 ml) for 30 min. The tubes with the precipitated proteins in them were centrifuged at 4°C for 15 min and the ethanolic supernatants were decanted into incubation tubes. After evaporation of the organic solvent with nitrogen, the residues were reconstituted in 300  $\mu$ l of assay buffer. Immunoreactive cortisol titres were determined by a cortisol RIA that included an iodinated tracer and a double-antibody separation of bound and unbound ligand, and phosphate-buffered saline (0.01 mol/L, pH 7.0) with 0.1% gelatin as the assay buffer.

The concentrations of G-CSF, GM-CSF and IL-1-ra were determined using commercially available enzyme immunoassays (Quantikine Kit from R&D Systems, Minneapolis, MN, USA). The quantity of TNF- $\alpha$  was measured by an enzyme-linked immunosorbent assay (Bioassay TNF- $\alpha$  US ELISA Kit ultrasensitive, NBS Biologicals Ltd, Cambs, UK) and S-100 by a luminescence-immunoassay (LIA-mat Sangtec-100, BYK-Sangtec Diagnostica GmbH, Germany).

### *Statistical analyses*

The changes with time in the concentrations of cortisol, TNF- $\alpha$ , IL-1-ra, G-CSF, GM-CSF and S-100 were examined by repeated measurements analysis of variance using version 6.11 of the mainframe computer-implemented SAS procedure GLM (SAS, 1989). For each parameter, the values of the concentrations during and after the infusion were compared with either the first or second pre-infusion value by use of the option *contrast* and adjacent levels of the parameters were compared by use of the option *profile*. The level of significance was fixed at  $p < 0.05$ .

## RESULTS

The pathophysiological effects and the changes in the serum concentrations of total and ionized calcium and inorganic phosphate induced by the EDTA infusion have been reported elsewhere (Riond *et al.*, 1997). The data (mean  $\pm$  SD) for cortisol, cytokines, growth factors and S-100 are presented in Figure 1A–F. During the days before the EDTA infusion, during the infusion and for 2 days thereafter, the mean plasma concentrations of cortisol were significantly higher than those from day 4 to day 10 after the infusion (Figure 1A). The SAS option *contrast* indicated a significant decrease in cortisol concentrations by day 3. The plasma concentrations of TNF- $\alpha$  and IL-1-ra followed a similar profile with significantly lower concentrations on post-infusion day 5 (Figure 1B–C). The SAS option *profile* revealed decreasing concentrations on day 2 for TNF- $\alpha$  and in the first part of day 1 for IL-1. During the last 2 h of the EDTA infusion, low concentrations of G-CSF were detected in one cow (Figure 1D). On days 3 and 4, the mean serum concentrations of G-CSF were significantly




Figure 1. The effects of EDTA infusion and stress on the plasma concentrations (means  $\pm$  SD with minima and maxima shown as dotted lines) of (A) cortisol, (B) TNF- $\alpha$ , (C) IL-1-ra, (D) G-CSF, (E) GM-CSF and (F) S-100 in six dairy cows




Figure 1. See legend on opposite page

higher than the pre-infusion values, but this was followed by a significant decrease on day 5. No significant changes with time were observed in the concentrations of GM-CSF (Figure 1E). From days 4 to 7, the serum concentrations of S-100 were significantly lower than the pre-infusion values (Figure 1F).

## DISCUSSION

As the plasma concentrations of TNF- $\alpha$  and IL-1-ra, which were not directly related to the EDTA infusion, were unexpectedly elevated during the first part of the experiment, the cortisol concentrations were measured retrospectively. Psychological stress is the most likely cause of the elevated plasma concentrations of TNF- $\alpha$ , IL-1-ra and cortisol following activation of the sympathetic/adrenomedullary system with subsequent systemic release of the inflammatory cytokines IL-6, IL-1 and TNF- $\alpha$  (Bellinger *et al.*, 1992; Stratakis and Chrousos, 1995). The inflammatory cytokines are known to activate the hypothalamic-pituitary-adrenal axis, with subsequent elevation of the concentrations of ACTH and cortisol (Besedovsky *et al.*, 1986; Sapolsky *et al.*, 1987; Perlstein *et al.*, 1993; Woloski *et al.*, 1985; Mastorakos *et al.*, 1993). Glucocorticoids in turn suppress the stimulatory effects of cytokines on the hypothalamic-pituitary-adrenal axis (Lázár *et al.*, 1992; Besedovsky *et al.*, 1986; Beutler *et al.*, 1986; Dubois *et al.*, 1989). In the present study, the concentrations of IL-1 and IL-6 were not determined. The elevated plasma concentrations of IL-1-ra before, during and for 4 days after the infusion may be explained by the fact that increased production of IL-1-ra is usually associated with an increased production of IL-1 (Mantavoni *et al.*, 1996; Nikolic-Paterson *et al.*, 1996). The source of the stress for these cows was probably their accommodation in a new place in the barn and/or new people. In contrast to the present study, cortisol concentrations increased significantly in association with EDTA infusion when the EDTA was infused at twice the rate (Desmecht *et al.*, 1996) and cortisol concentrations were also significantly increased in postparturient cows with severe hypocalcaemia (Horst and Jorgensen, 1982). To properly evaluate the degree of the stress, it would be better to make serial determinations of cortisol plasma concentrations in the course of the day because episodic secretion, inducing large fluctuations of cortisol concentrations, occurs in cattle (Thun *et al.*, 1981; Thun, 1986, 1987). Furthermore, a pronounced circadian rhythm has been observed for plasma cortisol. In the present study, TNF- $\alpha$  and IL-1-ra concentrations decreased significantly before the cortisol concentrations did so. This observation confirms the controlling role of these two cytokines on the adrenomedullary axis.

The decreased G-CSF serum concentrations following the EDTA infusion may be the consequence of hypocalcaemia or hypophosphataemia or of the chelation of other ions, such as zinc. The relatively small volume of the carrier saline solution in comparison to the volume of the extracellular space is unlikely to have induced changes in the extracellular fluid that would explain such a change in the concentration of G-CSF. At the end of the infusion, the production of G-CSF by monocytes and endothelial cells was possibly decreased or their elimination may have been enhanced. The importance of this finding in the pathophysiology of postparturient paresis of dairy cows remains to be established.

The period over which S-100 concentrations were significantly decreased corresponds to the beginning of the period with low concentrations in the serum of cortisol, TNF- $\alpha$  and IL-1-ra. This observation suggests a transient effect from low concentrations of cortisol, TNF- $\alpha$  and/or IL-1-ra which lasts only until a readjustment has occurred, following feedback mechanisms.

## CONCLUSION

Kits developed for the determination of human TNF- $\alpha$ , IL-1-ra, G-CSF, GM-CSF and S-100 in plasma may be used for cattle. The observations on the changes in the plasma concentrations of cortisol, TNF- $\alpha$ , IL-1-ra, G-CSF, GM-CSF and S-100 in relation to the infusion of EDTA may be useful for the design of future trials, which are still needed for a better understanding of the pathophysiology of postparturient paresis in dairy cows. This study strongly suggests an involvement of TNF- $\alpha$  and IL-1-ra in the stress response of cows. These two cytokines could be used as markers for stress in cattle. Also, the importance of stress as a confounding factor should not be neglected in this type of investigation. Finally, the role of G-CSF in postparturient paresis remains to be established.

## ACKNOWLEDGEMENTS

This study was supported by the Swiss National Science Foundation grants numbers 32-33853.92 and 32-46791.96 and by funds from the canton of Zurich. The assistance of Mrs Marianne Mathys in the preparation of the Figures was greatly appreciated.

## REFERENCES

- Aggarwal, B.B. and Natarajan, K., 1996. Tumor necrosis factors: developments during the last decade. *European Cytokine Network*, **7**, 93–124
- Bahr, N.I., Price, C.R., Döbeli, M. and Martin, R.D., 1998. Evidence from urinary cortisol that maternal behavior is related to stress in gorillas. *Physiology and Behavior*, **64**, 429–437
- Bazzoni, F. and Beutler, B., 1996. The tumor necrosis factor ligand and receptor families. *New England Journal of Medicine*, **334**, 1717–1725
- Bellinger, D.L., Lorton, D., Felten, S.Y. and Felten, D.L., 1992. Innervation of lymphoid organs and implications in development, aging, and autoimmunity. *International Journal of Immunopharmacology*, **14**, 329–344
- Besedovsky, H., del Rey, A., Sorkin, E. and Dinarello, C.A., 1986. Immunoregulatory feedback between interleukin-1 and glucocorticoid hormones. *Science (Washington DC)*, **233**, 652–654
- Beutler, B., Krochin, N., Milsark, I.W., Luedke, C. and Cerami, A., 1986. Control of cachectin (tumor necrosis factor) synthesis: mechanisms of endotoxin resistance. *Science (Washington DC)*, **232**, 977–980
- Desmecht, D.J.M., Linden, A.S., Godeau, J.M. and Lekeux, P.M., 1995. Experimental production of hypocalcaemia by EDTA infusion in calves: a critical appraisal assessed from the profile of blood chemicals and enzymes. *Comparative Biochemical Physiology A*, **110**, 115–130
- Desmecht, D.J.M., Linden, A.S. and Lekeux, P.M., 1996. Ruminal, cardiorespiratory and adrenocortical sequelae of Na<sub>2</sub>EDTA-induced hypocalcaemia in calves. *Veterinary Research Communications*, **20**, 47–60
- Dubois, C., Bissonnette, E. and Rola-Pleszczynski, M., 1989. Platelet-activating factor (PAF) enhances tumor necrosis factor production by alveolar macrophages. Prevention by PAF receptor antagonists and lipooxygenase inhibitors. *Journal of Immunology*, **143**, 964–970

- Goff, J.P., Reinhardt, T.A. and Horst, R.L., 1991. Enzymes and factors controlling vitamin D metabolism and action in normal and milk fever cows. *Journal of Dairy Science*, **74**, 4022–4032
- Hansson, M. and Söderström, T., 1993. The colony stimulating factor. *Medical Oncology and Tumor Pharmacotherapy*, **10**, 5–12
- Hill, C.M. and Lunec, J., 1996. The TNF-ligand and receptors superfamilies: controllers of immunity and the Trojan horses of autoimmune disease? *Molecular Aspects of Medicine*, **17**, 455–509
- Horst, R.L., Goff, J.P., Reinhardt, T.A. and Buxton, D.R., 1997. Strategies for preventing milk fever in dairy cattle. *Journal of Dairy Science*, **80**, 1269–1280
- Horst, R.L. and Jorgensen, N.A., 1982. Elevated plasma cortisol during induced and spontaneous hypocalcemia in ruminants. *Journal of Dairy Science*, **65**, 2332–2337
- Kehrli, M.E. and Goff, J.P., 1989. Periparturient hypocalcemia in cows: effects on peripheral blood neutrophil and lymphocyte function. *Journal of Dairy Science*, **72**, 1188–1196
- Lázár, G., Duda, E. and Lázár, G., 1992. Effect of RU 38486 on TNF production and toxicity. *FEBS Letters*, **308**, 137–140
- Mantovani, A., Muzio, M., Ghezzi, P., Colotta, F. and Introna, M., 1996. Negative regulators of the interleukin-1 system: receptor antagonists and a decoy receptor. *International Journal of Clinical and Laboratory Research*, **26**, 7–14
- Mastorakos, G., Chrousos, G.P. and Weber, J.S., 1993. Recombinant interleukin-6 activates the hypothalamic–pituitary–adrenal axis in humans. *Journal of Clinical Endocrinology and Metabolism*, **77**, 1690–1694
- Metcalf, D., 1997. *The Molecular Basis of Cellular Defense Mechanisms*, (Ciba Foundation Symposium 204; Wiley, Chichester), 40–50
- Mundy, G.R., 1993. Cytokines and growth factors in the regulation of bone remodelling. *Journal of Bone and Mineral Research*, **8**(supplement 2), S505–S510
- Nikolic-Paterson, D.J., Lan, H.Y. and Atkins, R.C., 1996. Interleukin-1 receptor antagonism. *Seminars in Nephrology*, **16**, 583–590
- Perlstein, R.S., Whitnall, M.H., Abrams, J.S., Mougey, E.H. and Neta, R., 1993. Synergistic roles of interleukin-6, interleukin-1, and tumor necrosis factor in the adrenocorticotropin response to bacterial lipopolysaccharide *in vivo*. *Endocrinology*, **132**, 946–952
- Riond, J.-L., Liesegang, A., Fakler, A. and Spichiger, U.E., 1997. Serum concentrations of total and ionized calcium and magnesium and inorganic phosphate during Na<sub>2</sub>EDTA infusion in dairy cows. *Journal of Animal Physiology and Animal Nutrition*, **75**, 161–166
- Sapolsky, R., Rivier, C., Yamamoto, G., Plotsky, P. and Vale, W., 1987. Interleukin-1 stimulates the secretion of hypothalamic corticotropin-releasing factor. *Science (Washington DC)*, **238**, 522–524
- SAS (1989). *SAS/STAT User's Guide Version 6*, 4th edn, Vols 1 and 2, (SAS Institute, Cary, NC, USA)
- Schäfer, B.W. and Heizmann, C.W., 1996. The S100 family of EF-hand calcium-binding proteins: functions and pathology. *Trends in Biochemical Sciences*, **21**, 134–140
- Stratakis, C.A. and Chrousos, G.P., 1995. Neuroendocrinology and pathophysiology of the stress system. In: G.P. Chrousos, R. McCarty, K. Pacák, G. Cizza, E. Sternberg, P.W. Gold and R. Kvetnanský (eds), *Stress, Basic Mechanisms and Clinical Implications*, (The New York Academy of Sciences, New York), 1–18
- Thun, R., 1986. Tagesrhythmische Cortisolsekretion vor und nach der Geburt beim Rind. *Fertilität*, **2**, 200–202
- Thun, R., 1987. *Untersuchungen über die Tagesrhythmik von Cortisol beim Rind*, (Ferdinand Enke Verlag, Stuttgart, Germany)
- Thun, R., Eggenberger, E., Zerobin, K., Lüscher, T. and Vetter, W., 1981. Twenty-four-hour secretory pattern of cortisol in the bull: evidence of episodic secretion and circadian rhythm. *Endocrinology*, **109**, 2208–2212
- van de Braak, A.E., van't Klooster, A.T., van Hal-van Gestel, J.C. and Malestein, A., 1984. Influence of stage of lactation and calcium level of the ration on mobilisation rate of calcium and excretion of hydroxyproline in urine in dairy cows. Studies with Na<sub>2</sub>-EDTA infusions in monozygotic twins. *Zentralblatt für Veterinärmedizin Reihe A (Journal of Veterinary Medicine Series A)*, **31**, 725–739
- Woloski, B.M.R.N.J., Smith, E.M., Meyer III, W.J., Fuller, G.M. and Blalock, J.E., 1985. Corticotropin-releasing activity of monokines. *Science (Washington DC)*, **230**, 1035–1037
- Zimmer, D.B., Cornwall, E.H., Landar, A. and Song, W., 1995. The S100 protein family: history, function and expression. *Brain Research Bulletin*, **37**, 417–429