

## Stomatal conductance in mature deciduous forest trees exposed to elevated CO<sub>2</sub>

Sonja Gisela Keel · Steeve Pepin · Sebastian Leuzinger · Christian Körner

Published online: 30 January 2007  
© Springer-Verlag 2007

---

### Erratum to: Trees DOI 10.1007/s00468-006-0106-y

In the last paragraph in the Discussion section on page 8, starting: “In conclusion”, two sentences are missing (in bold). The whole passage should read as follows:

But based on our and other field data (those for conifers in particular) and accounting for atmospheric feedback, we believe that theory based signals are likely to overestimate actual effects substantially. **Furthermore, gas exchange**

**theory (Farquhar and Wong 1984) does not imply species specificity. However, the trends observed in leaf conductance and leaf water potential both point to a great sensitivity of *Carpinus* to elevated CO<sub>2</sub>, and no leaf level response in *Fagus* and *Quercus*.** In the long run, such species specific differences may lead to a change in species abundance driven by soil moisture (and nutrient) effects.

---

The online version of the original article can be found at  
<http://dx.doi.org/10.1007/s00468-006-0106-y>

---

S. G. Keel (✉)  
Laboratory of Atmospheric Chemistry, Paul Scherrer Institute,  
CH-5232 Villigen, Switzerland  
e-mail: sonja.keel@psi.ch

S. Pepin · S. Leuzinger · C. Körner  
Institute of Botany, University of Basel,  
Schönbeinstrasse 6,  
CH-4056 Basel, Switzerland

*Present address:*  
S. Pepin  
Département des sols et de génie agroalimentaire, Université  
Laval, Pavillon Paul-Comtois,  
Québec (Québec), Canada, G1K 7P4