

Computational Fluid Dynamics: Hemodynamic Changes in Abdominal Aortic Aneurysm After Stent-Graft Implantation

Thomas Frauenfelder,¹ Mourad Lotfey,² Thomas Boehm,¹ Simon Wildermuth¹

¹Institute of Diagnostic Radiology, University Hospital of Zurich, Zurich, Switzerland

²Fluent Corp., Darmstadt, Germany

RE: *Cardiovasc Intervent Radiol* (2006) 29: 613–623 (this issue) — Several errors appear in the published online and print versions of this article in Table 3. In the table, the top heading on the right side should read “1-EIA-ratio per cardiac cycle.” In column 3, row 1, the correct number is 0.75. In column 3, row 2, the correct number is 0.95. The complete Table 3 appears correctly below.

Table 3. Flow volume ratio of the right versus the left EIA

	EIA-ratio per cardiac cycle		1-EIA-ratio per cardiac cycle	
	Before	After	Before	After
	Aortic stent-graft implantation		Aortic stent-graft implantation	
Patient 1	0.68	0.75	0.318	0.246
Patient 2	0.66	0.95	0.33	0.05
Patient 3	1.28	0.86	0.273	0.241
Patient 4	1.06	0.81	0.057	0.189
Patient 5	0.47	0.58	0.522	0.418
Patient 6	1.05	1.27	0.044	0.272
Patient 7	1.23	0.83	0.237	0.181
Patient 8	1.24	0.96	0.242	0.039
Patient 9	1.27	0.84	0.272	0.156
Patient 10	1.27	0.80	0.274	0.197
Patient 11	0.42	1.04	0.579	0.137