

Andreas Gebhardt  
Robert Buehler  
Roland Wiest  
Friedemann Tewald  
Johann Sellner  
Sebastian Humpert  
Heinrich P. Mattle  
Alain Kaelin-Lang

## ***Mycoplasma pneumoniae* as a cause of neuromyelitis optica?**

Received: 8 July 2007  
Received in revised form:  
11 December 2007  
Accepted: 7 January 2008  
Published online: 23 June 2008

Sirs: Until recently neuromyelitis optica (NMO) was regarded as a severe form of multiple sclerosis (MS). But characteristic serologic and neuroradiological findings and particularly the discovery of the specific NMO antibody by Lennon and colleagues [8] indicate that NMO is an autoimmune disease different from MS.

In a few patients, typical NMO is associated with viral infection like HIV [3], Dengue fever [10], varicella infection [1], infectious mononucleosis [12] and hepatitis A virus [6]. The only bacterial agent that has been associated with NMO is *Mycobacterium tuberculosis* [2].

---

A. Gebhardt · R. Buehler · J. Sellner ·  
S. Humpert · H. P. Mattle ·  
A. Kaelin-Lang (✉)  
Dept. of Neurology  
University of Bern  
Inselspital  
3010 Bern, Switzerland  
Fax: +41-31/632-2752  
E-Mail: alain.kaelin@dkf.unibe.ch

R. Wiest  
Dept. of Neuroradiology  
University of Bern  
Inselspital  
3010 Bern, Switzerland

F. Tewald  
Labor Enders and Partners  
Rosenbergstrasse 85  
70193 Stuttgart, Germany

On the other hand, several neurological manifestations have been described following *Mycoplasma pneumoniae* [4, 5] infection, a common bacterium of the respiratory tract, including acute transverse myelitis (ATM) [11].

Here we present a patient with typical NMO after *M. pneumoniae* infection that may represent an underestimated etiology for NMO. A previously healthy 32-year-old woman developed fever and a non-productive cough. A chest x-ray showed an infiltrate in the right inferior lobe. Respiratory symptoms improved with erythromycin treatment. Ten days later she experienced paresthesia and weakness in both legs, urinary retention and was admitted to our department.

On neurological examination she was alert. Cranial nerves were normal, but there was tetraparesis with tetrahyperreflexia and hypaesthesia with a sensory level at the 4<sup>th</sup> thoracic level. No signs of meningism were present. Sphincter tone was reduced on rectal examination. The night after admission, the patient had to be transferred to our intensive care unit because of respiratory failure requiring intubation. MRI investigation revealed a severe myelitis (Fig. 1A and B).

CSF investigation showed 15 white blood cells per  $\mu\text{L}$ , 37% polymorphonuclear neutrophils. There was no quantitative intrathecal immunoglobulin production.

Serologic examinations showed a highly positive complement fixation test (CFT) to *M. pneumoniae* (1:5120). An indirect immunofluorescence assay was used to search for *M. pneumoniae* antibodies in CSF and serum. This investigation revealed an elevated titer of IgG antibodies specific to *M. pneumoniae* in the CSF. In combination with an antibody specificity index of 9.5 these results indicate an intrathecal production of IgG antibodies specific to *M. pneumoniae*. Neither in

CSF nor in serum were there signs of another acute infection or autoimmune disorder. NMO-IgG anti-aquaporin-4 antibodies [8] were negative. Within 4 days the tetraparesis progressed to tetraplegia. Sensory level of hypesthesia moved up to C4 and bilateral blindness occurred. An immunosuppressive therapy with methylprednisolone and later with plasma exchange and mitoxantron was initiated in addition to the initial antibiotics therapy. Blindness resolved slowly within the following months. After 1 year, the patient had regained almost full function of both upper extremities. Visual function was normal, but she was still paraspastic and -plegic. CFT to *M. pneumoniae* had decreased to 1:120 and spinal MRI revealed an atrophy (Fig. 1C).

In this case, the development of symptoms after pneumonia and associated increase of *M. pneumoniae* antibody titres both in serum and CSF strongly suggests that the mycoplasma infection is involved in this NMO syndrome.

Several neurological complications including ADEM and ATM have been described after *M. pneumoniae* infection [5, 9]. Both direct CNS infection and an autoimmune mechanism triggered by *M. pneumoniae* are possible. Since direct PCR evidence for *M. pneumoniae* is only seldom found in CSF [11], it is often difficult to differentiate between both possibilities [5]. The clinical syndrome with bilateral optic neuritis and longitudinally extensive myelitis is typical for NMO and, to the best of our knowledge, has not been reported to date in association with *M. pneumoniae* infection. Overall, an autoimmune parainfectious pathogenesis seems more likely than a direct infection since the symptoms started late after the pneumonia, progressed in spite of an adequate antibiotic treatment, and since CSF analysis



**Fig. 1** Spinal MRI. **A** Sagittal T2-weighted TSE images reveal progressive T2-hyperintense and progressive swelling of the spinal cord at the level of C2-Th1 and Th4-Th9 16 days after symptoms onset. **B** T1-weighted SE images show hemorrhagic transformation at the level of Th5–Th9. **C** Follow-up examination after 10 months reveal severe atrophy of the thoracic cord

revealed only weak inflammatory signs in spite of a strong anti-mycoplasma antibodies titre. In this case, the recently published revised diagnostic criteria for NMO are fulfilled [13] even though the anti-aquaporin-4 antibodies [7], which have a 90% specificity for NMO, were not detected in the CSF or blood.

We postulate that this case represents an anti-aquaporin-4 antibody negative NMO syndrome following *M. pneumoniae* infection. We propose that in patients with

NMO syndrome, especially cases with rapid symptom onset, a comprehensive search for possible infectious etiologies including *M. pneumoniae* infection should be performed.

## References

1. Ahasan HA, Rafiqueuddin AK, Chowdhury MA, Azhar MA, Kabir F (1994) Neuromyelitis optica (Devic's disease) following chicken pox. *Trop* 24(2): 757–756

2. Barbizet J, Degos JD, Meyrignac C (1980) Acute neuromyelitis optica and acute pulmonary tuberculosis. *Rev Neurol (Paris)* 136(4):303–309
3. Blanche P, Diaz E, Gombert B, Sicard D, Rivoal O, Brezin A (2000) Devic's neuromyelitis optica and HIV-1 infection. *J Neurol Neurosurg Psychiatry* 68(6): 795–796
4. Behan PO, Feldman RG, Segerra JM, Draper IT (1986) Neurological aspects of mycoplasma infection. *Acta Neurol Scand* 74(4):314–322
5. Guleria R, Nisar N, Chawla TC, Biswas NR (2005) *Mycoplasma pneumoniae* and central nervous system complications: a review. *J Lab Clin Med* 146(2): 55–63
6. Jouhadi Z, Ouazzani I, Abid A, El Moutawakil B, Rafai MA, Slassi I (2004) Devic's optic neuromyelitis and viral hepatitis type A. A paediatric case report. *Rev Neurol (Paris)* 160(12): 1198–1202
7. Lennon VA, Kryzer TJ, Pittock SJ, Verkman AS, Hinson SR (2005) IgG marker of optic-spinal multiple sclerosis binds to the aquaporin-4 water channel. *J Exp Med* 15; 202(4):473–477
8. Lennon VA, Wingerchuk DM, Kryzer TJ, Pittock SJ, Lucchinetti CF, Fujihara K, Nakashima I, Weinshenker BG (2004) A serum autoantibody marker of neuromyelitis optica: distinction from multiple sclerosis. *Lancet* 11–17; 364(9451):2106–2112
9. Menge T, Hemmer B, Nessler S, Wiendl H, Neuhaus O, Hartung HP, Kieseier BC, Stuve O (2005) Acute disseminated encephalomyelitis: an update. *Arch Neurol* 62(11):1673–1680
10. Miranda de Sousa A, Puccioni-Sohler M, Dias Borges A, Fernandes Adorno L, Papais Alvarenga M, Papais Alvarenga RM (2006) Post-dengue neuromyelitis optica: case report of a Japanese-descendent Brazilian child. *J Infect Chemother* 12(6):396–398
11. Tsiodras S, Kelesidis I, Kelesidis T, Stamboulis E, Giamarellou H (2005) Central nervous system manifestations of *Mycoplasma pneumoniae* infections. *J Infect* 51(5):343–354
12. Williamson PM (1975) Neuromyelitis optica following infectious mononucleosis. *Proc Aust Assoc Neurol* 12: 153–155
13. Wingerchuk DM, Lennon VA, Pittock SJ, Lucchinetti CF, Weinshenker BG (2006) Revised diagnostic criteria for neuromyelitis optica. *Neurology* 23; 66(10):1485–1489