

# Numerical methods for Lévy processes

N. Hilber · N. Reich · C. Schwab · C. Winter

Received: 25 March 2008 / Accepted: 30 May 2008 / Published online: 23 July 2009  
© Springer-Verlag 2009

**Abstract** We survey the use and limitations of some numerical methods for pricing derivative contracts in multidimensional geometric Lévy models.

**Keywords** Multidimensional Lévy processes · Numerical methods · Asset pricing

**Mathematics Subject Classification (2000)** 60J75 · 65N06 · 65N30

**JEL Classification** C63

## 1 Introduction

Over the last years financial models with jumps and especially Lévy models have seen a tremendous increase in popularity. By now it is well established that Lévy models are more suitable for capturing market fluctuations than the classical Black–Scholes model [18], see e.g. Cont and Tankov [38] and Schoutens [114] for an overview and empirical justification. The number of financial models with jumps is growing steadily; for the most popular and some recent examples we refer to [12, 32, 33, 49, 74, 86, 87, 114, 116]. However, even in the Black–Scholes setting, analytic solutions to derivative pricing problems are often unavailable or not easily computable, e.g. for American or path-dependent options. Furthermore, in models with jumps one usually cannot even construct analytic solutions for the pricing of plain European vanilla options. Therefore, numerical methods for option pricing have been studied by many authors and several techniques have been developed to obtain efficient pricing algorithms. In particular, models with jumps give rise to previously unconsidered numerical challenges which have led to a number of innovative numerical tools.

---

N. Hilber (✉) · N. Reich · C. Schwab · C. Winter  
Seminar for Applied Mathematics, ETH Zurich, Raemistrasse 101, 8092 Zurich, Switzerland  
e-mail: [norbert.hilber@sam.math.ethz.ch](mailto:norbert.hilber@sam.math.ethz.ch)

In this survey we shall focus on the use and limitations of numerical methods in multidimensional Lévy models for asset pricing. Naturally, there are many other areas where numerical tools can be applied to Lévy processes, e.g. portfolio optimization, but these will be considered elsewhere. In Sects. 2–4 we describe the basic ideas and mathematical background of the most important numerical pricing techniques and further illustrate some of their recent developments. In Sect. 5 we give a qualitative comparison of the described methods and discuss their advantages and shortcomings. For a comprehensive introduction to Lévy models and numerical methods (mainly in dimension  $d = 1$ ) we refer to Cont and Tankov [38].

Throughout this survey, we focus on the pricing of European and American options on  $d \geq 1$  assets with maturity  $T < \infty$  and Lipschitz-continuous payoff  $g(S)$ . Unless stated otherwise, we shall assume that the  $d$ -dimensional underlying is modeled by an exponential Lévy process  $S$  with state space  $\mathbb{R}_{>0}^d$ . The risk-neutral dynamics of  $S = (S^1, \dots, S^d)$  are given by

$$S_t^i = S_0^i e^{rt + X_t^i}, \quad i = 1, \dots, d, \tag{1.1}$$

where  $X$  is an  $\mathbb{R}^d$ -valued Lévy process with characteristic triplet  $(\gamma, Q, \nu)$  under a (non-unique) risk-neutral measure  $\mathbb{P}$  such that  $(e^{X^1}, \dots, e^{X^d})$  is a martingale with respect to the canonical filtration  $\mathcal{F}_t^0 := \sigma(X_s, s \leq t)$ ,  $t \geq 0$ , of the multivariate process  $X$ . By the fundamental theorem of asset pricing (see [46]), an arbitrage-free price  $V(t, S)$  of the European option is given by

$$V(t, S) = \mathbb{E}[e^{-r(T-t)} g(S_T) | S_t = S], \tag{1.2}$$

where the expectation is taken with respect to the risk-neutral measure introduced above.

An arbitrage-free price of the corresponding American-type contract is given by an optimal stopping, free boundary problem, namely

$$V(t, S) = \operatorname{ess\,sup}_{t \leq \tau \leq T} \mathbb{E}[e^{-r(T-\tau)} g(S_\tau) | S_t = S], \tag{1.3}$$

with  $\tau$  ranging over stopping times. Note that the martingale condition implies

$$\int_{|z|>1} e^{z_i} \nu(dz) < \infty, \quad i = 1, \dots, d.$$

We therefore assume that the Lévy measure  $\nu$  of  $X$  admits *semiheavy tails* in the following sense. Let  $\nu_i$ ,  $i = 1, \dots, d$  denote the marginal Lévy measures of the Lévy measure  $\nu$  of  $X$ . Then we assume that there are constants  $\beta_i^- > 0$ ,  $\beta_i^+ > 0$ ,  $i = 1, \dots, d$  such that

$$\int_1^\infty e^{\beta_i^+ z} \nu_i(dz) < \infty, \quad \text{and} \quad \int_{-\infty}^{-1} e^{-\beta_i^- z} \nu_i(dz) < \infty. \tag{1.4}$$

Note that this assumption is satisfied by a wide range of Lévy models, cf. e.g. [91]. By [104, Proposition 3.2], it carries over to the multidimensional case:

**Lemma 1.1** *Let  $X$  be a Lévy process with state space  $\mathbb{R}^d$  and Lévy measure  $\nu$  such that the marginal measures  $\nu_i$  satisfy (1.4). Then the Lévy measure  $\nu$  also decays exponentially, i.e.,*

$$\int_{|z|>1} e^{\eta(z)} \nu(dz) < \infty, \quad \text{with } \eta(z) = \sum_{i=1}^d (\mu_i^+ 1_{\{z_i>0\}} + \mu_i^- 1_{\{z_i<0\}}) |z_i|,$$

and  $0 < \mu_i^- < \frac{\beta_i^-}{d}$  and  $0 < \mu_i^+ < \frac{\beta_i^+}{d}$ ,  $i = 1, \dots, d$ .

To fix notation, we recall some essential definitions and properties of Lévy processes. For an extensive description we refer to the monographs [16, 112].

A càdlàg stochastic process  $\{X_t : t \geq 0\}$  on  $\mathbb{R}^d$  such that  $X_0 = 0$  a.s. is called a Lévy process if it has independent and stationary increments and is stochastically continuous. For the characteristic function  $\Phi_t(\cdot)$  of  $X$  at time  $t \geq 0$  we have the Lévy–Khinchin representation (cf. [112]),

$$\begin{aligned} \Phi_t(\xi) &:= \mathbb{E}[e^{i\langle \xi, X_t \rangle}] = e^{t\psi(\xi)}, \quad \xi \in \mathbb{R}^d, \\ \text{with } \psi(\xi) &= i\langle \gamma, \xi \rangle - \frac{1}{2}\langle \xi, Q\xi \rangle + \int_{\mathbb{R}^d} (e^{i\langle \xi, z \rangle} - 1 - i\langle \xi, z \rangle 1_{\{|z| \leq 1\}}) \nu(dz), \end{aligned} \tag{1.5}$$

where  $Q \in \mathbb{R}^{d \times d}$  denotes the covariance matrix,  $\gamma \in \mathbb{R}^d$  the drift vector and  $\nu$  is the Lévy measure which satisfies

$$\int_{\mathbb{R}^d} (1 \wedge |z|^2) \nu(dz) < \infty. \tag{1.6}$$

The triplet  $(\gamma, Q, \nu)$  is called characteristic triplet of the process  $X$  and  $\psi$  the characteristic exponent.

Also note that the dependence structure of the jump part of a Lévy process  $X$  can be described by a Lévy copula  $F$ . These were introduced in Tankov [116] and developed in Kallsen and Tankov [71]. Lévy copulas are functions  $F : \bar{\mathbb{R}}^d \rightarrow \bar{\mathbb{R}}$  which are finite (except at infinity), grounded,  $d$ -increasing and preserve the margins. There is a Lévy copula associated to each process  $X$  and it satisfies the relationship

$$U(x_1, \dots, x_d) = F(U_1(x_1), \dots, U_d(x_d)), \tag{1.7}$$

where  $U$  denotes the tail integral and  $U_i, i = 1, \dots, d$  are the marginal tail integrals of the Lévy process  $X$ . Therefore, the jump part of a Lévy process is always uniquely defined by its  $d$  univariate marginal processes together with the Lévy copula.

## 2 Monte Carlo methods

One does not make a mistake by saying that the Monte Carlo method is the most used simulation tool in financial engineering practice. The popularity of Monte Carlo stems mainly from its simplicity, the applicability for parallel computing and the

independence of the convergence rate with respect to the dimension of the underlying problem. In the context of derivative pricing under diffusion driven market models, the Monte Carlo method is especially attractive since it is very simple to simulate Brownian motions. However, when market models are extended to Lévy processes, Monte Carlo simulation becomes more subtle since the laws of the increments of Lévy processes are in general not known explicitly. This fact becomes even more pronounced in  $d > 1$  dimensions. As a consequence, the paths of most Lévy processes can only be simulated approximately, see e.g. [9, 38, 114].

A Monte Carlo approximation  $\tilde{V}$  of the price  $V$  of a European option in (1.2) with  $t = 0$  consists of the following three steps:

1. Let  $N \in \mathbb{N}$ . For  $j = 1, \dots, N$  simulate the realizations  $\tilde{X}_{T,j}$  of the log-stock price process  $X$  at maturity.
2. For each realization  $\tilde{X}_{T,j}$ , evaluate the payoff

$$g_j := g(S_0^1 e^{rT + \tilde{X}_{T,j}^1}, \dots, S_0^d e^{rT + \tilde{X}_{T,j}^d}).$$

3. Take the discounted mean of  $g_1, \dots, g_N$  to obtain  $\tilde{V} = e^{-rT} N^{-1} \sum_{j=1}^N g_j$ .

The key in the above algorithm is the first step. The simulation of  $\tilde{X}_{T,j}$  is an easy task as long as the law of the increments of  $X$  is known explicitly. However in dimension  $d > 1$ , except for subordinated Brownian motion, explicit formulas to simulate the increments of multidimensional processes are not available in general and one has to rely on approximation methods. We discuss Gaussian approximation, where the small jumps of the Lévy process are approximated by a Brownian motion, as well as series expansions.

*Subordinated Brownian motion* A popular class of processes is obtained by subordinating a Brownian motion with drift with an independent positive Lévy process  $G$ . Then the resulting process is given by

$$X_t = \Sigma W_{G_t} + \gamma G_t, \quad \Sigma \in \mathbb{R}_{\geq 0}^{d \times d}, \quad \gamma \in \mathbb{R}^d, \quad t \in [0, T],$$

where  $W = (W^1, \dots, W^d)$  is a vector of  $d$  independent Brownian motions. If the subordinator is a gamma or a generalized inverse Gaussian process we obtain a multidimensional variance gamma [86] or generalized hyperbolic [49] process.

Such processes can directly be simulated at fixed times  $t_0, \dots, t_M$ . To do so, generate the increments of the subordinator  $\Delta G_m = G_{t_m} - G_{t_{m-1}}$ ,  $m = 1, \dots, M$  with  $G_{t_0} = 0$ . These can be obtained easily as described in e.g. [114]. After that, draw independent random variables  $N_m^i \sim \mathcal{N}(0, 1)$ ,  $m = 1, \dots, M$ ,  $i = 1, \dots, d$  and set  $\Delta X_m = \Sigma N_m \sqrt{\Delta G_m} + \gamma \Delta G_m$ . The discretized path of  $X$  is then given by  $X(t_m) = \sum_{j=1}^m \Delta X_m$ . In particular, to obtain the realizations  $\tilde{X}_{T,j}$  of  $X$  at maturity one may set  $M = 1$  and  $t_M = T$ .

*Gaussian approximation* Let  $X$  be an  $\mathbb{R}^d$ -valued Lévy process with characteristic triplet  $(\gamma, 0, \nu)$ . For  $\varepsilon \in (0, 1]$  let  $\nu_\varepsilon \leq \nu$  be a measure such that  $\nu^\varepsilon := \nu - \nu_\varepsilon$  is a finite

measure and  $\int_{\mathbb{R}^d} |x|^2 \nu_\varepsilon(dx) < \infty$ . Decompose  $X$  correspondingly into its small and large jump parts as  $X = X_\varepsilon + X^\varepsilon$ . The process  $X^\varepsilon$  can be written as

$$X_t^\varepsilon = \gamma_\varepsilon t + N_t^\varepsilon, \tag{2.1}$$

where  $\gamma_\varepsilon := \gamma + \int_{|z|>1} z \nu_\varepsilon(dz) - \int_{|z|\leq 1} z \nu^\varepsilon(dz)$  and  $N^\varepsilon$  is a compound Poisson process. A first approximation of  $X_t$  is  $X_t \approx \gamma_\varepsilon t + N_t^\varepsilon$  where jumps of magnitude smaller than  $\varepsilon$  are neglected or replaced by their expected values in the finite activity case. This approximation is reasonable when the intensity of small jumps is low. If this is not the case, the small jump part  $X_\varepsilon$  can be approximated by an  $\mathbb{R}^d$ -valued standard Brownian motion  $W$  independent of  $N^\varepsilon$ . It is shown in [10, 109] that, under certain assumptions on the covariance matrix  $Q_\varepsilon := \int_{\mathbb{R}^d} z z^\top \nu_\varepsilon(dz)$ , the process  $Q_\varepsilon^{-1/2} X_\varepsilon$  converges in distribution to  $W$  as  $\varepsilon \rightarrow 0$ . Then, by [37], there holds

**Theorem 2.1** *Let  $X$  be an  $\mathbb{R}^d$ -valued Lévy process with characteristic triplet  $(\gamma, 0, \nu)$ . Assume that  $Q_\varepsilon$  is non-singular for every  $\varepsilon \in (0, 1]$  and that for every  $\delta > 0$  there holds*

$$\int_{\langle Q_\varepsilon^{-1}z, z \rangle > \delta} \langle Q_\varepsilon^{-1}z, z \rangle \nu_\varepsilon(dz) \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0.$$

*Assume further that for some family of non-singular matrices  $\{\Sigma_\varepsilon\}_{\varepsilon \in (0, 1]}$  there holds*

$$\Sigma_\varepsilon^{-1} Q_\varepsilon \Sigma_\varepsilon^{-\top} \rightarrow I_d, \quad \text{as } \varepsilon \rightarrow 0,$$

*where  $I_d$  denotes the identity matrix in  $\mathbb{R}^d$ . Then for all  $\varepsilon \in (0, 1]$  there exists a càdlàg process  $Y^\varepsilon$  such that (in the sense of finite dimensional distributions)*

$$X_t \stackrel{(d)}{=} \gamma_\varepsilon t + \Sigma_\varepsilon W_t + N_t^\varepsilon + Y_t^\varepsilon, \tag{2.2}$$

*and such that for all  $T > 0$ ,  $\sup_{t \in [0, T]} |\Sigma_\varepsilon^{-1} Y_t^\varepsilon| \xrightarrow{(\mathbb{P})} 0$ , as  $\varepsilon \rightarrow 0$ . Here,  $\gamma_\varepsilon, N^\varepsilon$  are given in (2.1), and  $W$  is an  $\mathbb{R}^d$ -valued standard Brownian motion independent of  $N^\varepsilon$ .*

Thus, a more refined approximation of  $X_t$  is given by  $X_t \approx \Sigma_\varepsilon W_t + \gamma_\varepsilon t + N_t^\varepsilon$ . The compound Poisson process  $N_t^\varepsilon$  can be simulated using a series representation.

For  $X$  being a tempered  $\alpha$ -stable process,  $\alpha \in (0, 2)$ , it is shown in [37] that the remainder process  $Y^\varepsilon$  in (2.2) satisfies  $\varepsilon^{\alpha/2-1} \sup_{t \in [0, T]} |Y_t^\varepsilon| \rightarrow 0$  as  $\varepsilon \rightarrow 0$ . Hence, the convergence rate of the Monte Carlo method may be rather low as  $\alpha$  is close to 2.

**Series representations** In this subsection, we briefly discuss the representation of an  $\mathbb{R}^d$ -valued pure jump Lévy process with characteristic triplet  $(0, 0, \nu)$  by an (infinite) series of random variables. Truncating such a series gives the possibility to simulate the process. In particular, truncated series become useful in connection with Lévy copulas. We closely follow [108] and set  $T = 1$  for simplicity.

In order to illustrate the basic idea of series approximation we begin by constructing an infinite series representation for the Poisson process. This representation can then be generalized to more complex processes. Let  $\Gamma_1, \Gamma_2, \dots$  be the jumping times

of a Poisson process with unit rate and  $V_1, V_2, \dots$  i.i.d. uniform random variables independent of the  $\Gamma_m$ . Then, a Poisson process with intensity  $\lambda$  can be written as

$$X_t = \sum_{m=1}^{\infty} 1_{\{\Gamma_m \leq \lambda\}} 1_{\{V_m < t\}} = \sum_{m=1}^{\infty} U^{(-1)}(\Gamma_m) 1_{\{V_m < t\}},$$

since the tail integral  $U(\cdot)$  of a Poisson process is given by  $U(z) = \lambda 1_{\{z \leq 1\}}$ . Here,  $V_m$  can be interpreted as the jump times and  $U^{(-1)}(\Gamma_m)$  as the jump sizes of  $X$ . To get an implementable algorithm, the series expansion has of course to be truncated. The two most obvious possibilities are either to fix a number  $M$  and only consider  $M$  jumps or to fix a tolerance  $\varepsilon$  and only consider  $M_\varepsilon = \inf\{m : U^{-1}(\Gamma_m) > \varepsilon\}$  jumps. In the latter case jumps with jump size smaller than  $\varepsilon$  are omitted. Both approaches yield a compound Poisson approximation of the Lévy process  $X$ .

In general, split  $X$  into  $X = X_\varepsilon + X^\varepsilon$  as above and consider the Lévy–Itô decomposition

$$X_t^\varepsilon = \int_0^t \int_{|z|>1} z J(ds, dz) + \int_0^t \int_{\varepsilon \leq |z| \leq 1} z (J(ds, dz) - \nu(dz) ds).$$

The Poisson random measure  $J$  can be represented in the form  $J = \sum_{m=1}^{\infty} \delta_{(V_m, G_m)}$  as in [70, Proposition II.1.14], where  $\{G_i\}$  is a sequence of random variables independent of the i.i.d. sequence  $\{V_i\} \sim U(0, 1)$ . Using this representation of  $J$ , the process  $X^\varepsilon$  can be rewritten as

$$X_t^\varepsilon = \sum_{m \in \Lambda_\varepsilon(\omega)} G_m 1_{\{V_m \leq t\}} - t\gamma_\varepsilon, \quad t \in [0, T],$$

where  $\Lambda_\varepsilon(\omega) := \{m \geq 1 : |G_m(\omega)| \geq \varepsilon\}$  and  $\gamma_\varepsilon := \int_{\varepsilon \leq |z| \leq 1} z \nu(dz)$ . As  $\varepsilon \rightarrow 0$ ,  $X^\varepsilon \xrightarrow{\text{a.s.}} X$ . One obtains a series representation of  $X$  as

$$X_t = \sum_{m=1}^{\infty} G_m 1_{\{V_m \leq t\}} - t\gamma_m, \tag{2.3}$$

with a suitable sequence of centers  $\{\gamma_m\}$ . Note that the sequences  $\{G_m\}$  and  $\{\gamma_m\}$  in (2.3) are not unique. There are several methods to represent them, e.g. LePage’s method, Bondesson’s method, the rejection method and the shot noise method. For details we refer to [108]. Here, for the sake of brevity, we only illustrate LePage’s series representation (see [38, 108]) and focus on the representation of  $G_m$ .

LePage’s method is based on the radial decomposition of the Lévy measure as

$$\nu(A) = \int_{\mathbb{S}^{d-1}} \int_0^\infty 1_A(zs) \mu(dz, s) \lambda(ds), \quad A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\}), \tag{2.4}$$

where  $\lambda$  is a probability measure on the unit sphere  $\mathbb{S}^{d-1}$  and  $\mu(\cdot, s)$  is a Lévy measure on  $(0, \infty)$  for each  $s \in \mathbb{S}^{d-1}$ . Define the generalized inverse tail integral

$$U^{(-1)}(x, s) := \inf\{z > 0 : U(z, s) < x\}, \quad \text{where } U(z, s) := \int_z^\infty \mu(dx, s).$$

With this notation, the  $G_m$  in (2.3) are given by  $G_m = U^{(-1)}(\Gamma_m, Y_m)Y_m$ , where  $\{\Gamma_m\}$  is a sequence of jumping times of some standard Poisson process and  $\{Y_m\}$  is an independent sequence of i.i.d. random vectors taking values in  $\mathbb{S}^{d-1}$  with distribution  $\lambda$  given by (2.4). There are two difficulties arising in the practical simulation of  $G_m$ . Firstly, it may be hard to draw the random vectors  $Y_m$  in  $\mathbb{S}^{d-1}$  with distribution  $\lambda$ . Secondly, one needs a closed form expression (or a reasonable method to compute it numerically) for the inverse tail integral  $U^{(-1)}$  of the measure  $\mu(\cdot, s)$ , for each  $s$ .

It is also possible to use the series representation (2.3) to simulate Lévy processes given by Lévy copulas [38, 117]. Here, the jump sizes  $G_m^j$  of each component  $j = 1, \dots, d$  are calculated conditionally on the components  $i = 1, \dots, j - 1$ . For the sake of simplicity, we discuss the method for  $\mathbb{R}^2$ -valued Lévy processes having positive jumps, following [38]. For a more general treatment, we refer to [117]. Let  $F$  be the Lévy copula associated to  $X$  as given in (1.7) with marginal tail integrals  $U_1, U_2$ . Assume  $F$  is continuous on  $[0, \infty]^2$ . We again let  $\Gamma_1^1, \Gamma_2^1, \dots$  be the jumping times of a Poisson process with unit rate and  $V_1, V_2, \dots$  i.i.d. uniform random variables independent of  $\Gamma_m^1, m \in \mathbb{N}$ . Now, additionally consider  $\Gamma_1^2, \Gamma_2^2, \dots$ , which are independent of all other variables and distributed according to  $\partial_u F(u, v) |_{u=\Gamma_m^1}$ . Then, in law there holds

$$X_t^j = \sum_{m=1}^{\infty} U_j^{(-1)}(\Gamma_m^j) 1_{\{V_m \leq t\}}.$$

To simulate  $X^1, X^2$  we start as explained above. We fix a tolerance  $\varepsilon$  and only consider  $M_\varepsilon = \inf\{m : U_1^{-1}(\Gamma_m) > \varepsilon\}$  jumps. Then, for the  $m$ -th jump,  $m \in \{1, \dots, M_\varepsilon\}$ , compute the jump time of the Poisson process,  $\Gamma_m^1 = \sum_{j=1}^m T_j$ , where  $T_j$  are standard exponentially distributed random variables. Then, compute  $\Gamma_m^2 \sim \partial_u F(u, v) |_{u=\Gamma_m^1}$  and the jump time  $V_m \sim U(0, 1)$ . The discretized path is given by

$$X_t^j = \sum_{m=1}^{M_\varepsilon} U_j^{(-1)}(\Gamma_m^j) 1_{\{V_m \leq t\}}.$$

*Remark 2.2* Suppose the underlying stochastic process  $S$  is not modeled explicitly as in (1.1), but as the solution of a stochastic differential equation (SDE) of the form

$$S_t = S_0 + \int_0^t f(S_{s-}) dX_s, \quad t \in [0, T], \tag{2.5}$$

where  $f$  is some suitable given function and  $X$  is an  $\mathbb{R}^d$ -valued Lévy process. The process  $S$  might for instance arise from a local or stochastic volatility model. In general, it is no longer a Lévy process. To solve the pricing equation (1.2) in such models, one may use the well-known Euler–Maruyama scheme [89] to approximate the solution of (2.5). This allows to use Monte Carlo methods also in this setting. To implement the Euler–Maruyama scheme, one has to simulate increments of the driving Lévy process. As discussed above, in general this can be done only approximately, so that the approximation of the solution of the SDE bears two errors, one stemming

from the discretization through the Euler–Maruyama scheme, and one coming from the approximation of the underlying Lévy process, see e.g. [68, 69, 110] and the references therein.

In the Black–Scholes setting this approach has been widely studied; for an introductory overview we refer to [73].

*Remark 2.3* Quasi-Monte Carlo as well as randomized Quasi-Monte Carlo methods have gained a lot of interest over the last decade; see e.g. [58, 79] and the references therein. A recent survey by L’Ecuyer can be found in [76]. These methods rely on a careful choice of random numbers that drive the simulation and can improve the  $O(N^{-1/2})$  convergence rate of ordinary Monte Carlo methods. Recently, they have been applied in the context of Lévy processes in [11, 80].

*Variance reduction* In order to obtain more precise estimates, an alternative to simply increasing the number of simulations  $N$  is to use variance reduction techniques. For a description of the most relevant variance reduction techniques like control variates technique, antithetic sampling, importance sampling, stratified and bridge sampling, moment matching and low-discrepancy sequences we refer to the monographs [9, 58] as well as [120].

Here, we only illustrate the basic idea of control variates. Let  $\bar{V} = N^{-1} \sum_{m=1}^N X_m$  be a Monte Carlo estimator of the expectation  $V = \mathbb{E}[X]$  of a random variable  $X$ . The basic idea of the control variates technique is to look for a random variable  $Y$  which is highly correlated with  $X$  and has *known mean*  $\mathbb{E}[Y]$ . Then one may use the empirical means  $\bar{V}, \bar{Y}$  to obtain an estimator with lower variance. To do so, for  $\alpha \in \mathbb{R}$  consider the random variable

$$X_\alpha := X + \alpha(Y - \mathbb{E}[Y]).$$

Its variance is  $\text{Var}[X_\alpha] = \text{Var}[X] + \alpha^2 \text{Var}[Y] + 2\alpha \text{Cov}(X, Y)$ , which is minimized by the value  $\alpha^* := -\frac{\text{Cov}(X, Y)}{\text{Var}[Y]}$ . In this case there holds  $\text{Var}[X_{\alpha^*}] = (1 - \rho^2) \text{Var}[X]$ , where  $\rho$  denotes the correlation between  $X$  and  $Y$ . Thus, the variance of  $X$  is reduced by a factor of  $1 - \rho^2$  and the variance reduction performs well if  $\rho$  is close to  $\pm 1$ . The *unbiased* control variate estimator  $\bar{V}_\alpha$  for  $V = \mathbb{E}[X]$  is hence defined by

$$\bar{V}_\alpha := \bar{V} + \alpha(\bar{Y} - \mathbb{E}[Y]).$$

In practice,  $\alpha^*$  is usually not known and is therefore replaced by its sample counterpart

$$\bar{\alpha}^* := -\frac{\sum_{m=1}^N (X_m - \bar{X})(Y_m - \bar{Y})}{\sum_{m=1}^N (Y_m - \bar{Y})^2}.$$

In the context of option pricing, the control variate  $Y$  may, for example, be chosen to be lower and/or upper bounds for the option price itself (see e.g. [38]) or the price of the underlying asset at maturity  $S_T$  (see e.g. [58]). A combination of control variates and importance sampling for Lévy processes has been applied in [72].



*American options* Monte Carlo evaluation of American options has a “Monte Carlo on Monte Carlo” feature, since the determination of the optimal exercise time depends on an average over future events. To see this, consider a point  $(X_t, t)$  on a single simulated path  $X$ . In order to decide whether to exercise at this point, one has to evaluate the expectation (1.3). This requires continuation from  $(X_t, t)$  on many branching paths, and makes this direct approach infeasible.

In the Black–Scholes setting several methods have been suggested to overcome this difficulty. For a comparison of the main approaches we refer to [55]. The following have been introduced. The path-bundling technique provides lower and upper bounds for the true price of an American contract; see e.g. [26, 27]. The martingale optimization approach replaces the maximization over stopping times by minimizing over martingales and provides also an upper bound, see [35, 62, 107]. In [84], the future expectation is replaced by a least square interpolation. This method has been applied to value an American option with  $S$  in (1.1) following the Merton model ( $d = 1$ ). Furthermore, in [107] some Monte Carlo results for the valuation of American options under spectrally one-sided Lévy processes are presented.

### 3 Fast Fourier methods

Contrary to the classical Black–Scholes case in Lévy models there are usually no closed form option prices since the probability density of a Lévy process is typically not known in closed form. However, the characteristic function of this density can be expressed in terms of elementary functions for the majority of one-dimensional Lévy processes discussed in the literature. This has led to the development of Fourier-based option pricing methods where the Fourier transform and its inverse,

$$\mathcal{F}[g(x)](z) = (2\pi)^{-d} \int_{\mathbb{R}^d} e^{i\langle z, x \rangle} g(x) dx, \quad z \in \mathbb{R}^d,$$

$$\mathcal{F}^{-1}[g(z)](x) = \int_{\mathbb{R}^d} e^{-i\langle x, z \rangle} g(z) dz, \quad x \in \mathbb{R}^d,$$

are efficiently evaluated numerically by using the FFT algorithm, see [19, 20, 34, 47, 67, 78, 83, 85, 95, 100]. We present two approaches here. First, following the line of [20, 34, 47, 78] we transform the option value with respect to the log strike price. Secondly, it is also possible to transform the option value with respect to the log spot price as in [67, 83, 100]. For simplicity we set the interest rate  $r = 0$ .

*Transformation with respect to the log strike price* Consider an option with strike  $K = e^k$ , payoff  $g(k) = (e^{a \cdot X_T} - e^k)^+$  with  $a \in [0, 1]^d$  and maturity  $T$ . For the sake of notational simplicity suppose  $t = 0$ . The option price is given by

$$V(0, k) = \int_{\mathbb{R}^d} (e^{a \cdot s} - e^k)^+ p_T(s) ds, \quad k \in \mathbb{R},$$

where  $p_T(s)$  denotes the corresponding joint density of  $X_1, \dots, X_d$  at  $T$ .

Since the payoff  $g(k)$  tends to a positive constant as  $k \rightarrow -\infty$ , the Fourier transformation of  $g(k)$  does not exist in general. Therefore, instead of  $g(k)$  one has to consider the damped payoff  $e^{\alpha k} g(k)$  with a damping constant  $\alpha > 0$ . The Fourier transformation of the damped option price can then be written as

$$\begin{aligned} \mathcal{F}[e^{\alpha k} V(0, k)](z) &= (2\pi)^{-1} \int_{\mathbb{R}} e^{(\alpha+iz)k} V(0, k) dk \\ &= (2\pi)^{-1} \int_{\mathbb{R}^d} p_T(s) \int_{-\infty}^{(a,s)} e^{(\alpha+iz)k} (e^{(a,s)} - e^k) dk ds \\ &= (2\pi)^{-1} \int_{\mathbb{R}^d} p_T(s) \frac{e^{(1+\alpha+iz)(a,s)}}{(\alpha + iz)(1 + \alpha + iz)} ds \\ &= (2\pi)^{-1} \frac{\Phi_T((1 + \alpha + iz)a)}{(\alpha + iz)(1 + \alpha + iz)} \end{aligned}$$

and therefore

$$V(0, k) = \frac{e^{-\alpha k}}{2\pi} \mathcal{F}^{-1} \left[ \frac{\Phi_{T-t}((1 + \alpha + iz)a)}{(\alpha + iz)(1 + \alpha + iz)} \right], \quad k \in \mathbb{R}.$$

Note that, by Lemma 1.1, for  $\beta_i^- > (1 + \alpha)d, i = 1, \dots, d$  the characteristic function  $\Phi_T((1 + \alpha + iz)a)$  exists.

*Transformation with respect to the log spot price* We now consider a general European option with maturity  $T$  and payoff  $g(x)$  in log spot price  $x$ . The option price is then given by

$$V(t, x) = \mathbb{E}[g(X_T)|X_t = x] = \int_{\mathbb{R}^d} g(x + s) p_{T-t}(s) ds, \quad x \in \mathbb{R}^d, t \geq 0.$$

Now, if  $\mathcal{F}[g]$  exists, the Fourier transform of the option price can be written as

$$\begin{aligned} \mathcal{F}[V(t, x)](z) &= (2\pi)^{-d} \int_{\mathbb{R}^d} e^{i\langle z, x \rangle} V(t, x) dx \\ &= (2\pi)^{-d} \int_{\mathbb{R}^d} e^{i\langle z, y \rangle} g(y) dy \int_{\mathbb{R}^d} e^{-i\langle z, s \rangle} p_{T-t}(s) ds \\ &= \mathcal{F}[g(y)](z) \cdot \Phi_{T-t}(-z), \end{aligned}$$

where  $y = x + s$ . Therefore

$$V(t, x) = \mathcal{F}^{-1}[\mathcal{F}[g(y)](z) \cdot \Phi_{T-t}(-z)], \quad x \in \mathbb{R}^d, t \geq 0. \tag{3.1}$$

The restriction for the Fourier transformation of the payoff to exist is quite strong, since it is not even satisfied for a simple basket option. To weaken this assumption one may again try to dampen the payoff. But this approach is only practicable in dimension  $d = 1$ , since for most multivariate payoff functions there exist some coordinate directions  $j \in \{1, \dots, d\}$  such that  $\lim_{y_j \rightarrow \pm\infty} g(y) > 0$ , e.g. for basket options. Therefore, in the  $d$ -dimensional case one usually has to localize the payoff to

a bounded domain. One strategy of doing so with an explicit error analysis is described in Sect. 4 below. Thereafter, in most cases the Fourier transformation of  $g(y)$  must be evaluated numerically and one hence has to calculate both  $\mathcal{F}$  and  $\mathcal{F}^{-1}$  in (3.1) numerically. The computational cost is doubled. In dimension  $d = 1$  however the Fourier transformation of most payoffs can be obtained analytically and only one Fourier transformation, i.e.,  $\mathcal{F}^{-1}$  in (3.1), has to be evaluated numerically.

*Discretization* The multidimensional discrete Fourier transform of a given series of data points  $f_{\underline{j}}$  is given by the collection

$$\hat{f}_{\underline{k}} = \sum_{j_1=0}^{N-1} \dots \sum_{j_d=0}^{N-1} e^{2\pi i \langle \underline{k}, \underline{j} \rangle / N} f_{\underline{j}}, \quad k_n = 0, \dots, N - 1, \quad n = 1, \dots, d.$$

To compute  $\hat{f}_{\underline{k}}$ ,  $k_n = 0, \dots, N - 1, n = 1, \dots, d$  one a priori needs  $\mathcal{O}(N^{2d})$  operations. Utilizing the so-called *fast Fourier transform* [41, 98] this computational cost can be reduced to  $\mathcal{O}(N^d \log N)$ . For instance, suppose we want to approximate the inverse Fourier transform of a function  $f(z)$  with a discrete Fourier transform (to solve (3.1) one may choose  $f(z) = \mathcal{F}[g(y)](z) \cdot \Phi_{T-t}(-z)$ ). Then, the integral can be truncated and discretized using the trapezoidal rule, to give

$$\begin{aligned} \mathcal{F}^{-1}[f(z)](x) &= \int_{\mathbb{R}^d} e^{-i \langle x, z \rangle} f(z) \, dz \approx \int_{[-R, R]^d} e^{-i \langle x, z \rangle} f(z) \, dz \\ &\approx \sum_{j_1=0}^{N-1} \dots \sum_{j_d=0}^{N-1} \omega_{\underline{j}} f(z_{\underline{j}}) e^{-i \langle x, z_{\underline{j}} \rangle}, \end{aligned}$$

with discretization step  $\Delta z = \frac{2R}{N-1}$ ,  $z_{j_n}^n = -R + j_n \Delta z$  in Fourier space and suitable weights  $w_j$ ; see e.g. [67].

Herewith, in order to obtain an approximate value of  $V(t, x)$  in (3.1) for any  $x \in \mathbb{R}^d$ , we also have to discretize the spot price or  $x$ -domain  $\mathbb{R}^d$ . For this, we define an additional grid by setting  $x_{j_n}^n = -R_2 + k_n \Delta x$  with step size  $\Delta x = \frac{2R_2}{N-1}$  and given  $R_2 > 0$ . With the relation

$$\Delta z \cdot \Delta x = \frac{2\pi}{N} \tag{3.2}$$

we then find

$$\begin{aligned} \mathcal{F}^{-1}[f(z)](x_{\underline{k}}) &\approx e^{iR \langle x_{\underline{k}}, \mathbf{1} \rangle} \sum_{j_1=0}^{N-1} \dots \sum_{j_d=0}^{N-1} e^{-i2\pi \langle \underline{k}, \underline{j} \rangle / N} \omega_{\underline{j}} f(z_{\underline{j}}) e^{iR_2 \Delta z \langle \underline{1}, \underline{j} \rangle} \\ &= e^{iR \langle x_{\underline{k}}, \mathbf{1} \rangle} \hat{f}_{\underline{k}}. \end{aligned}$$

This expression can now be evaluated very efficiently using the fast Fourier transform as mentioned above. Also note that by (3.2) the discretization of the Fourier space and the spot price (or strike price) space are related and cannot be chosen independently. No time stepping is required and for  $d = 1$  dimension only  $\mathcal{O}(N \log N)$

work is needed to obtain the price at  $N$  spot (or strike) prices. For convergence rates and error analysis see [78].

*American options* In order to compute American option prices, one can approximate them by a sequence of Bermudan options with increasing number of exercise dates (as introduced in [56]).

For this, let  $t = t^0 < \dots < t^M = T$  be a time discretization which can be thought of as exercise times of a Bermudan option. Its price can be computed by backward induction as

$$\begin{aligned}
 V(t^M, x) &= g(x), \\
 V(t^m, x) &= \max(\mathbb{E}[V(t^{m+1}, X_{t^{m+1}}) | X_{t^m} = x], g(x)), \quad m = M - 1, \dots, 0,
 \end{aligned}
 \tag{3.3}$$

where at each time point equation (3.1) is solved with time step  $t^{m+1} - t^m$  as in [67, 85]. The overall computational cost for this approximation then is  $\mathcal{O}(MN^d \log N)$ . In [19] the Wiener–Hopf factorization is used to compute the values of perpetual American options.

### 4 PIDE-based methods

In the Black–Scholes setting, when the underlying process  $S$  is a geometric Brownian motion, it is well known that the solution  $V(t, S)$  of the pricing equation (1.2) can be described as the solution of a parabolic partial differential equation (PDE) also known as the Black–Scholes equation, namely

$$\frac{\partial V}{\partial t}(t, S) + \frac{1}{2} \sum_{i,j=1}^d S_i S_j Q_{ij} \frac{\partial^2 V}{\partial S_i \partial S_j} + r \sum_{i=1}^d S_i \frac{\partial V}{\partial S_i}(t, S) - rV(t, S) = 0, \tag{4.1}$$

with suitable boundary conditions depending on the payoff function  $g(\cdot)$ . Since PDEs of the form (4.1) are well studied objects in engineering and numerical mathematics, there exists a whole zoo of sophisticated and very general so-called finite difference and finite element methods for their numerical solution (at least in dimension  $d \leq 3$ ). For an introduction see e.g. [23] or for a more finance-related perspective [2, 48, 115]. Over the last years many authors have developed several variants of these methods, especially for the efficient treatment of linear and non-linear problems arising in diffusion-driven markets, see e.g. [1, 2, 51, 52, 97, 111, 118] and the references therein.

In this survey article we focus on markets driven by general Lévy processes. Therefore, we illustrate in this section how numerical methods to solve (4.1) can be modified and extended in order to yield a general and efficient pricing technique which can be applied to general Lévy models also in moderate dimension  $d > 3$ . The main difference is that corresponding to the existence of jumps in a Lévy model an additional integral term has to be introduced to (4.1) resulting in a partial integro-differential equation (PIDE). More precisely, if  $S$  is driven by an exponential Lévy process  $X$  as in (1.1) then, by [104, Theorem 4.2], the solution  $V(t, S)$  of (1.2) can be characterized by

**Theorem 4.1** *Let  $X$  be a Lévy process with state space  $\mathbb{R}^d$  and characteristic triplet  $(\gamma, Q, \nu)$ . Assume that the function  $V(t, S)$  in (1.2) satisfies*

$$V(t, S) \in C^{1,2}((0, T) \times \mathbb{R}_{>0}^d) \cap C^0([0, T] \times \mathbb{R}_{\geq 0}^d).$$

Then  $V(t, S)$  satisfies the PIDE

$$\begin{aligned} \frac{\partial V}{\partial t}(t, S) + \frac{1}{2} \sum_{i,j=1}^d S_i S_j Q_{ij} \frac{\partial^2 V}{\partial S_i \partial S_j} + r \sum_{i=1}^d S_i \frac{\partial V}{\partial S_i}(t, S) - rV(t, S) \\ + \int_{\mathbb{R}^d} \left( V(t, Se^z) - V(t, S) - \sum_{i=1}^d S_i (e^{z_i} - 1) \frac{\partial V}{\partial S_i}(t, S) \right) \nu(dz) = 0 \end{aligned} \tag{4.2}$$

in  $(0, T) \times \mathbb{R}_{\geq 0}^d$  where  $V(t, Se^z) := V(t, S_1 e^{z_1}, \dots, S_d e^{z_d})$ , and the terminal condition is given by

$$V(T, S) = g(S) \quad \forall S \in \mathbb{R}_{\geq 0}^d.$$

For its numerical solution, by [104, Corollary 4.3], the PIDE (4.2) can be transformed into a simpler form.

**Corollary 4.2** *Let  $X$  be a Lévy process with state space  $\mathbb{R}^d$  and characteristic triplet  $(\gamma, Q, \nu)$  and marginal Lévy measures  $\nu_i, i = 1, \dots, d$  satisfying (1.4) with  $\beta_i^+ > 1, \beta_i^- > 0, i = 1, \dots, d$ . Furthermore, let*

$$u(\tau, x) = e^{r\tau} V(T - \tau, e^{x_1 + (\gamma_1 - r)\tau}, \dots, e^{x_d + (\gamma_d - r)\tau}), \tag{4.3}$$

where

$$\gamma_i = \frac{Q_{ii}}{2} + \int_{\mathbb{R}} (e^{z_i} - 1 - z_i) \nu_i(dz_i).$$

Then  $u$  satisfies the PIDE

$$\frac{\partial u}{\partial \tau} + \mathcal{A}_{BS}[u] + \mathcal{A}_J[u] = 0 \tag{4.4}$$

in  $(0, T) \times \mathbb{R}^d$  with initial condition  $u(0, x) := u_0$ . The differential operator is defined for  $\varphi \in C_0^2(\mathbb{R}^d)$  by

$$\mathcal{A}_{BS}[\varphi] = -\frac{1}{2} \sum_{i,j=1}^d Q_{ij} \frac{\partial^2 \varphi}{\partial x_i \partial x_j}, \tag{4.5}$$

and the integro-differential operator by

$$\mathcal{A}_J[\varphi] = - \int_{\mathbb{R}^d} \left( \varphi(x + z) - \varphi(x) - \sum_{i=1}^d z_i \frac{\partial \varphi}{\partial x_i}(x) \right) \nu(dz). \tag{4.6}$$

The initial condition is given by

$$u_0 = g(e^x) := g(e^{x_1}, \dots, e^{x_d}). \tag{4.7}$$

Furthermore, for American options instead of a PIDE-based characterization one obtains an *inequality* representation provided the price  $u(\cdot, \cdot)$  solves a partial integro-differential inequality, i.e.,

$$\begin{aligned} \frac{\partial u}{\partial \tau} + (\mathcal{A}_{BS} + \mathcal{A}_J)[u] &\leq 0, \\ u(\tau, \cdot) &\geq \tilde{g}_\tau, \\ (u - \tilde{g}_\tau) \left( \frac{\partial u}{\partial \tau} + (\mathcal{A}_{BS} + \mathcal{A}_J)[u] \right) &= 0, \end{aligned} \tag{4.8}$$

with  $\mathcal{A}_{BS}$  and  $\mathcal{A}_J$  as in (4.4), where  $\tilde{g}_\tau$  denotes the payoff function  $g$  transformed according to (4.3), i.e.,

$$\tilde{g}_\tau(x) = e^{r\tau} g(e^{x_1 + (\gamma_1 - r)\tau}, \dots, e^{x_d + (\gamma_d - r)\tau}), \quad x \in \mathbb{R}^d. \tag{4.9}$$

For the derivation of (4.8) see e.g. [90] and [38, Sect. 12.1.3] based on the methodology of [14, 15].

The implementation of any finite difference or finite element scheme for (4.4), (4.8) requires the *localization* of the log price domain  $\mathbb{R}^d$  to a bounded domain  $B_R := [-R, R]^d$ ,  $R > 0$ . For this, we find that in finance truncation of  $\mathbb{R}^d$  to  $B_R$  corresponds to approximating the solution  $u$  of (4.4) by the price  $u_R$  of a corresponding barrier option on  $B_R$ . In log-prices  $u_R$  is given by

$$u_R(t, x) = \mathbb{E}[g(e^{X_T}) 1_{\{T < \tau_{B_R, t}\}} | X_t = x],$$

where  $\tau_{B_R, t} = \inf\{s \geq t | X_s \notin B_R\}$  denotes the first exit time of  $X$  from  $B_R$  after time  $t$ . In case the underlying stochastic process  $X$  admits semiheavy tails (1.4), the solution  $u_R$  of the localized problem converges pointwise exponentially to the solution  $u$  of (4.4), i.e., there exist constants  $c_1, c_2 > 0$  such that

$$|u(t, x) - u_R(t, x)| \lesssim e^{-c_1 R + c_2 \|x\|_\infty}.$$

It therefore indeed suffices to replace the original price space domain  $\mathbb{R}^d$  by  $B_R$  with sufficiently large  $R > 0$ . For details we refer to [104]. Furthermore, note that for a barrier option on  $B_R$  there is no localization error.

### 4.1 Finite difference methods

After localization of the original space domain  $\mathbb{R}^d$  to  $B_R$  as described above, the numerical solution of (4.4) by finite differences is obtained in three main steps:

1. The integration domain  $\mathbb{R}^d$  in (4.6) must also be localized to a bounded domain.
2. The small jumps must be approximated by a Brownian motion.
3. The solution is computed at discrete grid points and the derivatives in (4.4)–(4.6) are replaced by finite differences.

*Localization of the integration domain* The integration domain  $\mathbb{R}^d$  in (4.6) is truncated to a bounded domain  $\Lambda_Z = [-Z, Z]^d$ ,  $Z > 0$ . Similarly to the localization of the spatial domain it can be shown that the error decays exponentially with respect to  $Z$ ; see [39, 104].

*Approximation of small jumps* In order to numerically integrate the jump measure in the finite difference discretization described below, the small jumps of the Lévy process need to be truncated. By doing this, the Lévy measure becomes finite. More precisely, we introduce a truncation parameter  $\varepsilon \in (0, 1]$  and split the process  $X$  into  $X = X_\varepsilon + X^\varepsilon$  as defined in Sect. 2. The process  $X_\varepsilon$  is approximated by a Brownian motion and for the remainder process  $X^\varepsilon$  we then obtain the triplet  $(\gamma + \gamma_\varepsilon, Q + Q_\varepsilon, \nu^\varepsilon)$ , where the Lévy measure  $\nu^\varepsilon$  is now a finite measure, i.e.,  $\nu^\varepsilon(\mathbb{R}^d) < \infty$ .

Note that the truncation of the small jumps based on the non-financial parameter  $\varepsilon > 0$  introduces an additional error to the discretization which can have a significant impact on the accuracy and stability of the numerical scheme. For examples, such as barrier contracts under pure jump Lévy models, we refer to [82, Sects. 6.2, 8.3].

*Discretization* Consider a uniform grid on  $[0, T] \times [-R, R]^d$  with time step  $\Delta t = \frac{T}{M}$  and mesh width  $\Delta x = \frac{2R}{N}$  for  $N, M \in \mathbb{N}$ . Then, the time and space points are given by  $t_m = m\Delta t$  and  $x_{n_j}^j = -R + n_j\Delta x$  where  $m = 0, \dots, M$ ,  $n_j = 0, \dots, N$  and  $j = 1, \dots, d$ . Let  $\underline{u}_n^m = u(t_m, x_{n_1}^1, \dots, x_{n_d}^d)$  be the solution on the grid which is zero outside of  $[-R, R]^d$ . The spatial derivatives in (4.5) can be approximated using finite differences by

$$\frac{\partial^2 u}{\partial x^j \partial x^j}(x^j) \approx \frac{\underline{u}_{n_j+1} - 2\underline{u}_{n_j} + \underline{u}_{n_j-1}}{(\Delta x)^2},$$

$$\frac{\partial^2 u}{\partial x^j \partial x^i}(x^j, x^i) \approx \frac{\underline{u}_{(n_j+1, n_i+1)} - \underline{u}_{(n_j+1, n_i-1)} - \underline{u}_{(n_j-1, n_i+1)} + \underline{u}_{(n_j-1, n_i-1)}}{4(\Delta x)^2},$$

and the integral in (4.6) is numerically integrated using a trapezoidal quadrature rule with the same grid resolution  $\Delta x$ . Using this, the jump operator (4.6) reads

$$- \int_{\Lambda_Z} (u(x_n + z) - u(x_n)) \nu^\varepsilon(dz) \approx - \sum_{\ell_1} \dots \sum_{\ell_d} (\underline{u}_{n+\underline{\ell}} - \underline{u}_n) \underline{\nu}_\underline{\ell},$$

where  $\underline{\nu}_\underline{\ell} = \int_{(\ell_1-1/2)\Delta x}^{(\ell_1+1/2)\Delta x} \dots \int_{(\ell_d-1/2)\Delta x}^{(\ell_d+1/2)\Delta x} \nu^\varepsilon(dz)$ . Note that  $\underline{u}_{n+\underline{\ell}} = 0$  for  $n_j + \ell_j \notin \{0, \dots, N\}$ . Non-zero boundary conditions are treated in [39]. Denoting by  $\mathbf{A}_{BS}$  and  $\mathbf{A}_J$  the discretization matrices representing the differential and jump part, we can write a  $\theta$ -time stepping scheme as

$$\frac{\underline{u}^{m+1} - \underline{u}^m}{\Delta t} + \theta_1 \mathbf{A}_{BS} \underline{u}^{m+1} + (1 - \theta_1) \mathbf{A}_{BS} \underline{u}^m + \theta_2 \mathbf{A}_J \underline{u}^{m+1} + (1 - \theta_2) \mathbf{A}_J \underline{u}^m = 0. \tag{4.10}$$

Note that the matrix  $\mathbf{A}_{BS}$  is sparse whereas the matrix  $\mathbf{A}_J$  is densely populated. For  $\theta_1 = \theta_2 = 0$  the scheme is explicit but not unconditionally stable. Therefore, to obtain stability small time steps are required. For  $\theta_1 = \theta_2 = 1$  the scheme is implicit and

there are no stability constrains. However, at each time step a linear system with a full matrix has to be solved. Therefore, Cont and Voltchkova [39] propose an explicit-implicit scheme with  $\theta_1 = 1, \theta_2 = 0$  in dimension  $d = 1$  and prove convergence of the fully discrete scheme. In particular, they show that under certain smoothness assumptions on the payoff function  $g(\cdot)$  one obtains first order convergence in  $\Delta x$ .

Similar techniques for  $d = 1$  and  $d = 2$  are also shown in [24]. In dimension  $d = 1$ , it is also possible to use the fast Fourier transform and exploit the Toeplitz structure of the matrix  $A_J$ , cf. [3, 7].

However, the described discretization suffers from the so-called ‘‘curse of dimension,’’ i.e., the number of grid points grows like  $\mathcal{O}(N^d)$ . Therefore, high-dimensional problems with  $d > 3$  cannot be solved. To break this curse, for the Black–Scholes equation Reisinger and Wittum [105] employed so-called sparse grids (see [30, 125]) based on the combination technique (see e.g. [31]). Here, the number of grid points only grows like  $\mathcal{O}(N(\log N)^{d-1})$ . To our knowledge, no finite difference schemes for Lévy models of dimension  $d > 2$  have been considered in the literature up to now.

*Remark 4.3* (Multinomial tree methods) Among the most popular and intuitive numerical pricing techniques in the Black–Scholes setting are the so-called multinomial tree or Markov chain methods. The tree method originally dates back to Cox et al. [42] and was subsequently extended to finite activity jump diffusion models [5, 93]. The pricing of European and American options with multinomial approximation has thereafter been considered by several authors, see e.g. [6, 63, 88]. For multidimensional models however the method fails due to a rapid, exponential growth of complexity with the dimension. An overview can be found in e.g. [75]. The basic idea reads as follows. For the exponential Lévy process  $S$  one can construct a discrete-time Markov chain  $s$  by setting

$$s^{n+1} = S_{t+\Delta t}^{\Delta t, \Delta x} = S_t^{\Delta t, \Delta x} \exp(\varepsilon^n) = s^n \exp(\varepsilon^n),$$

where  $\varepsilon^n$  denotes an i.i.d. family of random variables taking  $k \in \mathbb{N}$  values. Usually, the values of  $\varepsilon^n$  are chosen to be multiples of the given step size  $\Delta x$ ,

$$\varepsilon^n \in \{-k_1 \Delta x, \dots, -\Delta x, 0, \Delta x, \dots, k_2 \Delta x\}.$$

Then there holds  $k = k_1 + k_2 + 1$ . The numbers  $k_1, k_2 \in \mathbb{N}$  are allowed to differ to account for asymmetry of jumps. The paths of the Markov chain  $\ln s$  fall on a lattice with step size  $(\Delta t, \Delta x)$ . They can therefore be seen as an explicit finite difference scheme in  $(t, \ln S)$ -space. The values of  $k_1, k_2$  and the transition probabilities have to be chosen in such a way that absence of arbitrage is guaranteed, cf. [5]. For example, if we denote quantities on the lattice by  $A_j^n$ , where  $n$  denotes the time index of a node and  $j$  the price space index, the approximate value  $V$  of a European option can be computed by backward induction; starting at maturity from the final node  $N \Delta t = T$ , at each node in the tree the option value is given by the discounted expectation of the values on the branches, i.e.,

$$V_j^n = e^{-r \Delta t} \sum_{i=-k_1}^{k_2} q_i V_{j+i}^{n+1},$$



where  $q_i, i = -k_1, \dots, k_2$ , denote the transition probabilities  $q_i = \mathbb{P}[\varepsilon^n = i \Delta x]$ . For American options this backward step has to be replaced by taking the maximum of  $V_j^n$  and the corresponding payoff from exercising at the current time. See also [93]. If the transition probabilities are chosen such that  $S^{\Delta t, \Delta x}$  converges weakly to  $S$  as  $(\Delta t, \Delta x) \rightarrow 0$ , convergence of the discrete time European and American option prices to their continuous time counterparts can be shown to hold, cf. [75, 99].

### 4.2 Finite element methods

Instead of solving the PIDE (4.4) directly, the finite element method is based on the reformulation of (4.4) into the corresponding variational Galerkin equation. To this end, for  $u, v \in C_0^\infty(\mathbb{R}^d)$  we associate with  $\mathcal{A}_{BS}$  in (4.5) the bilinear form

$$\mathcal{E}_{BS}(u, v) = \frac{1}{2} \sum_{i,j=1}^d Q_{ij} \int_{\mathbb{R}^d} \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_j} dx.$$

To the jump part  $\mathcal{A}_J$  in (4.6) we associate the bilinear jump form

$$\mathcal{E}_J(u, v) = - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \left( u(x+z) - u(x) - \sum_{i=1}^d z_i \frac{\partial u}{\partial x_i}(x) \right) v(x) dx \nu(dz), \quad (4.11)$$

and set

$$\mathcal{E}(u, v) = \mathcal{E}_{BS}(u, v) + \mathcal{E}_J(u, v).$$

Denoting by  $\mathcal{D}(\mathcal{E})$  the domain of  $\mathcal{E}(\cdot, \cdot)$ , the variational problem associated to (4.4) reads

Find  $u \in L^2((0, T); \mathcal{D}(\mathcal{E})) \cap H^1((0, T); \mathcal{D}(\mathcal{E})^*)$  such that

$$\left\langle \frac{\partial u}{\partial \tau}, v \right\rangle_{\mathcal{D}(\mathcal{E})^*, \mathcal{D}(\mathcal{E})} + \mathcal{E}(u, v) = 0, \quad \text{a.e. } \tau \in (0, T), \quad \forall v \in \mathcal{D}(\mathcal{E}), \quad (4.12)$$

$$u(0) = u_0,$$

where  $u_0$  is defined as in (4.7). For the well-posedness of (4.12) we refer to [91] for one-dimensional and to [50, 104] for certain multidimensional Lévy models. For instance, if  $Q > 0$  in (1.5) the domain  $\mathcal{D}(\mathcal{E})$  coincides with the Sobolev space  $H^1(\mathbb{R}^d)$ , and for  $Q = 0$  and tempered stable margins  $\mathcal{D}(\mathcal{E})$  can be seen to be some anisotropic Sobolev space.

*Remark 4.4* One key advantage of the variational formulation (4.12) is that the bilinear form  $\mathcal{E}(\cdot, \cdot)$  allows for a *naturally singularity free* discretization of general Lévy measures, i.e., the small jumps of the process do not need to be approximated by a Brownian motion. In contrast to Sect. 4.1, no additional truncation error is introduced. Consider, for example, a symmetric Lévy measure  $\nu$ . Then, by [50, Proposition 4.1],

the bilinear form  $\mathcal{E}_J(\cdot, \cdot)$  in (4.11) can be rewritten as

$$\mathcal{E}_J(u, v) = - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (u(x+z) - u(x))(v(x+z) - v(x))\nu(dz) dx. \tag{4.13}$$

Using (1.6), one readily infers that merely Lipschitz-continuous functions  $u, v$  are sufficient for the integrals in (4.13) to exist in the Lebesgue sense. In addition, in [104, Proposition 4.11] it is shown that the original bilinear form (4.11) exists for all standard, continuous finite element basis functions and general non-symmetric Lévy measures.

The finite element method for solving the pricing equations (4.4) and (4.12) has been studied by Achdou and Pironneau using adaptive mesh refinement techniques, see [2, Chaps. 4 and 5]. In one dimension, Matache et al. [91, 92] have introduced a very general wavelet-based finite element scheme to solve (4.12). This was subsequently applied to American-type contracts (cf. [90, 121]) as well as stochastic volatility models (cf. [64]). In [50, 101, 104] the wavelet-based approach was extended to multidimensional models based on sparse tensor products and wavelet compression techniques as described in [96, 101] and the references therein.

In the following we explain the basic finite element approach (cf. [2, 115]) and further illustrate the use of wavelet basis functions in this context. After localization of the original space domain  $\mathbb{R}^d$  to  $B_R$  as described above, the numerical solution of (4.12) by the finite element method is obtained in two main steps:

1. The infinite dimensional space  $\mathcal{D}(\mathcal{E})$  needs to be *discretized* by finite dimensional subspaces  $V_N \subset \mathcal{D}(\mathcal{E})$  corresponding to a finite element mesh with  $N$  degrees of freedom.
2. A *time stepping* scheme has to be applied to discretize in time.

*Space discretization* Let  $V_N \subset \mathcal{D}(\mathcal{E})$  be a subspace of dimension  $N^d := \dim V_N$  generated by a finite element basis  $\Phi := \{\phi_j : j = 1, \dots, N^d\}$  on a tensor product mesh of width  $\Delta x = \frac{2R}{N}$  on  $B_R$ . For classical examples of basis functions see e.g. [115, Chapter 5] or [23]. We use the Galerkin approach and approximate the solution  $u$  by a function  $u_N(t, x) = \sum_{j=1}^{N^d} \underline{u}_j(t)\phi_j(x) \in V_N$ . Then, for each time  $t \in [0, T]$  the semidiscrete problem of finding the coefficient vector  $\underline{u}(t)$  is an initial value problem for  $N^d$  ordinary differential equations

$$\mathbf{M} \frac{\partial}{\partial t} \underline{u}(t) + \mathbf{A} \underline{u}(t) = 0, \quad \underline{u}(0) = \underline{u}^0, \tag{4.14}$$

where  $\underline{u}^0$  denotes the coefficient vector of  $u_0$ , and  $\mathbf{M}, \mathbf{A}$  denote the mass and stiffness matrices with respect to the basis of  $V_N$ , i.e.,

$$\mathbf{M} = ((\phi_i, \phi_j))_{1 \leq i, j \leq N^d}, \quad \mathbf{A} = (\mathcal{E}(\phi_i, \phi_j))_{1 \leq i, j \leq N^d}. \tag{4.15}$$

*Time stepping using the  $\theta$ -scheme* As in the finite difference case, for the time discretization of (4.15) we consider a uniform grid with time step  $\Delta t = \frac{T}{M}$  and time

points  $t_m = m\Delta t$ ,  $m = 0, \dots, M$  for some  $M \in \mathbb{N}$ . Furthermore, we again use the  $\theta$ -scheme with  $\theta = \theta_1 = \theta_2$  to obtain

$$\frac{u^{m+1} - u^m}{\Delta t} \mathbf{M} + \theta \mathbf{A}u^{m+1} + (1 - \theta)\mathbf{A}u^m = 0, \quad m = 0, \dots, M - 1. \quad (4.16)$$

For  $\theta = 1/2$ , the scheme in (4.16) coincides with the popular Crank–Nicholson scheme. Note that, in contrast to the finite difference methods, also implicit time stepping schemes are admissible here provided one chooses a suitable (e.g. wavelet) basis for  $V_N$ ; see e.g. [91]. Furthermore, one can also use finite elements for the time discretization as in [64, 113] where an  $hp$ -discontinuous Galerkin method is used. This yields exponential convergence rates instead of only algebraic ones as in the  $\theta$ -scheme, and therefore only  $M = \log(N^d)$  time steps are required.

*Wavelet-based finite element methods* Wavelet-based finite element methods (or *wavelet methods*) provide a very general PIDE-based numerical pricing technique. The methods owe their name to the choice of a wavelet basis for the spaces  $V_N$  in the finite element method. A basic survey of wavelet-based finite element methods in finance can be found in [65], and we shall follow it here.

The motivation for applying wavelet methods rather than classical finite elements can be summarized as follows. As for the finite difference method, in high-dimensional models finite elements suffer from the “curse of dimension,” i.e., the number of degrees of freedom on a tensor product finite element mesh grows like  $\mathcal{O}(N^d)$ . For jump models the non-locality of the underlying operator  $\mathcal{A}_J$  implies that the standard finite element stiffness matrix  $\mathbf{A}$  consists of  $\mathcal{O}(N^{2d})$  non-zero entries, which is not practicable even in one dimension with small mesh widths.

For this reason, wavelet basis functions come into play. They can overcome these issues while still being easy to compute. In addition to great analytical tractability, choosing a wavelet basis for the discrete space  $V_N$  has three main advantages in practice:


- Break the curse of dimension using sparse tensor products (see e.g. [29, 96])  $\Rightarrow$  Dimension-independent complexity (up to log-factors).
- Multiscale compression of jump measure of  $X \Rightarrow$  Complexity of jump models can asymptotically be reduced to Black–Scholes complexity.
- Efficient preconditioning.

To illustrate what a wavelet basis is, we introduce the notion of a *multiscale basis* for the finite dimensional space  $V_N \subset \mathcal{D}(\mathcal{E})$ . In one dimension, suppose the mesh width  $\Delta x$  of  $V_N$  can be represented by a negative power of two,  $\Delta x = 2^{-L}$ , corresponding to a level index  $L \in \mathbb{N}_0$ . To simplify notation, write  $V_L := V_N$  with basis  $\Phi_L := \Phi$ . Then by decreasing the mesh width  $\Delta x = 2^{-L}$  one obtains a sequence of spaces

$$V_0 \subset V_1 \subset V_2 \subset \dots \subset L^2(B_R), \quad \bigcup_{L \in \mathbb{N}_0} V_L = L^2(B_R),$$

generated by bases  $\Phi_L$ ,  $L \in \mathbb{N}_0$ . Hence, bi-orthogonal *complement* or *wavelet bases*  $\Psi_L = \{\psi_{j,L} : j \in \nabla_L\}$ , with suitable index sets  $\nabla_L$ , can be constructed from the

**Fig. 1** Schematic of single-scale space  $V_L$  and its decomposition into multiscale wavelet spaces  $W_\ell$



single-scale bases  $\Phi_L$ ; for details see [36, 44, 94]. Denoting by  $W_L$  the span of  $\Psi_L$ , the spaces  $V_{L+1}$  admit a splitting

$$V_{L+1} = W_L \oplus V_L, \quad L > 0. \tag{4.17}$$

Each wavelet space  $W_L$  can be thought of as describing the increment of information when refining the finite element approximation from  $V_L$  to  $V_{L+1}$ . Furthermore, (4.17) implies that for any  $L > 0$  the finite element space  $V_L$  can be written as a direct *multilevel* sum of the wavelet spaces  $W_\ell$ ,  $\ell < L$ . Thus, any  $u_L \in V_L = V_N$  has the representation

$$u_L = \sum_{\ell=0}^{L-1} \sum_{j \in \nabla_\ell} d_{j,\ell} \psi_{j,\ell},$$

with suitable coefficients  $d_{j,\ell} \in \mathbb{R}$ . Figure 1 illustrates the decomposition of the finite element space  $V_L$ ,  $L = 4$ , spanned by continuous, piecewise linear (nodal) basis functions  $\phi_{i,L}$  into its increment spaces  $W_\ell$ ,  $\ell = 0, \dots, 3$ , spanned by wavelets  $\psi_{j,\ell}$ .

In the multidimensional setting we obtain multivariate wavelet basis functions by using tensor products. The finite element spaces  $V_L$  can then be characterized by

$$V_L = \text{span}\{\psi_{j_1,\ell_1}(x_1) \cdots \psi_{j_d,\ell_d}(x_d) : \ell_1, \dots, \ell_d \leq L, j_i \in \nabla_{\ell_i}\}.$$

Since these multivariate wavelet bases consist of products of one-dimensional wavelets, they form *hierarchical bases* as in [61]. Thus, the spaces  $V_L$  can be replaced by *sparse tensor product spaces*

$$\hat{V}_L = \text{span}\{\psi_{j_1,\ell_1}(x_1) \cdots \psi_{j_d,\ell_d}(x_d) : \ell_1 + \cdots + \ell_d \leq L, j_i \in \nabla_{\ell_i}\}.$$

In [29, 96] it is shown that, under certain smoothness assumptions on the solution  $u$  of (4.12), the sparse tensor product spaces preserve the approximation properties of the full tensor product spaces while there holds  $\hat{N} := \dim \hat{V}_L = \mathcal{O}(N |\log N|^{d-1}) \ll N^d$ . Therefore, the complexity of the finite element stiffness matrix can be reduced to  $\mathcal{O}(\hat{N}^2)$  instead of originally  $\mathcal{O}(N^{2d})$  non-zero entries.

Furthermore, wavelet basis functions give rise to certain *cancelation properties* and *norm equivalences* as illustrated in e.g. [21, 36]. One therefore obtains sharp estimates for the entries of the corresponding stiffness matrix, cf. [91, 101]. Herewith a priori compression schemes can be defined that further reduce the complexity of the stiffness matrix. The compression exploits the fact that the position of large entries in the stiffness matrix arising from a model with jumps resembles the structure of a Black–Scholes stiffness matrix. The remaining entries can a priori be proved to be negligible. Therefore, the compression scheme (asymptotically) reduces the complexity of a model with jumps to that of the Black–Scholes model.

Combining the compression scheme with the sparse tensor product spaces results in a computational complexity of  $\mathcal{O}(\hat{N})$  instead of the original  $\mathcal{O}(N^{2d})$ . It is proved that these wavelet schemes preserve stability and convergence of the classical finite element schemes, cf. [101–103].

Since the convergence of finite element methods has been studied intensively for many decades, sophisticated numerical analysis is available here. For example, in [50, 101] it is shown that when using piecewise polynomial basis functions of degree  $p \geq 1$ , the finite element scheme described above converges at rate  $p + 1$  in  $\Delta x$  (provided sufficient smoothness of the solution  $u$  of (4.4)). In particular, employing a piecewise linear basis as illustrated in Fig. 1 one obtains a second order scheme.

*American options* Similarly to the variational formulation for European contracts, for finite element implementation the partial integro-differential inequality (4.8) is reformulated into the corresponding variational inequality

$$\begin{aligned} &\text{Find } u(\tau, \cdot) \in K_{\tilde{g}_\tau} := \{v \in \mathcal{D}(\mathcal{E}) : v \geq \tilde{g}_\tau \text{ a.e.}\} \quad \text{such that} \\ &\left\langle \frac{\partial u}{\partial \tau} v - u \right\rangle_{\mathcal{D}(\mathcal{E})^*, \mathcal{D}(\mathcal{E})} + \mathcal{E}(u, v - u) \leq 0 \quad \text{a.e. in } [0, T], \text{ for all } v \in K_{\tilde{g}_\tau}, \end{aligned}$$

with  $\tilde{g}_\tau$  given by (4.9).

Choosing finite dimensional subspaces  $V_N \subset V$ , as above, to discretize in space and applying the  $\theta$ -scheme defined in (4.16) with  $\theta = 1$  to discretize in time leads to a sequence of matrix linear complementary problems (LCPs)

$$\text{Given } \underline{u}^0, \text{ find } \underline{u}^m \in \underline{K} := \{\underline{v} \in \mathbb{R}^{\dim V_N} : \underline{v} \geq \underline{\tilde{g}}_t\} \quad \text{such that for all } \underline{v} \in \underline{K}, \tag{4.18}$$

$$(\underline{v} - \underline{u}^{m+1})^\top (\mathbf{M} + \Delta t \mathbf{A}) \underline{u}^{m+1} \geq (\underline{v} - \underline{u}^{m+1})^\top \mathbf{M} \underline{u}^m, \quad m = 0, \dots, M - 1,$$

with  $\mathbf{M}, \mathbf{A}$  as in (4.15) and where  $\underline{u}^0$  denotes the coefficient vector of  $u_0$ . As already described in (3.3) this discretization can be interpreted as approximating the value of the American option by a sequence of Bermudan option values.

For a large number  $N$  of degrees of freedom, standard solution methods like projected SOR [43] for the matrix LCP (4.18) are not suitable, since their rate of convergence depends on  $N$ . The wavelet-based solution algorithm suggested in [90] relies on a fixed point iteration where in each iteration step a  $V_L$ -projection  $P_{\underline{K}}$  onto the convex cone  $\underline{K}$  has to be realized. Due to norm equivalences of the wavelet basis, the outer fixed point iteration convergences at a rate independent of the number of degrees of freedom. The projection  $P_{\underline{K}}$  is based on a wavelet generalization of the classical Cray algorithm [43].

## 5 Comparison

So far we have presented the setup and general methodology of the major numerical schemes for multidimensional Lévy models. In this final section we explain the main differences and problem-dependent advantages of these methods.

*Admissible exotic contracts* The applicability of the different numerical methods to exotic contracts in Lévy models is essentially subject to the same challenges as in the classical Black–Scholes model.

As we have already seen in the above sections, nowadays *American contracts* can be handled more or less efficiently by all described methods. For Monte Carlo however this was long thought to be an impossible task, since it is based on simulation forward in time. In mesh-based methods one employs backward time stepping. With this, the efficient pricing of American contracts is straightforward. Therefore, Monte Carlo schemes are still considered inferior to mesh-based methods when applied to American contracts.

Contracts with *discontinuous payoffs* such as digital options do not pose any additional challenges for the finite element method, since it solves the variational problem (4.12), which is  $L^2$ -based. Under certain conditions, however, discontinuous payoffs can result in a decreased rate of convergence for Monte Carlo methods; for details see e.g. [25].

Pricing of *barrier options* can be handled very easily with a PIDE-based approach, since one simply has to restrict the price-space domain of the discretization to that of the barrier and impose suitable boundary conditions. Here, very complex multidimensional barriers can be handled instantly. If the barrier is monitored continuously, pricing barrier options is somewhat more complicated for Monte Carlo methods, since we simulate forward in time and one cannot see what happens in between sampling dates. Therefore, straightforward Monte Carlo methods overestimate the prices of knock-out options and underestimate those of knock-in contracts. Correction techniques have been considered for several years now; see e.g. [8, 28, 106]. In the case where the barrier is monitored at discrete days (which is common in practice) one may choose the sampling dates to coincide with the observation dates and hence the bias in the Monte Carlo estimates vanishes. For Fourier methods the Wiener–Hopf factorization can be used; see e.g. [19].

Monte Carlo methods are easily applied to *path-dependent derivatives* such as Asian or lookback options, since sample paths are simulated forward in time and

the history at each time step is known. One only has to take into account that, as for barrier options, Monte Carlo methods might result in biased estimates if the contract is monitored continuously. Since mesh-based methods solve backward in time, they are intrinsically not well suited for path-dependent options. However, several methods have been introduced to provide PIDE-based tools for such contracts. For example, to price Asian options one may introduce an additional variable, i.e., increase the problem dimension, to handle the averaging of the underlying's prices; see e.g. [119, 123].

For all numerical methods, a continuously paid *dividend*  $q$  can simply be handled by changing the interest rate  $r \mapsto r - q$ . If the dividend is paid at discrete points in time it can still be handled easily by Monte Carlo methods, since one is proceeding forward in time and payment of a dividend results in an immediate decrease of the underlying's value. For Fourier and PIDE-based techniques the situation of discretely paid dividends is usually more complicated, since one needs to enforce additional no-arbitrage conditions; see e.g. [122].

*Admissible market models* Monte Carlo methods are applicable to any Lévy process  $X$  as long as there is an efficient way of simulating its trajectories. Furthermore, the simulation techniques illustrated in Sect. 2 provide a very general framework for Monte Carlo simulations applicable to more exotic models. Such simulations can however be rather involved for general multidimensional models. They have to be considered separately for each model.

In contrast to constructing a new simulation scheme for each process, PIDE-based methods provide a standard approach whenever the PIDE (4.4) admits a unique solution and the Lévy measure  $\nu$  of  $X$  is available in a suitable form, i.e., if the density or tail integrals of  $\nu$  are known explicitly. These requirements have been proved for all major one-dimensional models (see [40, 91]) and multidimensional Lévy copula models (see [50, 104]). One major advantage of the PIDE-based methods is that changing the market model only amounts to changing the stiffness matrix  $\mathbf{A}$  in (4.15) and (4.16). The matrix  $\mathbf{A}$  needs to be assembled only once for each model and can be re-used for different payoffs. PIDE-based methods are hence well suited for the analysis of model risk.

As long as the characteristic function is known (see [114] for an overview) Fourier methods using the fast Fourier transformation provide a very efficient standardized approach in moderate dimensions. Except for driving processes with independent marginals or subordinated Brownian motion, however, the characteristic function is generally not known in Lévy models of dimension  $d > 1$ .

*Implementation* It is well known that in the Black–Scholes setting one advantage of the Monte Carlo method is its intuitive and rather simple implementation. In fact for the one-dimensional Black–Scholes model also Fourier, finite difference and classical finite element methods can be implemented by straightforward standard techniques. A great amount of fundamental literature is available in this case; see e.g. [2, 22, 115]. However, for (multidimensional) Lévy models more work is required for all methods. Even though the convergence rate of the Monte Carlo method is dimension-independent, its implementation in general requires special considerations for different techniques; see e.g. [117] for multidimensional simulation based on copulas.

As illustrated in the previous sections, all naive grid-based techniques such as standard Fourier and PIDE methods suffer from an exponential growth of complexity with the dimension. Applying sparse grids (cf. [29, 50]) or the so-called multigrid technique (cf. [17, 105]) to overcome this issue requires the implementation of a non-trivial data structure to handle the degrees of freedom efficiently. Furthermore, being the most general grid-based method, the implementation of wavelet-based finite elements requires additional handling of the compression techniques. For implementation details see [124].

*Model sensitivities and Greeks* Calculating price sensitivities (e.g. the Greeks) is a central modeling and computational task for risk management and hedging. We distinguish two classes: Sensitivities of the price  $V$  in (1.2) to variations of a model parameter, like the Greek Vega  $\partial_\sigma V$ ; and sensitivities of  $V$  to variations of the state space such as the Greek Delta  $\partial_S V$ . Mesh-based methods are known to be well suited for the fast and accurate calculation of sensitivities whereas Monte Carlo methods are facing a certain challenge in this respect.

For PIDE-based methods, for instance, suppose the market model and hence the operator  $\mathcal{A} = \mathcal{A}_{BS} + \mathcal{A}_J$  in (4.4) depends on some model parameter  $\eta$ . We want to calculate the sensitivity of the solution  $u$  of (4.4) with respect to  $\eta$ . To this end, write  $u(\eta_0)$  for a fixed realization  $\eta_0$  of  $\eta$  in order to emphasize the dependence of  $u$  on  $\eta_0$  in (4.4). Then, as shown in [66], the derivative  $\tilde{u}(\delta\eta)$  of  $u$  with respect to  $\eta$ , i.e.,  $\tilde{u}(\delta\eta) := \lim_{s \rightarrow 0^+} \frac{1}{s}(u(\eta_0 + s\delta\eta) - u(\eta_0))$ , is the solution of the PIDE

$$\frac{\partial \tilde{u}(\delta\eta)}{\partial \tau} + \mathcal{A}(\eta_0)\tilde{u}(\delta\eta) = -D_\eta \mathcal{A}u(\eta_0), \quad \tilde{u}(\delta\eta)(0, \cdot) = 0 \quad \text{in } \mathbb{R}^d,$$

where  $D_\eta \mathcal{A}$  is the derivative of  $\mathcal{A}$  with respect to  $\eta$ . Therefore, the derivative of  $u$  with respect to  $\eta$  can be obtained as a solution of the same PIDE as the price  $u$  itself, where now the right hand side depends on  $u$ . Thus, sensitivities with respect to model parameters can be calculated with the same computational effort as the price itself. Furthermore, it is shown in [66] that all computed sensitivities converge with the same rate as the original price  $u$ . For sensitivities with respect to a variation of the state space, a finite difference-like differentiation procedure is presented in [66] which allows to obtain the sensitivities from the finite element forward price with the same convergence rate but without additional work.

Using a similar approach, also Fourier methods are capable of calculating sensitivities with respect to state space variation efficiently; see e.g. [4] for some numerical examples.

For Monte Carlo methods the computing time required for the calculation of sensitivities can be significantly greater than the time needed to calculate the prices themselves (to the same accuracy). For example, suppose one wants to estimate the Delta  $\partial_S V(S)$ . Then, one can compute a Monte Carlo estimator for  $V(0, S + \delta)$  for some small perturbation  $\delta$ . With this, the Delta  $\partial_S V(S)$  may then be approximated by a forward finite difference estimator  $(V(0, S + \delta) - V(0, S))/\delta$ . In the Black–Scholes setting, it is proved in [60] that the best possible convergence rate for such an approximation is  $N^{-1/4}$  if the simulations of the two estimators are drawn independently



(here  $N$  denotes the number of sample paths). By replacing the forward finite difference estimator with a central difference estimator this rate can be improved to  $N^{-1/3}$ . Furthermore, using common random numbers for both Monte Carlo estimators, one can obtain  $N^{-1/2}$ ; see [59, 60, 77]. However, an important drawback of using common random numbers is that it may perform very poorly when the payoff is not sufficiently smooth, e.g. for digital options. For a more sophisticated method of calculating the Delta we refer to e.g. [57].

Also in the Black–Scholes setting, Malliavin calculus has become a popular tool for the Monte Carlo calculation of certain Greeks. This approach is based on the observation that most of the sensitivities of interest can be expressed as  $\mathbb{E}[g(S_T)\pi]$ , where  $\pi$  is a random variable depending on the sensitivity to be calculated. For details we refer to [45, 53, 54, 81] and the references therein. In case the probability density of  $S$  is known and differentiable, the likelihood ratio method (see e.g. [58]) may be used to replace the Malliavin calculus. See also [58, Sect. 7.2] for the pathwise differentiation technique.

*Extension beyond Lévy models* Recently, certain market models have been introduced where the driving process is still Markovian but not Lévy anymore. In [33] for instance, the underlying  $S$  is modeled by a time-inhomogeneous so-called Sato process. Furthermore, in certain stochastic volatility models such as the BNS model [13] the driving process is not stationary anymore.

Both time-inhomogeneity and non-stationarity can be handled easily by the PIDE-based methods. Non-stationarity can be treated by efficient evaluation of the entries of the stiffness matrices  $\mathbf{A}$  in (4.10) and (4.14). Furthermore, if the stiffness matrix  $\mathbf{A}$  in (4.10), (4.14) is time-dependent, then the time stepping scheme (4.16) can still be applied with  $\mathbf{A} = \mathbf{A}(t)$  evaluated at each time step.

If the characteristic function of a non-stationary process is known explicitly (as e.g. for certain instances of the BNS model, see [114]), Fourier methods are applicable. Also if the process is time-inhomogeneous, Fourier methods can be applied as in [33].

Monte Carlo discretization can easily be extended to non-stationary or time-inhomogeneous processes, e.g. using suitable Markov chain approximations. With such a rather naive approach, however, the computational complexity grows significantly.

## References

1. Achdou, Y., Franchi, B., Tchou, N.: A partial differential equation connected to option pricing with stochastic volatility: regularity results and discretization. *Math. Comput.* **74**(251), 1291–1322 (2005) (electronic)
2. Achdou, Y., Pironneau, O.: *Computational Methods for Option Pricing*. *Frontiers in Applied Mathematics*, vol. 30. SIAM, Philadelphia (2005)
3. Almendral, A., Oosterlee, C.: Numerical valuation of options with jumps in the underlying. *Appl. Numer. Math.* **53**, 1–18 (2005)
4. Almendral, A., Oosterlee, C.: Highly accurate evaluation of European and American options under the variance gamma process. *J. Comput. Finance* **10**, 21–42 (2006)
5. Amin, K.: Jump diffusion option valuation in discrete time. *J. Finance* **48**, 1833–1863 (1993)

6. Amin, K., Khanna, A.: Convergence of American option values from discrete- to continuous-time financial models. *Math. Finance* **4**, 289–304 (1994)
7. Andersen, L., Andreasen, J.: Jump-diffusion processes: volatility smile fitting and numerical methods for pricing. *Rev. Deriv. Res.* **4**, 231–262 (2000)
8. Andersen, L., Brotherton-Ratcliffe, R.: Exact exotics. *Risk* **9**, 85–89 (1996)
9. Asmussen, S., Glynn, P.: *Stochastic Simulation: Algorithms and Analysis*. Stochastic Modelling and Applied Probability, vol. 57. Springer, New York (2007)
10. Asmussen, S., Rosiński, J.: Approximations of small jumps of Lévy processes with a view towards simulation. *J. Appl. Probab.* **38**, 482–493 (2001)
11. Avramidis, A., L'Ecuyer, P.: Efficient Monte Carlo and quasi-Monte Carlo option pricing under the variance gamma model. *Manag. Sci.* **52**(12), 1930–1944 (2006)
12. Barndorff-Nielsen, O.: Normal inverse Gaussian distributions and stochastic volatility modelling. *Scand. J. Stat.* **24**(1), 1–13 (1997)
13. Barndorff-Nielsen, O., Shephard, N.: Non-Gaussian Ornstein–Uhlenbeck based models and some of their uses in financial economics. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **63**, 167–241 (2001)
14. Bensoussan, A., Lions, J.L.: *Applications of Variational Inequalities in Stochastic Control*. Studies in Mathematics and its Applications, vol. 12. North-Holland, Amsterdam (1982). Translated from the French
15. Bensoussan, A., Lions, J.L.: *Impulse Control and Quasivariational Inequalities*. Gauthier-Villars, Montrouge (1984). Translated from the French by J.M. Cole
16. Bertoin, J.: *Lévy Processes*. Cambridge University Press, Cambridge (1996)
17. Bin Zubair, H., Oosterlee, C.W., Wienands, R.: Multigrid for high-dimensional elliptic partial differential equations on non-equidistant grids. *SIAM J. Sci. Comput.* **29**, 1613–1636 (2007) (electronic)
18. Black, F., Scholes, M.: The pricing of options and corporate liabilities. *J. Polit. Econ.* **81**, 637–659 (1973)
19. Boyarchenko, S., Levendorskiĭ, S.: *Non-Gaussian Merton–Black–Scholes Theory*. Advanced Series on Statistical Science & Applied Probability, vol. 9. World Scientific, River Edge (2002)
20. Boyarchenko, S., Levendorskiĭ, S.: On errors and bias of Fourier transform methods in quadratic term structure models. *Int. J. Theor. Appl. Finance* **10**, 273–306 (2007)
21. Bramble, J.H., Cohen, A., Dahmen, W.: *Multiscale Problems and Methods in Numerical Simulations*. Lecture Notes in Mathematics, vol. 1825. Springer, Berlin (2003)
22. Brandimarte, P.: *Numerical Methods in Finance and Economics*, 2nd edn. Statistics in Practice. Wiley-Interscience, Hoboken (2006). A MATLAB<sup>®</sup>-based introduction
23. Brenner, S., Scott, L.: *The Mathematical Theory of Finite Element Methods*, 2nd edn. Texts in Applied Mathematics, vol. 15. Springer, New York (2002)
24. Briani, M., Natalini, R., Russo, G.: Implicit-explicit numerical schemes for jump–diffusion processes. *Calcolo* **44**, 33–57 (2007)
25. Broadie, M., Glasserman, P.: Estimating security price derivatives using simulation. *Manag. Sci.* **42**, 269–285 (1996)
26. Broadie, M., Glasserman, P.: Pricing American-style securities using simulation. *J. Econ. Dyn. Control* **21**, 1323–1352 (1997)
27. Broadie, M., Glasserman, P., Jain, G.: Enhanced Monte Carlo estimates for American option prices. *J. Deriv.* **5**, 25–44 (1997)
28. Broadie, M., Glasserman, P., Kou, S.: A continuity correction for discrete barrier options. *Math. Finance* **7**, 325–349 (1997)
29. Bungartz, H.J., Griebel, M.: A note on the complexity of solving Poisson's equation for spaces of bounded mixed derivative. *J. Complex.* **15**, 167–199 (1999)
30. Bungartz, H.J., Griebel, M.: Sparse grids. *Acta Numer.* **13**, 147–269 (2004)
31. Bungartz, H.J., Griebel, M., Röschke, D., Zenger, C.: Two proofs of convergence for the combination technique for the efficient solution of sparse grid problems. In: Keyes, D., Xu, J. (eds.) *Domain Decomposition Methods in Scientific and Engineering Computing*, University Park, PA, 1993. *Contemp. Math.*, vol. 180, pp. 15–20. Amer. Math. Soc., Providence (1994)
32. Carr, P., Geman, H., Madan, D., Yor, M.: The fine structure of assets returns: an empirical investigation. *J. Bus.* **75**, 305–332 (2002)
33. Carr, P., Geman, H., Madan, D., Yor, M.: Self-decomposability and option pricing. *Math. Finance* **17**, 31–57 (2007)
34. Carr, P., Madan, D.: Option valuation and the fast Fourier transform. *J. Comput. Finance* **2**(4), 61–73 (1999)

35. Chen, N., Glasserman, P.: Additive and multiplicative duals for American option pricing. *Finance Stoch.* **11**, 153–179 (2007)
36. Cohen, A.: *Numerical Analysis of Wavelet Methods*. Elsevier, Amsterdam (2003)
37. Cohen, S., Rosiński, J.: Gaussian approximation of multivariate Lévy processes with applications to simulation of tempered stable processes. *Bernoulli* **13**, 195–210 (2007)
38. Cont, R., Tankov, P.: *Financial Modelling with Jump Processes*. Chapman & Hall/CRC Financial Mathematics Series. Chapman & Hall/CRC, Boca Raton (2004)
39. Cont, R., Voltchkova, E.: A finite difference scheme for option pricing in jump diffusion and exponential Lévy models. *SIAM J. Numer. Anal.* **43**, 1596–1626 (2005)
40. Cont, R., Voltchkova, E.: Integro-differential equations for option prices in exponential Lévy models. *Finance Stoch.* **9**, 299–325 (2005)
41. Cooley, J., Tukey, J.: An algorithm for the machine calculation of complex Fourier series. *Math. Comput.* **19**, 297–301 (1965)
42. Cox, J., Ross, S., Rubinstein, M.: Option pricing: A simplified approach. *J. Financ. Econ.* **48**, 289–304 (1993)
43. Cryer, C.: The solution of a quadratic programming problem using systematic overrelaxation. *SIAM J. Control* **9**, 385–392 (1971)
44. Dahmen, W., Kunoth, A., Urban, K.: Biorthogonal spline wavelets on the interval—stability and moment conditions. *Appl. Comput. Harmon. Anal.* **6**, 259–302 (1999)
45. Davis, M., Johansson, M.: Malliavin Monte Carlo Greeks for jump diffusions. *Stoch. Process. Appl.* **116**, 101–129 (2006)
46. Delbaen, F., Schachermayer, W.: A general version of the fundamental theorem of asset pricing. *Math. Ann.* **300**, 463–520 (1994)
47. Duffie, D., Pan, J., Singleton, K.: Transform analysis and asset pricing for affine jump-diffusions. *Econometrica* **68**, 1343–1376 (2000)
48. Duffy, D.J.: *Finite Difference Methods in Financial Engineering*. Wiley Finance Series. Wiley, Chichester (2006). A partial differential equation approach, with 1 CD-ROM (Windows, Macintosh and UNIX)
49. Eberlein, E., Prause, K.: The generalized hyperbolic model: financial derivatives and risk measures. In: Geman, H., Madan, D., Pliska, S., Vorst, T. (eds.) *Mathematical finance—Bachelier Congress, 2000 (Paris)*. Springer Finance, vol. 44, pp. 245–267. Springer, Berlin (2002)
50. Farkas, W., Reich, N., Schwab, C.: Anisotropic stable Lévy copula processes—analytical and numerical aspects. *Math. Models Methods Appl. Sci.* **17**, 1405–1443 (2007)
51. Figlewski, S., Gao, B.: The adaptive mesh model: a new approach to efficient option pricing. *J. Financ. Econ.* **53**, 313–351 (1999)
52. Forsyth, P.A., Vetzal, K.R.: Quadratic convergence for valuing American options using a penalty method. *SIAM J. Sci. Comput.* **23**, 2095–2122 (2002) (electronic)
53. Fournié, E., Lasry, J.M., Lebuchoux, J., Lions, P.L.: Applications of Malliavin calculus to Monte Carlo methods in finance, II. *Finance Stoch.* **5**, 201–236 (2001)
54. Fournié, E., Lasry, J.M., Lebuchoux, J., Lions, P.L., Touzi, N.: Applications of Malliavin calculus to Monte Carlo methods in finance. *Finance Stoch.* **3**, 391–412 (1999)
55. Fu, M., Laprise, S., Madan, D., Su, Y., Wu, R.: Pricing American options: a comparison of Monte Carlo simulation approaches. *J. Comput. Finance* **4**, 39–88 (2001)
56. Geske, R., Johnson, H.: The American put option valued analytically. *J. Finance* **39**, 1511–1524 (1984)
57. Giles, M., Glasserman, P.: Smoking adjoints: fast Monte Carlo Greeks. *Risk* **19**, 88–92 (2006)
58. Glasserman, P.: *Monte Carlo Methods in Financial Engineering*. Applications of Mathematics, vol. 53. Springer, New York (2004)
59. Glasserman, P., David, D.Y.: Some guidelines and guarantees for common random numbers. *Manag. Sci.* **38**, 884–908 (1992)
60. Glynn, P.W., Iglehart, D.L.: Importance sampling for stochastic simulations. *Manag. Sci.* **35**, 1367–1392 (1989)
61. Griebel, M., Oswald, P.: Tensor product type subspace splittings and multilevel iterative methods for anisotropic problems. *Adv. Comput. Math.* **4**, 171–206 (1995)
62. Haugh, M., Kogan, L.: Pricing American options: a duality approach. *Oper. Res.* **52**, 258–270 (2004)
63. Heston, S., Zhou, G.: On the rate of convergence of discrete-time contingent claims. *Math. Finance* **10**, 53–75 (2000)
64. Hilber, N., Matache, A.M., Schwab, C.: Sparse wavelet methods for option pricing under stochastic volatility. *J. Comput. Finance* **8**(4), 1–42 (2005)

65. Hilber, N., Reich, N., Winter, C.: Wavelet methods. In: Cont, R. (ed.) *Encyclopedia of Quantitative Finance*. Wiley, Chichester (to appear). <http://www.wiley.com/legacy/wileychi/eqf/>
66. Hilber, N., Schwab, C., Winter, C.: Variational sensitivity analysis of parametric Markovian market models. In: *Special issue on Mathematics of Finance, Banach Center Publ.*, vol. 83, pp. 85–106 (2008)
67. Jackson, K., Jaimungal, S., Surkov, V.: Fourier space time-stepping for option pricing with Lévy models. Tech. rep., SSRN (2007). <http://ssrn.com/abstract=1020209>
68. Jacod, J.: The Euler scheme for Lévy driven stochastic differential equations: limit theorems. *Ann. Probab.* **32**, 1830–1872 (2004)
69. Jacod, J., Kurtz, T.G., Méléard, S., Protter, P.: The approximate Euler method for Lévy driven stochastic differential equations. *Ann. Inst. H. Poincaré Probab. Stat.* **41**, 523–558 (2005)
70. Jacod, J., Shiryaev, A.: *Limit Theorems for Stochastic Processes*, 2nd edn. *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, vol. 288. Springer, Berlin (2003)
71. Kallsen, J., Tankov, P.: Characterization of dependence of multidimensional Lévy processes using Lévy copulas. *J. Multivar. Anal.* **97**, 1551–1572 (2006)
72. Kawai, R.: Adaptive Monte Carlo variance reduction for Lévy processes with two-time-scale stochastic approximation. *Methodol. Comput. Appl. Probab.* **10**, 199–223 (2008)
73. Kloeden, P., Platen, E.: *Numerical Solution of Stochastic Differential Equations*, 3rd edn. *Applications of Mathematics*, vol. 23. Springer, Berlin (1999)
74. Kou, G.: A jump diffusion model for option pricing. *Manag. Sci.* **48**, 1086–1101 (2002)
75. Kushner, H., Dupuis, P.: *Numerical Methods for Stochastic Control Problems in Continuous Time*, 2nd edn. *Applications of Mathematics (New York)*, vol. 24. Springer, New York (2001)
76. L'Ecuyer, P.: Quasi-Monte Carlo methods with applications in finance. *Finance Stoch.* **13**, 307–350 (2009)
77. L'Ecuyer, P., Perron, G.: On the convergence rates of IPA and FDC derivative estimators. *Oper. Res.* **42**, 643–656 (1994)
78. Lee, R.: Option pricing by transform methods: Extensions, unification, and error control. *J. Comput. Finance* **7**(3), 51–86 (2004)
79. Lemieux, C., L'Ecuyer, P.: On the use of Quasi-Monte Carlo methods in computational finance. In: Alexandrov, V., Dongarra, J., Juliano, B., Renner, R., Tan, C. (eds.) *ICCS '01: Proceedings of the International Conference on Computational Sciences, Part I*, pp. 607–618. Springer, London (2001)
80. Leobacher, G.: Stratified sampling and Quasi-Monte Carlo simulation of Lévy processes. *Monte Carlo Methods Appl.* **12**, 231–238 (2006)
81. León, J., Solé, J., Utzet, F., Vives, J.: On Lévy processes. *Malliavin calculus and market models with jumps. Finance Stoch.* **6**, 197–225 (2002)
82. Levendorskiĭ, S.: Early exercise boundary and option prices in Lévy driven models. *Quant. Finance* **4**, 525–547 (2004)
83. Lewis, A.: A simple option formula for general jump-diffusion and other exponential Lévy processes. Tech. rep., SSRN (2001). <http://papers.ssrn.com/abstract=282110>
84. Longstaff, F., Schwartz, E.: Valuing American options by simulation. *Rev. Financ. Stud.* **14**, 113–147 (2001)
85. Lord, R., Fang, F., Bervoets, F., Oosterlee, C.: A fast method for pricing early-exercise options with the FFT. In: Shi, Y., van Albada, G., Dongarra, J., Sloot, P. (eds.) *ICCS '07, Part II*, pp. 415–422. Springer, Berlin (2007)
86. Luciano, E., Schoutens, W.: A multivariate jump-driven financial asset model. *Quant. Finance* **6**, 385–402 (2006)
87. Madan, D., Carr, P., Chang, E.: The variance gamma process and option pricing. *Eur. Finance Rev.* **2**, 79–105 (1998)
88. Maller, R.A., Solomon, D.H., Szimayer, A.: A multinomial approximation for American option prices in Lévy process models. *Math. Finance* **16**, 613–633 (2006)
89. Maruyama, G.: Continuous Markov processes and stochastic equations. *Rend. Circ. Mat. Palermo* **4**, 48–90 (1955)
90. Matache, A.M., Nitsche, P.A., Schwab, C.: Wavelet Galerkin pricing of American options on Lévy driven assets. *Quant. Finance* **5**, 403–424 (2005)
91. Matache, A.M., Petersdorff, T., Schwab, C.: Fast deterministic pricing of options on Lévy driven assets. *M2AN Math. Model. Numer. Anal.* **38**, 37–71 (2004)
92. Matache, A.M., Schwab, C., Wihler, T.P.: Linear complexity solution of parabolic integro-differential equations. *Numer. Math.* **104**, 69–102 (2006)

93. Mulinacci, S.: An approximation of American option prices in a jump-diffusion model. *Stoch. Process. Appl.* **62**, 1–17 (1996)
94. Nguyen, H., Stevenson, R.: Finite elements on manifolds. *IMA J. Numer. Math.* **23**, 149–173 (2003)
95. Papapantoleon, A.: Applications of semimartingales and Lévy processes in finance: duality and valuation. PhD thesis, University of Freiburg (2006). <http://www.freidok.uni-freiburg.de/volltexte/2919/>
96. von Petersdorff, T., Schwab, C.: Numerical solution of parabolic equations in high dimensions. *M2AN Math. Model. Numer. Anal.* **38**, 93–127 (2004)
97. Pooley, D.M., Forsyth, P.A., Vetzal, K.R.: Numerical convergence properties of option pricing PDEs with uncertain volatility. *IMA J. Numer. Anal.* **23**, 241–267 (2003)
98. Press, W., Teukolsky, S., Vetterling, W., Flannery, B.: *Numerical Recipes in C*, 2nd edn. Cambridge University Press, Cambridge (1992)
99. Prigent, J.L.: *Weak Convergence of Financial Markets*. Springer Finance. Springer, Berlin (2003)
100. Raible, S.: Lévy processes in finance: theory, numerics, and empirical facts. PhD thesis, University of Freiburg (2000). <http://www.freidok.uni-freiburg.de/volltexte/51/>
101. Reich, N.: Wavelet compression of anisotropic integrodifferential operators on sparse tensor product spaces. PhD thesis 17661, ETH Zürich (2008). <http://e-collection.ethbib.ethz.ch/view/eth:30174>
102. Reich, N.: Wavelet compression of anisotropic integrodifferential operators on sparse tensor product spaces (submitted). Research report No. 2008-26, Seminar for Applied Mathematics, ETH Zürich (2008). <http://www.sam.math.ethz.ch/reports/2008/26>
103. Reich, N.: Wavelet compression of integral operators on sparse tensor spaces: Construction, consistency and asymptotically optimal complexity (submitted). Research report No. 2008-24, Seminar for Applied Mathematics, ETH Zürich (2008). <http://www.sam.math.ethz.ch/reports/2008/24>
104. Reich, N., Schwab, C., Winter, C.: On Kolmogorov equations for anisotropic multivariate Lévy processes (submitted). Research report 2008-3, Seminar for Applied Mathematics, ETH Zürich (2008). <http://www.sam.math.ethz.ch/reports/2008/03>
105. Reisinger, C., Wittum, G.: Efficient hierarchical approximation of high-dimensional option pricing problems. *SIAM J. Sci. Comput.* **29**, 440–458 (2007)
106. Ribeiro, C., Webber, N.: Correcting for simulation bias in Monte Carlo methods to value exotic options in models driven by Lévy processes. *Appl. Math. Finance* **13**, 333–352 (2006)
107. Rogers, L.: Monte Carlo valuation of American options. *Mathematical Finance* **12**, 271–286 (2002)
108. Rosiński, J.: Series representations of Lévy processes from the perspective of point processes. In: Barndorff-Nielsen, O., Mikosch, T., Resnick, S. (eds.) *Lévy processes*, pp. 401–415. Birkhäuser, Boston (2001)
109. Rosiński, J.: Tempering stable processes. *Stoch. Process. Appl.* **117**, 677–707 (2007)
110. Rubenthaler, S.: Numerical simulation of the solution of a stochastic differential equation driven by a Lévy process. *Stoch. Process. Appl.* **103**, 311–349 (2003)
111. Sanfelici, S.: Galerkin infinite element approximation for pricing barrier options and options with discontinuous payoff. *Decis. Econ. Finance* **27**, 125–151 (2004)
112. Sato, K.I.: *Lévy Processes and Infinitely Divisible Distributions*. Cambridge Studies in Advanced Mathematics, vol. 68. Cambridge University Press, Cambridge (1999)
113. Schötzau, D., Schwab, C.: *hp*-discontinuous Galerkin time-stepping for parabolic problems. *C. R. Acad. Sci. Paris Sér. I Math.* **333**(12), 1121–1126 (2001)
114. Schoutens, W.: *Lévy Processes in Finance*. Wiley, Chichester (2003)
115. Seydel, R.: *Tools for Computational Finance*, 3rd edn. Universitext/Springer, Berlin (2006)
116. Tankov, P.: Dependence structure of Lévy processes with applications to risk management. Rapport Interne No. 502, CMAPX École Polytechnique (2003). <http://www.cmap.polytechnique.fr/preprint>
117. Tankov, P.: Simulation and option pricing in Lévy copula models. Tech. rep., Laboratoire de Probabilités et Modèles Aléatoires, Université Paris VII (2005). [http://people.math.jussieu.fr/~tankov/levycopulas\\_ima.pdf](http://people.math.jussieu.fr/~tankov/levycopulas_ima.pdf)
118. Topper, J.: Worst case pricing of rainbow options. *J. Mong. Math. Soc.* **5**, 96–109 (2001)
119. Večeř, J., Xu, M.: Pricing Asian options in a semimartingale model. *Quant. Finance* **4**, 170–175 (2004)
120. Webber, N.: Simulation methods with Lévy processes. In: Kyprianou, A., Schoutens, W., Willmott, P. (eds.) *Exotic Option Pricing and Advanced Lévy Models*, pp. 29–49. Wiley, Chichester (2005)
121. Wilhelm, M., Winter, C.: Finite element valuation of swing options. *J. Comput. Finance* **11**(3), 107–132 (2008)
122. Willmott, P.: *Derivatives: The Theory and Practice of Financial Engineering*. Wiley, Chichester (1998)

123. Wilmott, P., Dewynne, J., Howison, S.: *Option Pricing: Mathematical Models and Computation*. Oxford Financial Press, Oxford (1993)
124. Winter, C.: *Wavelet Galerkin schemes for option pricing in multidimensional Lévy models*. PhD thesis 18221, ETH Zürich (2009). <http://e-collection.ethbib.ethz.ch/view/eth:41555>
125. Zenger, C.: Sparse grids. In: *Parallel algorithms for partial differential equations*, Kiel, 1990. *Notes Numer. Fluid Mech.*, vol. 31, pp. 241–251. Vieweg, Braunschweig (1991)