

Amaurosis after spine surgery: survey of the literature and discussion of one case

Stephan Zimmerer · Markus Koehler ·
Stephanie Turtschi · Anja Palmowski-Wolfe ·
Thierry Girard

Received: 1 August 2009/Revised: 18 May 2010/Accepted: 14 August 2010/Published online: 1 September 2010
© Springer-Verlag 2010

Abstract Postoperative vision loss (POVL) associated with spine surgery is a well known, albeit very rare complication. POVL incidence after spinal surgery ranges from 0.028 to 0.2%; however, due to the increase in number and duration of annual complex spinal operations, the incidence may increase. Origin and pathogenesis of POVL remain frequently unknown. A 73-year-old patient presented with lumbar disc herniation with associated neurological deficits after conservative pre-treatment at a peripheral hospital. Known comorbidities included arterial hypertension, moderate arterial sclerosis, diabetes mellitus type 2, mildly elevated blood lipids and treated prostate gland cancer. During lumbar spine surgery in modified prone position the patient presented with an acute episode of severe hypotension, which required treatment with catecholamines and Trendelenburg positioning. Three hours postoperatively, a visual loss in the right eye occurred, resulting in a complete amaurosis. Anti-hypertensive medication, arteriosclerosis and intraoperative hypotension are possible causes for the POVL. Intraoperative administration of catecholamines and Trendelenburg positioning for treatment of systemic hypotension might further

compromise ocular perfusion. In patients with comorbidities compromising arterial blood pressure, blood circulation and microcirculation, POVL must be considered as a severe postoperative complication. It is recommended to inform patients about such complications and obtain preoperative informed consent regarding POVL. Any recent modification of antihypertensive medication must be reported and analysed for potential intraoperative hemodynamic consequences, prior to spine surgery in prone position.

Keywords Amaurosis · Blood pressure · Spinal surgery · Prone position · Postoperative vision loss (POVL)

Introduction

Postoperative vision loss (POVL) associated with spine surgery in prone position is a well known yet very rare complication [1, 2]. The exact origin and pathogenesis are still unknown in many cases [3, 4]. The most common ophthalmologic diagnosis associated with POVL is ischemic optic neuropathy (ION) [5, 6]. Other causes include central retinal artery occlusion (CRAO), cortical blindness as well as unknown aetiology [5, 6]. A practice advisory for perioperative visual loss associated with spine surgery is provided e.g. by the “American Society of Anesthesiologists” [7].

Survey of literature

There are several reports of vision impairment after surgery in prone position or cardiac surgery. The incidence of POVL after non-ocular surgeries has been reported as 0.002% among all surgeries and reaches 0.2% among cardiac and spine surgeries [8–11]. Due to an increase in number and duration of annual complex spinal operations,

S. Zimmerer (✉)
Department of Neurosurgery, University Hospital Basel,
Spitalstrasse 21, 4031 Basel, Switzerland
e-mail: zimmerers@uhbs.ch

M. Koehler
Bethesda Spital Basel, Gellertstr. 144, 4052 Basel, Switzerland

S. Turtschi · A. Palmowski-Wolfe
Department of Ophthalmology, University Hospital Basel,
Spitalstrasse 21, 4031 Basel, Switzerland

T. Girard
Department of Anaesthesia, University Hospital Basel,
Spitalstrasse 21, 4031 Basel, Switzerland

POVL incidence may increase [1, 6, 12–14]. In a recently published retrospective population-based American study with data of 4,728,815 patients over the years 1997–2002, the overall incidence of visual disturbance after spine surgery was 0.094% [15]. In addition to collections of up to 93 cases from several groups, there are a number of single case reports. Within these case reports, 27.3% refer to thoracic [16–18], 18.2% to lumbar [19–21] and 54.5% to cervical surgery [5, 22–25].

Similar reports are given after orthopaedic and general surgery [26–29]. Patil et al. performed a multivariate mult-center analysis based on diagnosis codes and reported that spine surgery had the highest rates of POVL, with 0.28% for scoliosis correction and 0.14% for posterior lumbar fusion. Paediatric patients (<18 years) and elderly patients (>84 years) were 5.8 times and 3.2 times, respectively, more likely than patients 18–44 years of age to develop non-ION, non-CRAO visual loss after spine surgery. Patients with peripheral vascular disease (OR = 2.0), hypertension (OR = 3), and those who received blood transfusions (OR = 2.2) were more likely to develop non-ION, non-CRAO vision loss after spine surgery. ION was present in 0.006% of patients. Hypotension (OR = 10.1), peripheral vascular disease (OR = 6.3) and anaemia (OR = 5.9) were the strongest risk factors identified for the development of ION [30]. An overview of the published studies of POVL after spine surgery is shown in Table 1.

Case

Special history

A 73-year-old patient presented with radicular pain and sensorimotor deficits was diagnosed with a large

mediolateral lumbar disc herniation at lumbar level 4/5 and a caudally displaced sequester that compressed the right-sided root of the nerve L5. The patient was treated conservatively as an in-patient with physiotherapy and analgesics for over 6 weeks. Lack of improvement led to indication for discectomy and the patient was transferred to our department for surgical treatment.

General history

The patient presented with the following comorbidities: arterial hypertension, generalised arteriosclerosis, type 2 diabetes mellitus, obesity (BMI 31 kg/m²) and radiation therapy for prostate gland cancer 3 years ago. His medication was NORVASC® 10 mg (Amlodipine, Besylate), ATACAND PLUS® 16 mg (combination of Candesartan and Hydrochlorothiazide), BELOC-ZOK® 50 mg (Metoprolol), MODURETIC® (Hydrochlorothiazide 50 mg and Amilorid HCl 5 mg), ASPIRIN CARDIO® 100 mg (Acetylsalicylic acid), AMARYL® 2 mg (Glimerid). He had no history of angina pectoris or other cardiac symptoms and the patient's general physical condition was appropriate for his age.

The day before surgery, the patient underwent preoperative anaesthetic evaluation. Blood pressure was 130/80 mmHg and haemoglobin level was 143 g/l with a haematocrit of 0.38. The indication for surgery was verified by the responsible surgeon and affirmed with written consent, including information about the risks of a POVL. The only drugs given on the day of surgery were oral midazolam 7.5 mg, metoprolol and candesartan.

Surgery

Standard monitoring was established in the OR consisting of non-invasive blood pressure monitoring, ECG and

Table 1 Overview of the published studies of POVL after spine surgery [15, 56, 34, 4, 13, 57, 38, 36, 6]

Study	Number of POVL-patients	Incidence of POVL	Comments
Retrospective studies			
Patil et al. [15] 2008	4,452	0.094% (4,452/4,728,815)	1993–2002 US population analysis, ICD based
Chang and Miller [56] 2005	4	0.03% (4/14,102)	John Hopkins
Warner et al. [34] 2001	8	0.07% (8/11,492)	Mayo clinic
Roth and Barach [4] 2001	1	0.09% (1/1,100)	University of Chicago
Stevens et al. [13] 1997	7	0.20% (7/3,450)	Pooled data from 3 centres
Survey studies			
Delattre et al. [57] 2007	17	–	28 French orthopaedic spine centres
Lee et al. [38] 2006	93	–	ASA POVL-registry
Cheng et al. [36] 2000	24	–	AANS survey
Myers et al. [6] 1997	37	–	Scoliosis Research Society (60/400 responses)

arterial oxygen saturation (SaO₂). Blood pressure was 130/85 mmHg and heart rate 85 s⁻¹. Anaesthesia was induced with intravenous thiopental (4 mg/kg) and fentanyl (1 µg/kg). Succinylcholine (1 mg/kg) was used for neuromuscular relaxation. Anaesthesia was maintained with sevoflurane (0.5–1.5 MAC). Following endotracheal intubation, 10 min later, the patient developed hypotension of 80/45 mmHg that was treated with bolus doses of phenylephrine (100 µg) and ephedrine (10 mg). There was no change in heart rate. Blood pressure then stabilised at 100/50 mmHg. After prone positioning, occulo-bulbar pressure was carefully avoided. Shortly thereafter blood pressure dropped again to 75/40 mmHg. Despite bolus doses of phenylephrine and ephedrine, blood pressure remained at 85/40 mmHg. Therefore, bolus doses of epinephrine (10 µg) were repetitively injected during the following 2 h. The head of the patient was lifted and repositioned every 15 min to avoid prolonged pressure on the ocular bulbs. After flavectomy and during dissection of the sequester in the recessus beside the medialised nerve L5, hypotension recurred and the patient was put into a Trendelenburg position to improve cardiac filling. Surgery was interrupted for approximately 10 min and then the patient was again positioned horizontally. Total blood loss was less than 100 ml. Intravenous fluids provided were 3,100 ml of Ringer's lactate and 0.9% NaCl. Postoperatively, the patient was positioned supine and extubated. The patient was immediately conscious and showed no neurological abnormality except for the pre-existing sensorimotor deficits of nerve L5 on the right side.

Postoperative sequel

Three hours after surgery the patient complained to a nurse of blurred vision in both eyes, more pronounced on his right side. Blood pressure was 130/85 mmHg. No physician was informed. Then the patient was transferred to the neurosurgical ward for further treatment. There he reported once more a reduction of vision in both eyes, worse on his right side. Nurses misinterpreted these symptoms as an effect of the eye ointment as no external problem was observable. Six hours after surgery, the nurses informed the surgeon who diagnosed a complete vision loss of the right eye and a reduction of vision of the left eye. Vital parameters were in a normal range and blood pressure was 140/80 mmHg. The right eye showed no visual reaction to any direct light stimulus though the left eye reacted promptly. Pupils were both of a middle diameter. Vision in the left eye was about 60%. The ophthalmologist diagnosed a complete occlusion of central ophthalmic artery on the right side and a partial occlusion on the left side. An antithrombotic therapy with low-molecular-weight heparin was judged superior to a lytic therapy.

Further sequel and final outcome

An ultrasound examination of the carotid artery showed no stenosis or existing plaques in the carotid bifurcation on either side. There was no evidence of any embolic event. A thoracic CT scan showed no sign of pulmonary embolism. Laboratory values 1 day after surgery for haemoglobin and haematocrit were 117 g/l and 0.32 l/l, respectively. Reduced vision in the left eye was probably pre-existing and improved with corrective lenses. The complete amaurosis of the right eye remained unchanged. In the follow-up period of 3 years with neurological and ophthalmologic examinations at 2, 6 months, 1, 2 and 3 years after surgery, the ophthalmologic findings were unchanged. The neurological examination showed that the preoperative existing deficits of nerve L5 had completely regressed.

The complete patient's history was carefully reinvestigated after the incident. It became apparent that the anti-hypertensive medication had been modified in the peripheral hospital, just before the patient's transfer into our clinic. The patient's initial medication consisted of hydrochlorothiazide 50 mg and amilorid 5 mg, captopril 12.5 mg and atenolol 100 mg combined with chlorthalidone 25 mg. Two days before surgery, the patient's antihypertensive treatment had been supplemented with a calcium antagonist (amlodipine), a beta-blocker (metoprolol) and the ACE inhibitor had been replaced by an AT-II antagonist (candesartan).

Discussion

The combination of extensive antihypertensive medication, arteriosclerosis, intraoperative hypotension and catecholamine treatment [31] is the most probable cause of POVL in this patient. Supplementation of antihypertensive medication 2 days before surgery was certainly an aggravating factor. During this 2-day period, the blood pressure might have been in a normal range for a 40-year-old man but not for a 73-year-old who is used to high blood pressure in daily life. The 3 h delay prior to visual loss is not possible to enlighten definitely, most probable causes are the missing documentation and/or the lack of awareness of the responsible nurses. Other authors report release of hallucinations and visual loss as first manifestations of postoperative unilateral blindness [32]. A greater awareness of the patient's first reported visual impairment by the nurses may have improved his chances of a better outcome [33].

Several groups have undertaken previous retrospective investigations of POVL: an ex post facto study was performed by the Mayo Clinic, Rochester, from 1986 to 1998 [34]. Vision loss was considered present in this investigation if any part of the visual field was affected. Initial database screening found 405 cases of new-onset vision

loss and visual changes in 410 cases; 216 of these patients regained full visual acuity within 30 days. In this study, only 1 per 125,234 patients undergoing non-cardiac surgery developed vision loss persisting for longer than 30 days. The researchers concluded that vision loss and blindness after surgery and anaesthesia are still very rare events.

The same group reported other postoperative visual changes such as blurred vision, which was reversible in most cases [33, 35]. In another extended survey, a questionnaire focussing on intraoperative factors that may predispose patients to perioperative vision loss was sent to current members of the American Association of Neurological Surgeons/Congress of Neurological Surgeons, Section on Disorders of the Spine and Peripheral Nerves [36]. Out of 290 returned surveys, 22 surgeons reported 24 patients with visual loss after spine surgery. Although many of these patients had probable causative factors for POVL (e.g. hypotension, low haematocrit level, coexisting disease), some did not ($n = 8$) [3, 4, 36, 37], suggesting the necessity of high vigilance for evolving POVL even in the absence of risk factors.

In 2006, the “American Society of Anesthesiologists” POVL-Registry did an analysis of spine surgery cases with POVL showing ION in 83 out of 93 spine surgery cases [38]. The mean age of the patients was 50 ± 14 years, and most patients were relatively healthy. Mayfield pins supported the head during surgery in 16 out of 83 cases. The mean anaesthetic duration was 9.8 ± 3.1 h and the median estimated blood loss was 2.0 l (range 0.1–25 l). Bilateral disease was present in 55 patients, with complete visual loss in the affected eye(s) in 47 cases. Patients with ION had significantly longer anaesthetic duration, more blood loss, were more frequently positioned with Mayfield pins and had more bilateral POVL compared to the remaining 10 cases with CRAO ($P < 0.05$). This suggests a different aetiology for ION and CRAO. The conclusion was that ION was the most common cause of POVL after spine surgeries reported while most of the affected patients in this study were relatively healthy. Blood loss of over 1,000 ml or anaesthetic duration of over 6 h was reported in 96% of these cases. The authors recommended that for patients undergoing prolonged spine surgery in the prone position, the risk of POVL should be discussed preoperatively with the patients.

Intraoperative visual-evoked potential monitoring has been proposed [30], but has not yet proven to be effective.

The aetiology of POVL is probably multifactorial [39]. However, patients with an extensive blood loss resulting in hypotension and anaemia along with prolonged operative durations [29] may be more at risk to develop visual disturbances. An acute anaemic state may have an additive or

synergistic effect with other factors such as medical comorbidities leading to visual disturbances [11, 40].

Different hypotheses for the pathophysiological causes of POVL were discussed and investigated. Even without surgery, ION is a common cause of visual loss in the elderly population and shows certain risk factors like hypertension, nocturnal hypotension, diabetes mellitus, atherosclerosis and small cup in the optic disc [41, 42], which should be taken into account in spinal surgery patients. In addition to patient positioning during surgery [43–46], surgery duration and loss of blood [47] seem to be important risk factors [48]. Other factors such as micro-embolisms [49], decrease of haematocrit [47] or poor oxygen saturation [50], hypocapnia [50] and prolonged external ocular compression during surgery are also described [51]. These factors may occur as a single or multifactorial cause [39, 52, 53].

Our hypothesis is that in the reported patient, POVL was caused by an unusually low intraoperative blood pressure where the perfusion pressure was too low for the arteriosclerotically altered vessels and effector organs. Missing information regarding the very recent alteration of anti-hypertensive medication was an additional factor.

Conclusion

In patients with cardiovascular comorbidities such as arterial hypertension, decreased blood circulation and microcirculation, severe postoperative complications such as POVL should be taken into account while planning surgical treatment [38]. Adequate transfusion guidelines should be established for at-risk patients. Unacceptable values of haemoglobin and haematocrit should be corrected preoperatively (e.g. by substituting iron and erythropoietin) and monitored during surgery to avoid intraoperative anaemia in at-risk patients. The blood pressure of patients with predisposing diseases should be kept within normal limits. To avoid POVL as a devastating complication, it is imperative that anaesthesiologists understand contributing factors and prevention strategies [54, 55].

It is recommended for spine surgeons to obtain preoperative informed consent regarding POVL. Since visual loss following spine surgery may be reversible in the early stages, awareness, evaluation and prompt management of this rare but potentially devastating complication is critical [15]. Therefore, we chose the following recommendations from the literature to be regarded by surgeons and anaesthesiologists.

Recommendations according to the literature [1, 7, 15]:

- Avoid direct pressure on the globes
- Avoid perioperative hypotension

- Avoid perioperative anaemia
- Consider 10 degrees of reverse Trendelenburg during prone surgery
- Lower transfusion threshold to keep haematocrit above 30% in at-risk patients
- Avoid infusions of large amounts of crystalloid
- Consider staging long spinal surgeries (above 8 h)
- Maintain mean arterial pressure at patient's baseline
- Avoid changes in any perfusion-related medication shortly before surgery
- Perform a postoperative visual exam as early as possible in at-risk patients

At-risk patients are defined as patients who suffer from a pre-existing cardiovascular disease and additional metabolic diseases, where a prolonged duration of surgery in prone position (e.g. >2 h) or an increased blood loss is expected.

Acknowledgments The authors would like to thank the staff of the Department of Ophthalmology as well as Petra Schwenzer and Virginia Dittrich for their support in editing. The manuscript submitted does not contain information about medical device(s)/drug(s). No funds were received in support of this work. No benefits in any form have been or will be received from a commercial party related directly or indirectly to the subject of this manuscript.

References

1. Baig MN, Lubow M, Immesoete P, Bergese SD, Hamdy EA, Mendel E (2007) Vision loss after spine surgery: review of the literature and recommendations. *Neurosurg Focus* 23:E15
2. Petrozza PH (2002) Major spine surgery. *Anesthesiol Clin North Am* 20:405–415
3. Cheng MA, Tempelhoff R (2002) Postoperative visual loss, still no answers yet. *Anesthesiology* 96:1531–1532
4. Roth S, Barach P (2001) Postoperative visual loss: still no answers—yet. *Anesthesiology* 95:575–577
5. Kamming D, Clarke S (2005) Postoperative visual loss following prone spinal surgery. *Br J Anaesth* 95:257–260
6. Myers MA, Hamilton SR, Bogosian AJ, Smith CH, Wagner TA (1997) Visual loss as a complication of spine surgery. A review of 37 cases. *Spine* 22:1325–1329
7. Anesthesiologists Aso (2006) Practice advisory for perioperative visual loss associated with spine surgery: a report by the American society of anesthesiologists task force on perioperative blindness. *Anesthesiology* 104:1319–1328
8. Newman NJ (2008) Perioperative visual loss after nonocular surgeries. *Am J Ophthalmol* 145:604–610
9. Kawaguchi M, Hayashi H, Kurita N, Furuya H (2009) Postoperative visual disturbances after non-ophthalmic surgery. *Masui* 58:952–961
10. Shen Y, Drum M, Roth S (2009) The prevalence of perioperative visual loss in the United States: a 10-year study from 1996 to 2005 of spinal, orthopedic, cardiac, and general surgery. *Anesth Analg* 109:1534–1545. doi:[ane0b013e3181b0500b](https://doi.org/10.1097/BRS.0b013e3181b0500b)[pii]101213/ane0b013e3181b0500b
11. Williams EL (2002) Postoperative blindness. *Anesthesiol Clin North Am* 20:605–622
12. Stambough JL, Dolan D, Werner R, Godfrey E (2007) Ophthalmologic complications associated with prone positioning in spine surgery. *J Am Acad Orthop Surg* 15:156–165. doi:[15/3/156](https://doi.org/10.1097/BRS.0b013e3181b0500b)[pii]
13. Stevens WR, Glazer PA, Kelley SD, Lietman TM, Bradford DS (1997) Ophthalmic complications after spinal surgery. *Spine* 22:1319–1324
14. Roth S (2009) Perioperative visual loss: what do we know, what can we do? *Br J Anaesth* 103suppl 1:31–40. doi:[aep295](https://doi.org/10.1093/bja/aep295)[pii]101093/bja/aep295
15. Patil CG, Lad EM, Lad SP, Ho C, Boakye M (2008) Visual loss after spine surgery: a population-based study. *Spine Phila Pa* 1976 33:1491–1496. doi:[10.1097/BRS.0b013e318175d1bf00007632-200806010-00016](https://doi.org/10.1097/BRS.0b013e318175d1bf00007632-200806010-00016)[pii]
16. Helaine L, Cadic A, Magro E, Simon A, Kiss G, Gueret G, Arvieux CC (2009) Vision loss after spine surgery: a case report. *Ann Fr Anesth Reanim*
17. Katz DA, Karlin LI (2005) Visual field defect after posterior spine fusion. *Spine Phila Pa* 1976 30:83–85. doi:[00007632-200502010-00025](https://doi.org/10.1097/BRS.0b013e318175d1bf00007632-200502010-00025)[pii]
18. Kumar N, Jivan S, Topping N, Morrell AJ (2004) Blindness and rectus muscle damage following spinal surgery. *Am J Ophthalmol* 138:889–891
19. Katzman SS, Moschonas CG, Dzioba RB (1994) Amaurosis secondary to massive blood loss after lumbar spine surgery. *Spine* 19:468–469
20. Kothari MT, Maiti A (2007) Ophthalmic artery occlusion: a cause of unilateral visual loss following spine surgery. *Indian J Ophthalmol* 55:401–402
21. Lee LA, Lam AM (2001) Unilateral blindness after prone lumbar spine surgery. *Anesthesiology* 95:793–795
22. Hoski JJ, Eismont FJ, Green BA (1993) Blindness as a complication of intraoperative positioning A case report. *J Bone Jt Surg Am* 75:1231–1232
23. Kasodekar VB, Chen JL (2006) Monocular blindness: a complication of intraoperative positioning in posterior cervical spine surgery. *Singapore Med J* 47:631–633
24. Lubnin A, Serova NK, Proshutinskii SD, Konovalov NA, Moshchev DA (2008) Sudden loss of vision in one eye in a female patient after neurosurgical intervention into the cervical portion of the spinal cord. *Anesteziol Reanimatol*, pp 88–91
25. Manfredini M, Ferrante R, Gildone A, Massari L (2000) Unilateral blindness as a complication of intraoperative positioning for cervical spinal surgery. *J Spinal Disord* 13:271–272
26. Fournier JH, Velis E (2004) Visual loss after shoulder surgery under general anesthesia diagnosed as caused by ocular compression with electroretinography testing: case report and review of the literature. *Int Surg* 89:209–211
27. Gaillard MC, Zambaz BD, Borruat FX (2004) Posterior ischemic optic neuropathy: case report of a rare complication after general surgery. *Klin Monatsbl Augenheilkd* 221:421–423
28. Pandey V, Rao PS, Rao SK, Acharya KK (2008) Monocular blindness due to central retinal artery occlusion in bipolar hemireplacement arthroplasty of the hip. *Singapore Med J* 49:e96–e97
29. Grossman W, Ward WT (1993) Central retinal artery occlusion after scoliosis surgery with a horseshoe headrest. Case report and literature review. *Spine* 18:1226–1228
30. Kabbara AI (2007) What happened to the old visual evoked potential monitoring? *Anesthesiology* 106:1249; author reply 1251–1242
31. Lee LA, Nathens AB, Sires BS, McMurray MK, Lam AM (2005) Blindness in the intensive care unit: possible role for vasopressors? *Anesth Analg* 100:192–195
32. Montero JA, Ruiz-Moreno JM, Galindo A, Fernandez-Munoz M (2007) Release hallucinations and visual loss as first manifestations of postoperative unilateral blindness. *Eur J Ophthalmol* 17:844–846
33. Yi HJ, Kim DW (2004) Reversible unilateral blindness after lumbar spine surgery: a role for cerebrospinal fluid leakage? *J Neurosurg Anesthesiol* 16:322–323

34. Warner ME, Warner MA, Garrity JA, MacKenzie RA, Warner DO (2001) The frequency of perioperative vision loss. *Anesth Analg* 93:1417–1421
35. Warner ME, Fronapfel PJ, Hebl JR, Herman DC, Warner DO, Decker P, Warner MA (2002) Perioperative visual changes. *Anesthesiology* 96:855–859
36. Cheng MA, Sigurdson W, Tempelhoff R, Laurysen C (2000) Visual loss after spine surgery: a survey. *Neurosurgery* 46:625–630 discussion 630–621
37. Roth S, Tung A, Ksiazek S (2007) Visual loss in a prone-positioned spine surgery patient with the head on a foam headrest and goggles covering the eyes: an old complication with a new mechanism. *Anesth Analg* 104:1185–1187
38. Lee LA, Roth S, Posner KL, Cheney FW, Caplan RA, Newman NJ, Domino KB (2006) The American society of anesthesiologists postoperative visual loss registry: analysis of 93 spine surgery cases with postoperative visual loss. *Anesthesiology* 105:652–659 quiz 867–658
39. Benumof JL, Mazzei W, Roth S, Barach P, Lam AM, Lee LA (2002) Multifactorial etiology of postoperative vision loss. *Anesthesiology* 96:1531–1532 (author reply 1532–1533)
40. Gill B, Heavner JE (2006) Postoperative visual loss associated with spine surgery. *Eur Spine J* 15:479–484
41. Obuchowska I, Mariak Z (2006) Ischemic optic neuropathy. Pathogenesis, clinical features, diagnostics and treatment. *Klin Oczna* 108:238–242
42. Luneau K, Newman NJ, Biousse V (2008) Ischemic optic neuropathies. *Neurologist* 14:341–354
43. Bermejo-Alvarez MA, Carpintero M, Garcia-Carro G, Acebal G, Fervienza P, Cosio F (2007) Ischemic optic neuropathy after lumbar spine surgery. *Rev Esp Anestesiol Reanim* 54:621–625
44. Bekar A, Tureyen K, Aksoy K (1996) Unilateral blindness due to patient positioning during cervical syringomyelia surgery: unilateral blindness after prone position. *J Neurosurg Anesthesiol* 8:227–229
45. Chung MS, Son JH (2006) Visual loss in one eye after spinal surgery. *Korean J Ophthalmol* 20:139–142
46. Heitz JW, Audu PB (2008) Asymmetric postoperative visual loss after spine surgery in the lateral decubitus position. *Br J Anaesth* 101:380–382
47. Dunker S, Hsu HY, Sebag J, Sadun AA (2002) Perioperative risk factors for posterior ischemic optic neuropathy. *J Am Coll Surg* 194:705–710
48. Torossian A, Schmidt J, Schaffartzik W, Wulf H (2006) Loss of vision after non-ophthalmic surgery: systematic review of the literature on incidence, pathogenesis, treatment and prevention. *Anaesthesist* 55:457–464
49. Nakra D, Bala I, Pratap M (2007) Unilateral postoperative visual loss due to central retinal artery occlusion following cervical spine surgery in prone position. *Paediatr Anaesth* 17:805–808
50. Lee LA, Vavilala MS, Lam AM, Douville C, Moore A, Visco E, Newell DW (2004) Ophthalmic artery blood flow velocity increases during hypocapnia. *Can J Anaesth* 51:388–392
51. Tachfouti S, Karmane A, El Moussaif H, Boutimzine N, Daoudi R (2007) Orbital ischemia syndrome after surgical intervention of the spine. A case report. *Bull Soc Belge Ophthalmol*, pp 27–30
52. Biousse V, Newman NJ (2007) Vision loss: overview. *Semin Neurol* 27:199–210
53. Ho VT, Newman NJ, Song S, Ksiazek S, Roth S (2005) Ischemic optic neuropathy following spine surgery. *J Neurosurg Anesthesiol* 17:38–44. doi:0008506-200501000-00009[pii]
54. Rupp-Montpetit K, Moody ML (2004) Visual loss as a complication of nonophthalmologic surgery: a review of the literature. *AANA J* 72:285–292
55. Rupp-Montpetit K, Moody ML (2005) Visual loss as a complication of non-ophthalmic surgery: a review of the literature. *Insight* 30:10–17
56. Chang SH, Miller NR (2005) The incidence of vision loss due to perioperative ischemic optic neuropathy associated with spine surgery: the Johns Hopkins Hospital Experience. *Spine Phila Pa* 1976 30:1299–302. doi:00007632-200506010-00013[pii]
57. Delattre O, Thoreux P, Liverneaux P, Merle H, Court C, Götting M, Rouvillain JL, Catonné Y (2007) Spinal surgery and ophthalmic complications: a French survey with review of 17 cases. *J Spinal Disord Tech* 20:302–307. doi:10.1097/01.bsd.0000211290.21766.04; 00024720-200706000-00008[pii]