

## Development of microsatellite loci in the European Dipper, *Cinclus cinclus*

T. B. Bucher · P. Wandeler · J. Hegelbach ·  
L. F. Keller

Received: 9 July 2009 / Accepted: 31 July 2009 / Published online: 26 August 2009  
© Springer Science+Business Media B.V. 2009

**Abstract** Eighteen polymorphic microsatellite DNA loci were isolated in the Central European subspecies of the European Dipper (*Cinclus cinclus aquaticus*). The loci were tested for polymorphism using a test panel of 24 breeding birds. Numbers of alleles ranged from 2 to 21 per locus and expected heterozygosities varied between 0.47 and 0.83. Two loci (Cici10 and Cici12) proved to be Z-linked. Some pairs of loci exhibited significant linkage disequilibrium but not the two loci that are located on the Z-chromosome. This pattern suggests that demographic effects rather than physical linkage are likely responsible for the observed levels of linkage disequilibrium. These loci will be useful for applied conservation projects and for investigations of the dispersal and mating patterns of European and other dippers.

**Keywords** Dipper · Cinclidae · *Cinclus mexicanus* · Multiplex PCR · Z-linked loci

The only passerine birds able to dive in fast-flowing rivers are the five species of dippers (Cinclidae; Aves). Not surprisingly, perhaps, given the adaptations that are required to inhabit such a habitat, this genus with its five species is monophyletic (Voelker 2002). As extreme habitat specialists, dippers are sensitive to environmental change, e.g. to changes in water quality (Tyler and Ormerod 1994) or to climate change (Hegelbach 2001; Moreno-Rueda and Rivas 2007). Consequently, some populations in Europe

(Henderson et al. 2004) and North America (Anderson et al. 2008) have been declining substantially and are, therefore, of conservation concern.

The palaeartic representative, the European or White-throated Dipper *Cinclus cinclus*, has the widest geographical distribution. Its populations are distributed over Europe, Asia and a small strip of Northern Africa and several subspecies have been described based on morphology (Tyler and Ormerod 1994). The subspecies status of Central Europe's *Cinclus cinclus aquaticus* has been confirmed recently using *cytochrome b* sequence variation (Lauga et al. 2005).

Here, we describe the isolation and characterization of 18 variable di- and tetranucleotide microsatellite loci from a population of *Cinclus cinclus aquaticus* near Zurich, Switzerland. Blood samples were preserved in APS-buffer pH 8.0 (Sibley and Ahlquist 1981), containing 10% EDTA, 1% sodium fluoride and a small amount of thymol sufficient to ensure saturation. DNA was extracted using the QIAmp DNA mini kit (QIAGEN).

An enriched DNA library was made by ECOGENICS GmbH (Zürich, Switzerland) from size-selected genomic DNA ligated into TSPAD-linker (Tenzer et al. 1999) and enriched by magnetic bead selection with biotin-labelled (CA)<sub>13</sub> and (ACAG)<sub>7</sub> oligonucleotide repeats (Gautschi et al. 2000a, b). Out of the 864 recombinant colonies screened, 310 gave a positive signal after screening with fluorescently labelled CA or ACAG repeats. Plasmids from 140 positive clones were sequenced as described in Gautschi et al. (2000a) and primers were designed for 29 CA and six ACAG microsatellite inserts. Of these, 17 and 6, respectively turned out to be polymorphic in a panel of 24 unrelated individuals screened for polymorphism on ELCHROM gels (Table 1). Five loci were not developed further.

T. B. Bucher (✉) · P. Wandeler · J. Hegelbach · L. F. Keller  
Zoological Museum, University of Zurich,  
Winterthurerstrasse 190, 8057 Zurich, Switzerland  
e-mail: t.bucher@access.uzh.ch

**Table 1** Characteristics and summary statistics of 20 polymorphic microsatellite loci isolated from a *Cinclus cinclus aquaticus* population in Switzerland and genotyped in all breeding birds born in 1999 in one river system (Küsnacht)

Locus name/GenBank accession no.	Repeat motif	Primer sequence (5'-3') and fluorescent label	Multiplex panel no.	Primer concentration ( $\mu$ M)	$T_a$ ( $^{\circ}$ C)	Size range (bp)	$N$	$N_a$	$H_o$	$H_e$	HW (exact)
Cici 01 FJ177034	(CA)21	GTGGAGCAGGAATCTAAACTGG FAM-CCTGCCCAGAGTTTTTAATGC	1	2.00	63	100–150	24	16	0.2917	0.7839	<b>0.0001</b>
Cici 02 FJ177035	(GT)17	AGCATCACAGGAGCAAAAGTG VIC-CTACGGCTACATCTCACTTCC	1	1.00	63	90–140	24	10	0.5417	0.6658	0.1987
Cici 03 FJ177036	(CA)20 AA (CA)4	GGGGCTCATGATGGATAG VIC-CTGTGAAGTTATCAGAGGATGTTTG	2	2.00	61	140–190	23	14	0.8261	0.6947	0.5032
Cici 04 FJ177037	(GT)22	GGATGATCTGGGCTATGTGC FAM-CAGCTCTGGTGGTCAGGTG	1	1.00	63	155–210	24	14	1	0.8229	0.2669
Cici 05 FJ177038	(CA)21	FAM-CTCCGGTCACTCCCATACC GGTGTGGTGTGATGTTGCAG	1	6.00	63	220–270	24	16	0.875	0.7752	0.5196
Cici 06 FJ177039	(TGA)3 TGGG (TGA)12	PET-CAGCAAAAAGCATACCCCTGACT GCTTTATTACCCCTGCTGGAT	1	1.00	63	95–150	24	10	0.875	0.7752	0.0788
Cici 07 FJ177040	(GT)23	NED-TAAGTCCCGAGCACCATCTG TTTTCGTTCTCCAGAGCAGT	1	2.00	63	190–250	24	13	0.8333	0.7292	0.1842
Cici 08 FJ177041	(GT)14	FAM-CCCCAAATATCCTACCACACC TTTGGGTTCAAACACAGGATGG	2	2.00	61	210–250	24	9	0.6667	0.7465	0.6924
Cici 09 FJ177042	(CTGT)6 (CT)3	GAAGGCCAGATGACATTGCT NED-CCCTTGCCCTTGCAGACTGT	1	0.35	63	100–130	24	2	0.4167	0.4688	0.6644
Cici 10 <sup>a</sup> FJ177043	(ATCT)14 (GTCT)5 (ATCT)2 (ATCC)11	VIC-TGTGGATAAAGGATCTCAATGC CCCAGAAAAGGTACAGGATGG	1	1.00	63	145–202	24	10	0.2917	0.651	<b>&lt;0.0001</b>
Cici 11 FJ177044	(GT)23	PET-GAAGCGCAAGGATTTTCAC GCTGGACTCCTGTGTTTGACT	2	2.00	61	150–260	24	21	0.8333	0.7344	0.7129
Cici 12 <sup>a</sup> FJ177045	(TCAT)17 TATGGA (TCAT)4	VIC-CAAACCTGCACGTGGAATG GTCTGGCCCCACAGAGTTAC	1	0.50	63	205–280	24	15	0.375	0.6892	<b>0.0001</b>
Cici 13 FJ177046	(GTCT)11 GTACCCC (CA)5	NED-GTGTCCAGTCAGTCAAAAAGC TAGAGATTTCATGAGCACCTTGG	1	0.70	63	140–185	24	7	0.6667	0.5799	1.0000
Cici 14 FJ177047	(GTCT)9	ACCGCCAAACAAACAAAAGG PET-TCTCCTGTTAGCCCCGAAG	1	2.00	63	155–230	24	14	0.9583	0.7873	0.4615
Cici 16 FJ177049	(CA)12	FAM-TTAGTGGGGCTTCTGAGCAC CAGGCAAGTTCAGCCACAG	2	2.00	61	253–300	24	7	0.4167	0.48	0.1602
Cici 17 FJ177050	(CA)20	ACTTGGGATGGGAACCACT VIC-TGGAATCTACTTATGGCTGCAT	2	2.50	61	211–310	24	14	0.7083	0.6623	0.2907

**Table 1** continued

Locus name/GenBank accession no.	Repeat motif	Primer sequence (5'–3') and fluorescent label	Multiplex panel no.	Primer concentration (μM)	T <sub>a</sub> (°C)	Size range (bp)	N	N <sub>a</sub>	H <sub>O</sub>	H <sub>E</sub>	HW (exact)
Cici 22 FJ177055	(CA) <sub>19</sub>	NED-GCTGGACCTCTTGGGTACT AGCAGCACCAGTGGAAAGAGT	2	8.00	61	260–330	24	16	0.6667	0.7118	0.6577
Cici 23 FJ177056	(CA) <sub>16</sub>	FAM-ACCGTTTGAGGTGCCAGA AATGGCTGTGGTCTTGTTCA	2	2.00	61	108–160	24	15	0.9167	0.8325	0.4559

One primer of each pair was 5' fluorescently labelled with 6-FAM, NED, PET or VIC (ABI)

T<sub>a</sub> annealing temperature of primer pairs, N number of individuals genotyped at each locus, N<sub>a</sub> observed number of alleles, H<sub>O</sub> observed heterozygosity, H<sub>E</sub> Nei's unbiased expected heterozygosity; HW (exact), exact P values of a test for deviations from Hardy–Weinberg equilibrium at each locus, with tests significant after Bonferroni correction in bold type  
<sup>a</sup> Z-linked

The remaining 18 polymorphic microsatellites were split into two panels for multiplex PCR. Panel one included eleven and panel two seven *Cinclus* primer pairs. In addition, two loci designed for other bird species were also included in panel two: QmAAT31 (Hughes et al. 1998) and Ase64 (Richardson et al. 2000). Amplification was conducted in a final volume of 5 μl containing 2 μl of PCR multiplex kit (QIAGEN), between 0.1 and 2.2 μM of each primer, one of which was labeled with a fluorescent dye (Table 1), and 20–40 ng of template DNA, using the following cycling parameters: 15 min of denaturation at 95°C, followed by 28 cycles of 30 s at 94°C, 90 s at 61°C (panel 2) or 63°C (panel 1) and 60 s extension at 72°C and a final extension of 30 min at 60°C (GeneAmp® PCR System 9700, ABI). Fragment analyses were performed on an ABI Prism® 3100 Avant DNA analyzer and allele sizes scored against the size standard GS-500 LIZ™ using GeneMapper v3.7™ software (ABI).

All 18 loci were tested for polymorphism, deviations from Hardy–Weinberg equilibrium (HWE) and linkage disequilibrium (LD) using SAS/Genetics (SAS Institute Inc., version 9.1.3) and a test panel of 24 breeding birds that were all born in the same year and the same river. The loci showed between 2 and 21 alleles and expected heterozygosities ranged from 0.47 to 0.83 (Table 1). Significant deviations from HWE were detected at three loci (Cici01, Cici10, and Cici12). Sex-specific analyses revealed that two of these loci (Cici10 and Cici12) were Z-linked, i.e. only males were heterozygous. The heterozygote deficit at Cici01, however, seemed to be the result of null alleles.

Tests for linkage disequilibrium revealed that Cici11 and Cici17 were in significant LD after sequential Bonferroni correction (corrected P = 0.038). These loci could in principle be physically linked. However, these two loci did not exhibit significant LD in other cohorts (data not shown) and the two Z-linked loci that were known to be on the same chromosome did not exhibit significant LD. This pattern suggests that the levels of LD detected in our test panel were more likely caused by demographic factors such as bottlenecks and immigration rather than by physical linkage.

The high levels of genetic variation (number of alleles and heterozygosities) present at these 18 microsatellite loci render them ideal for the analysis of the dipper mating system. Moreover, they may also prove useful for the analysis of dispersal and population structure. Finally, ten loci (Cici02, Cici04, Cici05, Cici08, Cici10, Cici11, Cici12, Cici13, Cici15, and Cici16) also proved polymorphic in the American Dipper (*Cinclus mexicanus*; Anderson et al. 2008) suggesting that our loci will also work in other dipper species.

**Acknowledgments** We thank Barbara Hefti-Gautschi and Ecogenics GmbH (Zurich, Switzerland) for developing the library and a subset of the markers, and María José Ruiz-López for her help with the genotyping and the initial data analysis. Funding was provided by the “Stiftung für wissenschaftliche Forschung an der Universität Zürich”.

## References

- Anderson CM, Spellman GM, Ferrell CS, Strickler K, Sarver SK (2008) Conservation genetics of American Dipper (*Cinclus mexicanus*): the genetic status of a population in severe decline. *Conserv Genet* 9:939–944
- Gautschi B, Tenzer I, Müller JP, Schmid B (2000a) Isolation and characterization of microsatellite loci in the bearded vulture (*Gypaetus barbatus*) and cross-amplification in three Old World vulture species. *Mol Ecol* 9:2193–2195
- Gautschi B, Widmer A, Koella J (2000b) Isolation and characterization of microsatellite loci in the Dice Snake (*Natrix tessellata*). *Mol Ecol* 9:2191–2193
- Hegelbach J (2001) Water temperature and phytophenology indicate the earlier onset of oviposition in Eurasian Dipper (*Cinclus cinclus*) from the Swiss Lowlands. *J Ornithol* 142:284–294
- Henderson IG, Fuller RJ, Conway GJ, Gough SJ (2004) Evidence for declines in populations of grassland-associated birds in marginal upland areas of Britain. *Bird Study* 51:12–19
- Lauga B, Cagnon C, d’Amico F, Karama S, Mouchès C (2005) Phylogeography of the white-throated dipper *Cinclus cinclus* in Europe. *J Ornithol* 146:257–262
- Moreno-Rueda G, Rivas JM (2007) Recent changes in allometric relationships among morphological traits in the dipper (*Cinclus cinclus*). *J Ornithol* 148:489–494
- SAS Institute (2003) SAS/STAT, Version 9.1.3. SAS Institute Inc, Cary, NC
- Tenzer I, degli Ivanisovich S, Morgante M, Gessler C (1999) Identification of microsatellite markers and their application to population genetics of *Venturia inaequalis*. *Phytopathology* 89:748–753
- Tyler SJ, Ormerod SJ (1994) The dippers. Poyser, London
- Voelker G (2002) Molecular phylogenetics and the historical biogeography of dippers (*Cinclus*). *Ibis* 144:577–584