

Childhood adversity and chronicity of mood disorders

Jules Angst · Alex Gamma · Wulf Rössler ·
Vladeta Ajdacic · Daniel N. Klein

Received: 3 November 2009 / Accepted: 11 June 2010 / Published online: 30 June 2010
© Springer-Verlag 2010

Abstract To evaluate the potential impact of early childhood problems on the chronicity of mood disorders. A representative cohort from the population was prospectively studied from ages 19/20 to 39/40. Unipolar (UP) and bipolar disorders (BP) were operationally defined applying broad Zurich criteria for bipolarity. Chronicity required the presence of symptoms for more days than not over 2 years prior to an interview, or almost daily occurrence for 1 year. A family history and a history of childhood problems were taken at ages 27/28 and 29/30. Data include the first of multiple self-assessments with the Symptom-Checklist-90 R at age 19/20, and mastery and self-esteem assessed 1 year later. A factor analysis of childhood problems yielded two factors: family problems and conduct problems. Sexual trauma, which did not load on either factor, and conduct problems were unrelated to chronicity of UP or BP or both together. In contrast, childhood family problems increased the risk of chronicity by a factor of 1.7. An anxious personality in childhood and low self-esteem and mastery in early adulthood were also associated with chronicity. Childhood family problems are strong risk factors for the chronicity of mood disorders (UP and BP). The risk may be mediated partly by anxious personality traits, poor coping and low self-esteem.

Keywords Chronicity · Mood disorders · Childhood adversity · Coping · Self-esteem · Personality

J. Angst (✉) · A. Gamma · W. Rössler · V. Ajdacic
Zurich University Psychiatric Hospital, Lengstrasse 31,
P.O.Box 1931, 8032 Zurich, Switzerland
e-mail: jules.angst@uzh.ch

D. N. Klein
Department of Psychology and Psychiatry and Behavioral
Sciences, Stony Brook University, Stony Brook, NY, USA

Introduction

There is growing evidence that early adversity plays an important role in the onset of mood disorders [1, 23, 24]. Numerous studies have documented associations between mood disorders and reports of parental indifference, over-control, and neglect, and physical and sexual abuse [19, 22, 31]. Although most of these data are retrospective, several prospective studies have also reported associations between childhood maltreatment and the later development of major depression [11, 38].

Our paper examines the relationship between childhood adversity and the course of mood disorders. The literature presents increasing evidence that childhood adversity impacts on the course of unipolar (UP) and bipolar (BP) mood disorders. It has been shown to correlate with an earlier onset and with higher recurrence/cycling of BP [14, 17, 25, 35] and UP [5]. Childhood adversity and maladaptive parenting were also found to be associated with chronic depression [9, 10, 28]. The association between childhood adversity and chronicity may be mediated by conduct disorder and shame-withdrawal in childhood [7], and, more importantly, by an increased vulnerability to stress in adulthood [15, 20, 39, 40]. However, in a small retrospective study comparing the relative impact of childhood adversity on the chronic course of UP and BP, Vocisano et al. [41] found that patients with chronic forms of UP and BP did not differ in terms of childhood maltreatment.

Recently, we reported that subjects with long-term/chronic as opposed to acute depression had an earlier age of onset and higher somatic and psychiatric comorbidity [4]. This paper extends our previous work by examining the link between childhood adversity and chronicity in mood disorders, and widens the scope of the earlier research by including bipolar depression.

Methodology

Zurich study

The design of the Zurich study was recently described in detail [2]. Essentially, it is a cohort study from ages 19/20 to 40/41 with six waves of interviews conducted by trained professionals. The interview sample ($N = 591$) was drawn from 4,547 subjects (2,201 males, 2,346 females) aged 19 (m) and 20 (f), who were representative of the canton of Zurich in Switzerland and screened in 1978 with the Symptom Checklist 90-R (SCL-90-R) [12]. The sample was stratified by high versus low scores on the General Severity Index of the SCL-90-R (“high risk” vs. “low risk” subjects), with oversampling of risk cases.

Over 20 years, the attrition rate was 38% (men 44%, women 31%). Only subjects participating in waves 3–6, i.e. at the ages of 27/28–40/41 (1986–1999), could be considered for the present analysis, because the temporal criteria for dysthymia had not yet been developed. The sample size is relatively small ($N = 499$) but, when sample stratification is statistically offset, it represents 2,146 subjects of the same age.

Chronic depression

There is no general agreement on the definition of long-standing, long-term, persistent or chronic depression [29]. The suggested minimum duration varies between 1 and 3 years [2]. In this study, a diagnosis of chronic major and minor depressive episodes required the temporal chronicity criterion for DSM-IV dysthymia (being symptomatic more days than not over 2 years), or mood changes had to be present almost daily over the 12 months prior to an interview; in both cases, work or social impairment had to be present.

Childhood and conduct problems

Childhood problems were assessed in retrospect at ages 27/28 and 29/30. The items are listed in Table 1.

Other variables

Unipolar depression and bipolar disorders were defined by the “broad” concept for bipolarity published earlier [3], which identified about half of all depressed subjects as bipolar. The concept is termed “broad” because it uses subthreshold definitions of hypomania and identifies more depressives as bipolar than the current DSM-IV concept. It requires a diagnosis of a major depressive episode according to DSM-IV and the presence of any number of the diagnostic symptoms of DSM-IV hypomania. No other criteria are

Table 1 Zurich study: factor analysis of childhood problems (tetrachoric correlations)

	Family problems	Conduct problems
Tense family atmosphere	0.77	−0.03
Sexual trauma	0.30	0.13
Broken home	0.61	0.14
<i>Trouble with the police</i>	−0.03	0.73
Restless in class	0.10	0.23
<i>Unpunctual</i>	0.05	0.71
Problems within family	0.81	0.21
Problems outside family	−0.09	−0.81
Father with psychol. problems	0.59	0.14
Mother with psychol. problems	0.44	0.18
Conflicts between parents	0.73	0.26
Conflicts with parents	0.61	0.24
Conflicts with other persons	0.81	−0.07
More punished than peers	0.49	0.38
Disliked, rejected by peers	0.50	0.36
<i>Frequent physical fights</i>	0.09	0.58
<i>Discipline problems at school</i>	0.09	0.78
<i>Repeated running away</i>	0.32	0.79
<i>Theft</i>	0.02	0.80
<i>Truancy</i>	0.19	0.86

Family problems are given in bold

Conduct problems are given in italics

applied. Psychiatric disorders were diagnosed by algorithms on the basis of DSM-III criteria (GAD, panic disorder), DSM-III-R criteria (major depressive disorder, phobias, obsessive–compulsive disorder (OCD)), and DSM-IV criteria (post-traumatic stress disorder and substance abuse/dependence). Mastery and self-esteem were assessed by the scales of Pearlin and Schooler [33] at age 20/21. Mastery means the extent to which individuals perceive their life-chances and/or outcomes as being under their own control rather than externally determined. Self-esteem refers to an individual’s positive or negative attitude towards the self as a whole, and is assessed by statements such as “On the whole, I am satisfied with myself” and “I take a positive attitude towards myself”. The Symptom Checklist 90 R (SCL-90-R) [13] was used for the self-assessment of depression, anxiety, somatisation and vegetative lability [36] at age 19/20. We present T-transformed scores (mean = 50, SD = 10). An anxious personality in childhood/adolescence refers to an individual’s having been more anxious than his/her peers in childhood.

Statistics

Frequencies were compared across groups using χ^2 tests. Kruskal–Wallis tests were applied to continuous variables.

Apart from sample frequencies, we also report population estimates of cumulative incidence rates for a disorder. These are obtained as follows: first, the cumulative incidence rate of a disorder within the study sample is computed. The cumulative incidence rate is the percentage of subjects who *ever* reported (i.e. in any given interview) having had the disorder within the previous 12 months. It should be noted that, on account of the oversampling of risk cases, rates for most disorders are substantially higher in our sample than in the population. It should also be noted that the cumulative incidence rate is usually lower than the lifetime prevalence rate because it is based on the cumulation of (at most) 6–1-year prevalence rates. It can be considered as a lower bound for the lifetime prevalence rate. Moreover, the cumulative incidence rates in the sample are used to calculate population estimates using standard statistical procedures to offset sample stratification (i.e. to correct the bias introduced by oversampling risk cases). The results are cumulative incidence rates representative for the cohort of 19/20-year-olds in the population of the canton of Zurich in 1978.

The 20 dichotomous items on childhood adversity (Table 1) were factor analysed using a tetrachoric correlation matrix and oblique rotation. Eigenvalues, scree plot and results of Horn's parallel test were inspected to decide on the number of factors to be retained. The empirical and theoretical validity and interpretability of the factors were also considered in this decision.

A multivariate logistic regression was conducted to examine the influence of several predictors on the chronicity of mood disorder. The initial independent variables entered into the regression model were the level of childhood family problems, the level of childhood conduct problems, the presence of psychological problems in the father, the presence of psychological problems in the mother, the self-rating score of mastery (coping) at age 20/21, the self-rating score of self-esteem at age 20/21, the scores of the SCL scales for depression, anxiety, somatisation and interpersonal sensitivity at age 19/20 (t-transformed), the presence of an anxious personality in childhood/adolescence, the presence of a family history of depression, and the presence of a family history of mania or bipolar disorder. The models were reduced by eliminating, one by one, the least significant variable. If the deletion of a variable led to a change of more than 20% in one or more of the remaining regression coefficients, the variable was kept in the model. This procedure continued until only variables with a statistical significance level of at least $P < 0.1$ remained. Only these variables will be reported.

The overall analysis involved many tests of statistical significance, raising the problem of type I errors, i.e. of

tests being statistically significant by chance alone. We offer a correction for multiple comparisons but nevertheless present the uncorrected P -values, because if only corrected values are shown, important findings may be missed. A Bonferroni-corrected threshold for statistical significance is reported with each table. The correction can be applied by simply raising the level of statistical significance to this threshold.

Analyses were performed in Stata 10.2 and SAS 8.2 for Windows.

Results

Descriptive data on the sample

We identified 104 major or minor bipolar (BP) subjects and 110 major or minor unipolar depressives (UP). The respective weighted cumulative incidence rates including the last interview (which are statistical estimates of the lower bound of population prevalence) were 15.4% (BP) and 14.8% (UP). There was no significant gender difference: 60.6% of BP (weighted 62.6%) and 60.0% of UP (weighted 61.4%) were women.

Thirty-eight BP subjects were chronic and 66 acute; among UP there were 43 chronic and 67 acute.

The 38 chronic BP subjects included 25 with MDE (67%), 14 of whom also met criteria for dysthymia. In all, 22 (58%) of the 38 chronic BP cases suffered from dysthymia. The remaining five non-dysthymic chronic cases suffered from recurrent brief depression ($N = 4$) and minor depression ($N = 1$).

The 43 chronic UP subjects consisted of 34 with MDE (79.1%), 21 of whom were also dysthymics. In all, 27 subjects (62.8%) with chronic UP were diagnosed as dysthymics. Of the remaining three non-dysthymic chronic UP cases, two suffered from recurrent brief depression and 1 from minor depression.

Factor analysis of childhood family problems

A factor analysis ($N = 397$) of the 20 items on childhood family problems resulted in two factors: family problems (F1) and conduct problems (F2; Table 1). The reliability (Cronbach's alpha) was 0.76 (F1) and 0.62 (F2). The two factors were uncorrelated ($r = 0.04$). Three items did not load strongly enough on the two factors and were excluded from the scales: "sexual trauma", "restless in class", and "problems outside the family". "Problems outside the family" loaded negatively on F2; this item was not included in the scale because its interpretation is uncertain and it does not fit with the other items representing conduct behaviour.

Childhood family and conduct problems by gender

The frequencies of childhood problems by gender are shown in Table 2. There were some differences: women reported more sexual trauma and psychological problems in their fathers and men reported more discipline problems at school and more trouble with the police.

There were no significant gender differences in family problems scores (F1) but, as expected, men scored significantly higher than women on behavioural/conduct problems (F2).

Childhood family and conduct problems in BP and UP subjects

UP and BP subjects did not differ in their total scores on the two factors: family problems (mean 3.2 ($s = 2.4$) vs. mean 3.1 ($s = 2.6$), $N = 82$ vs. 95) and conduct problems (mean 0.8 ($s = 1.4$) vs. mean 1.0 ($s = 1.4$), $N = 93$ vs. 100). At item level, being unpunctual at school was the only behaviour on which BP subjects scored higher. Sexual trauma did not differ significantly ($P < 0.21$) between the 110 UP (15.5%) versus 104 BP (22.1%) probands, but the two diagnostic groups differed significantly ($P < 0.0006$)

Table 2 Scores for childhood family and conduct problems ($M = 292$, $F = 299$)

	<i>M</i>	<i>F</i>	<i>P</i>
Tense family atmosphere	22.9	25.4	ns
Sexual trauma	4.8	18.7	0.001
Broken home	17.0	16.0	ns
<i>Trouble with the police</i>	41.4	15.4	0.001
Restless in class	27.6	21.6	ns
<i>Unpunctual</i>	8.9	4.3	0.05
Problems within family	42.2	47.8	ns
Problems outside family	0.4	1.3	ns
Father with psychol. problems	11.6	21.4	0.007
Mother with psychol. problems	17.1	23.7	ns
Conflicts between parents	25.8	29.5	ns
Conflicts with parents	28.1	24.8	ns
Conflicts with other persons	23.1	26.8	ns
More punished than peers	13.6	20.1	ns
Disliked, rejected by peers	12.4	18.1	ns
<i>Frequent physical fights</i>	8.0	6.9	ns
<i>Discipline problems at school</i>	20.4	8.6	0.001
<i>Repeated running away</i>	3.1	4.3	ns
<i>Theft</i>	14.2	8.2	0.04
<i>Truancy</i>	11.1	5.6	0.03

Family problems are given in bold

Conduct problems are given in italics

Bonferroni threshold: $P = 0.0025$

from the 183 other subjects without mood disorders (6.6%). In bipolar women, an interaction between the age of onset of hypomanic symptoms and a family history of depression was found with regard to the level of childhood family problems: in women with a family history of depression, the age of onset of their hypomanic symptoms rose from about 10 to 17 years with increasing levels of childhood family problems, but *decreased* from about 17 to 13 years in women without such a family history.

Childhood family and conduct problems in subjects with chronic versus acute BP and UP disorders

Table 3 presents the scores and significant differences between chronic and acute subgroups of mood disorders. Sufferers from chronic mood disorders scored consistently higher on childhood family problems; the differences were significant in the UP and in the combined (UP + BP) groups.

Interestingly, there were no associations between chronicity and the other variables entered into the analyses: sexual trauma, gender, SCL-90-R depression, anxiety and interpersonal sensitivity scores, family history of mania, and anxious personality (Table 3). Remarkably also childhood conduct problems were not associated significantly with chronicity (Table 4).

Multivariable logistic regression analyses on chronicity

Three multivariable logistic regression analyses were carried out comparing subjects with and without chronic UP, BP, and UP + BP.

Table 5 summarises the results. The score for childhood family problems was the most consistent predictor of chronicity across all three diagnostic groups with odds ratios between 1.7 and 1.8. Being more anxious than one's peers was a risk factor for UP (OR = 4.0) and for UP + BP (OR = 2.3). Unexpectedly, a family history of depression was negatively associated with chronic BP at a trend level (OR = 0.4) and with UP + BP (OR = 0.5). Furthermore, lower self-esteem was associated with chronic BP, as was decreased mastery in the merged group (UP + BP). Finally, somatisation, as measured by the SCL-90-R at age 19/20, was positively associated with chronic UP (OR = 2.1).

Discussion

Our data are restricted to a relatively small but representative age cohort interviewed four times from the ages of 27/28 to 39/40. The relatively small Ns may have created some type II errors. We must, therefore, also consider any

Table 3 Childhood problems and other characteristics of chronic versus acute mood disorders ($N = 397$)

	BP		<i>P</i>	UP		<i>P</i>	BP + UP		<i>P</i>	Oth. $N = 183$
	Chronic ($N = 38$)	Acute ($N = 66$)		Chronic ($N = 43$)	Acute ($N = 67$)		Chronic ($N = 81$)	Acute ($N = 133$)		
Female sex	57.9	62.1	0.67	55.8	62.7	0.47	56.8	62.4	0.42	41.0
Childhood family problems	3.8	2.8	0.08	3.9	2.7	0.05	3.8	2.7	0.008	1.6
Childhood conduct problems	0.8	1.0	0.53	0.9	0.7	0.35	0.8	0.9	0.88	0.4
Sexual trauma	23.7	21.2	0.77	20.9	11.9	0.20	22.2	16.5	0.30	6.6
Self-esteem (age 20)	13.1	14.6	0.02	13.8	13.6	0.46	13.5	14.1	0.22	14.7
Mastery/coping (age 20)	11.9	13.4	0.04	12.3	13.0	0.31	12.2	13.2	0.03	14.6
FH + depression	55.3	65.2	0.32	60.5	67.2	0.47	58.0	66.2	0.23	34.4
SCL depression (age 19)	67.5	64.9	0.27	67.5	65.7	0.28	67.5	65.3	0.11	58.4
SCL anxiety (age 19)	67.8	62.6	0.03	65.8	62.8	0.07	66.8	62.7	0.005	56.6
SCL somatization (age 19)	58.3	56.7	0.46	61.0	54.5	0.005	59.7	55.6	0.01	51.4
SCL interspers. sensitivity (age 19)	68.0	65.3	0.26	67.2	65.6	0.49	67.6	65.5	0.20	59.2
Anxious personality	33.3	23.4	0.29	35.0	17.0	0.05	34.2	20.5	0.03	15.7

Bonferroni threshold $P = 0.0014$

Table 4 Childhood conduct problems and chronicity

Conduct item	BP		UP		BP + UP	
	Chronic ($N = 38$)	Acute ($N = 66$)	Chronic ($N = 43$)	Acute ($N = 67$)	Chronic ($N = 81$)	Acute ($N = 133$)
Childhood conduct problems (mean)	0.8	1.0	0.9	0.7	0.84	0.86
Trouble with police (%)	26.3	28.8	30.2	26.9	28.4	27.8
Restless in class (%)	30.6	40.6	17.5	26.4	23.7	34.2
Unpunctual (%)	11.1	17.2	7.5	3.8	9.2	11.1
Frequent physical fights (%)	11.1	12.5	12.5	3.8	11.8	8.6
Discipline problems at school (%)	16.7	23.4	17.5	13.2	17.1	18.8
Repeated running away (%)	8.3	7.8	7.5	5.7	7.9	6.8
Theft (%)	14.1	8.3	15.0	15.1	11.8	14.5
Truancy (%)	13.9	17.2	17.5	11.3	15.8	14.5

Bonferroni threshold $P = 0.0019$

All P ns

Table 5 Predictors of chronicity: multivariable logistic regression

	BP		UP		BP + UP	
	OR	P^\dagger	OR	P^\dagger	OR	P^\dagger
Childhood family problems	1.8	0.02	1.7	0.05	1.7	0.002
Self-esteem (age 20)	0.5	0.01				
Mastery/coping (age 20)					0.7	0.03
FH + depression	0.4	0.08			0.5	0.04
SCL somatization (age 19)			2.1	0.01		
Anxious personality			4.0	0.02	2.3	0.03

Bonferroni threshold $P = 0.0028$

† Chronic versus acute

statistically significant trends in two of the three comparisons across the three groups: UP, BP, UP + BP. Unsurprisingly, we found two very distinct factors: childhood family problems and conduct problems.

The main positive finding of our study is the consistently strong (1.7-fold) elevation of risk for chronic mood disorders in subjects with childhood family problems. Prospective studies have reported an association between childhood adversity and adult depression [11, 37], but to our knowledge, none has investigated the relationship between early adversity and chronicity of adult mood disorders. A large patient study on dysthymia and double depression by McCullough et al. [30] found no differences

in early adversity between subgroups of chronic depression (major depression, dysthymia, double depression etc.). A smaller patient study by Vocisano et al. [41] reported that chronic UP and BP patients did not differ as regards childhood maltreatment. Our findings extend these studies and suggest that in the community childhood adversity is associated with chronicity of both UP and BP disorders. Finally, the findings that childhood/adolescent anxious personality, low self-esteem, and poor mastery were all associated with chronicity raise the possibility that the latter two factors may be expressions of early trait anxiety, and that all three may mediate the association between early adversity and chronic mood disorder.

In the multivariable analysis, the SCL-90-R (self-assessment) score for anxiety was not significantly associated with chronicity, in contrast to a self-described anxious personality in childhood and adolescence. And a positive family history of depression not only failed to predict but was negatively associated with chronicity, meaning that there were higher positive family history rates in the acute types of mood disorders, as was found for schizophrenia by Bleuler [6].

There were no differences on conduct problem scores between UP and BP subjects or between acute and chronic subtypes of mood disorders, although the expected gender difference [34] was present. These negative findings regarding conduct problems may be surprising; BP subjects might have intuitively been expected to score higher than their UP counterparts. On the other hand, one has to consider that clear mania in childhood is extremely rare. As the excellent Canadian follow-up study of high risk children of parents with BP conducted over 15 years, bipolar disorder was not observed before the age of 12 [16] and in 80% of cases manifested first as depression.

All three diagnostic groups (BP/UP/BP + UP) differed clearly from the residual group without mood disorders in potential risk factors for chronicity (Table 3, statistics not shown)—i.e. family history of depression, female gender, higher scores on childhood family and conduct problems, lower self-esteem and mastery, higher rates of anxious personality, and higher scores on the SCL-90-R scales for anxiety, depression, somatisation and interpersonal sensitivity. The findings regarding anxious personality are reminiscent of the results of the prospective Christchurch Health and Development Study, which found anxious and withdrawn behaviours to be predictive of later anxiety and mood disorders [18] and there is epidemiological confirmation of the familial risks of anxiety [26]. Although we found a number of clear risk factors, most of the variables did not significantly distinguish between the chronic and acute subtypes of mood disorders.

Among the strengths of the study is that it is based on a representative sample, repeatedly assessed prospectively

over 20 years. It is limited, however, by the fact that the childhood variables were assessed retrospectively 15–20 years after the event. Although findings from retrospective studies must be regarded cautiously [21], depressed persons' reports of childhood adversity are stable despite changes in clinical state and over periods of up to 20 years [27, 42]. In addition, adults' retrospective reports of childhood adversity are often corroborated by informants or records [8, 21] and several prospective studies have reported associations between childhood maltreatment and later mood disorders [11, 37]. Furthermore, twin studies indicate that genetic factors cannot explain the association between early adversity and mood disorders [32].

Finally, we have performed many tests of statistical significance, raising the problem of type I errors (false positive tests). Our approach to this issue was to offer a correction for multiple comparisons but nevertheless to present the uncorrected *P*-values so that potentially important findings are not obscured by the correction.

Conclusion

Childhood family problems were strongly associated with chronicity of both bipolar and unipolar mood disorders. Our data are also in line with the hypothesis that childhood adversity may create a vulnerability in the form of childhood anxious personality and low self-esteem and mastery that predispose to chronic mood disorders. However, this requires further analysis and confirmation by other studies.

Acknowledgments This work was supported by Grant 3200-050881.97/1 of the Swiss National Science Foundation.

Conflict of interest statement Prof. Dr. Jules Angst has served on the advisory boards of Eli Lilly & Company, Janssen Cilag, Lundbeck and Sanofi Aventis, on the speakers' bureau of Eli Lilly & Company and AstraZeneca, and as a consultant for Sanofi Aventis. Dr. Alex Gamma, Dr. Vladeta, Prof. Dr. Wulf Rössler, Prof. Dr. Daniel Klein has no conflict of interest to declare.

References

1. Alloy LB, Abramson LY, Smith JM (2006) Role of parenting and maltreatment histories in unipolar and bipolar mood disorders: mediation by cognitive vulnerability to depression. *Clin Child Fam Psychol Rev* 9:23–64
2. Angst J, Gamma A, Baldwin DS, Ajdacic-Gross V, Rössler W (2009) The generalized anxiety spectrum: prevalence, onset, course and outcome. *Eur Arch Psychiatry Clin Neurosci* 259: 37–45
3. Angst J, Gamma A, Benazzi F, Ajdacic V, Eich D, Rössler W (2003) Toward a re-definition of subthreshold bipolarity: epidemiology and proposed criteria for bipolar-II, minor bipolar disorders and hypomania. *J Affect Disord* 73:133–146

4. Angst J, Gamma A, Rössler W, Ajdacic V, Klein DN (2009) Long-term depression versus episodic major depression: results from the prospective Zurich Study of a community sample. *J Affect Disord* 115:112–121
5. Bernet CZ, Stein MB (1999) Relationship of childhood maltreatment to the onset and course of major depression in adulthood. *Depress Anxiety* 9:169–174
6. Bleuler M (1972) Die schizophrenen Geistesstörungen im Lichte langjähriger Kranken- und Familiengeschichten. Thieme, Stuttgart
7. Brown GW, Craig TKJ, Harris TO, Handley RV, Harvey AL (2007) Development of a retrospective interview measure of parental maltreatment using the childhood experience of care and abuse (CECA) instrument—A life-course study of adult chronic depression—1. *J Affect Disord* 103:205–215
8. Brown GW, Craig TKJ, Harris TO, Handley RV, Harvey AL (2007) Validity of retrospective measures of early maltreatment and depressive episodes using the childhood experience of care and abuse (CECA) instrument—A life-course study of adult chronic depression—2. *J Affect Disord* 103:217–224
9. Brown GW, Harris TO, Hepworth C, Robinson RG (1994) Clinical and psychosocial origins of chronic depressive episodes. II. A patient enquiry. *Br J Psychiatry* 165:457–465
10. Brown GW, Moran P (1994) Clinical and psychosocial origins of chronic depressive episodes. I. A community survey. *Br J Psychiatry* 165:447–456
11. Brown J, Cohen P, Johnson JG, Smailes EM (1999) Childhood abuse and neglect: specificity and effects on adolescent and young adult depression and suicidality. *J Am Acad Child Adolesc Psychiatry* 38:1490–1496
12. Derogatis LR, Cleary PA (1977) Factorial invariance across gender for the primary symptom dimensions of the SCL-90. *Br J Soc Clin Psychol* 16:347–356
13. Derogatis RL (1977) Symptom checklist 90, R-version manual I: scoring, administration and procedures for the SCL-90. Johns Hopkins Press, Baltimore
14. Dienes KA, Hammen C, Henry RM, Cohen AN, Daley S (2006) The stress sensitization hypothesis: understanding the course of bipolar disorder. *J Affect Disord* 95:43–49
15. Dougherty LR, Klein DN, Davila J (2004) A growth curve analysis of the effects of chronic stress on the course of dysthymic disorder: moderation by adverse parent-child relationships and family history. *J Consult Clin Psychol* 72:1012–1021
16. Duffy A, Alda M, Hajek T, Grof P (2009) Early course of bipolar disorder in high-risk offspring: prospective study. *Br J Psychiatry* 195:457–458
17. Garno JL, Goldberg JF, Ramirez PM, Ritzler BA (2005) Impact of childhood abuse on the clinical course of bipolar disorder. *Br J Psychiatry* 186:121–125
18. Goodwin RD, Fergusson DM, Horwood LJ (2004) Early anxious/withdrawn behaviours predict later internalising disorders. *J Child Psychol Psychiatry* 45:874–883
19. Green JG, McLaughlin KA, Berglund PA, Gruber G, Sampson NA, Zaslavsky AM, Kessler RC (2010) Childhood adversities and adult psychiatric disorders in the National comorbidity survey replication I: associations with first onset of DSM-IV disorders. *Arch Gen Psychiatr* 67:113–132
20. Hammen C, Henry RM, Daley SE (2000) Depression and sensitization to stressors among young women as a function of childhood adversity. *J Consult Clin Psychol* 68:782–787
21. Hardt J, Rutter M (2004) Validity of adult retrospective reports of adverse childhood experiences: review of the evidence. *J Child Psychol Psychiatry* 45:260–273
22. Heider D, Matschinger H, Bernert S, ESEMEd/MHEDEA 2000 investigators (2006) Relationship between parental bonding and mood disorder in six European countries. *Psychiat Res* 143:89–98
23. Hill J (2003) Childhood trauma and depression. *Curr Opin Psychiatry* 16:3–6
24. Klein DN (2006) Depression and childhood adversity and abuse. *Depress Mind Body* 2:89–93
25. Leverich GS, McElroy SL, Suppes T, Keck PE, Denicoff KD, Nolen WA, Altshuler LL, Rush AJ, Kupka R, Frye MA, Autio KA, Post RM (2002) Early physical and sexual abuse associated with an adverse course of bipolar illness. *Biol Psychiatry* 51:288–297
26. Li XM, Sundquist J, Sundquist K (2008) Age-specific familial risks of anxiety. A nation-wide epidemiological study from Sweden. *Eur Arch Psychiatry Clin Neurosci* 258:441–445
27. Lizardi H, Klein DN (2005) Long term stability of parental representations in depressed outpatients utilizing the parental bonding instrument. *J Nerv Ment Dis* 193:183–188
28. Lizardi H, Klein DN, Crosby Ouimette P, Riso LP, Anderson RL, Donaldson SK (1995) Reports of the childhood home environment in early-onset dysthymia and episodic major depression. *J Abnorm Psychol* 104:132–139
29. Marneros A, Deister A (1990) Chronische depression: Psychopathologie, Verlaufsaspekte und prädisponierende Faktoren. In: Möller HJ (ed) *Therapieresistenz unter Antidepressiva-Behandlung*. Springer, Berlin, pp 1–12
30. McCullough JP, Borian FE, Riso LP, Klein DN, Howland RH, Keller MB (2003) Group comparisons of DSM-IV subtypes of chronic depression: validity of the distinctions, part 2. *J Abnorm Psychol* 112:614–622
31. Molnar BE, Buka SL, Kessler RC (2001) Child sexual abuse and subsequent psychopathology: results from the national comorbidity survey. *Am J Publ Health* 91:753–760
32. Nelson EC, Heath AC, Madden PAF et al (2002) Association between self-reported childhood sexual abuse and adverse psychosocial outcomes: result from a twin study. *Arch Gen Psychiatry* 59:139–145
33. Pearlin LI, Schooler C (1978) The structure of coping. *J Health Soc Behav* 19:2–21
34. Pitzer M, Esser G, Schmidt MH, Laucht M (2009) Temperamental predictors of externalizing problems among boys and girls: a longitudinal study in a high-risk sample from ages 3 months to 15 years. *Eur Arch Psychiatry Clin Neurosci* 259:445–458
35. Post RM, Leverich GS, Xing G, Weiss SRB (2001) Development vulnerabilities to the onset and course of bipolar disorder. *Dev Psychopathol* 13:581–598
36. Scheidegger P (1992) Reproduzierbarkeit von Strukturen in empirischen Daten. Medical Thesis. In, University of Zurich
37. Spatz-Widom C, DuMont K, Czaja SJ (2007) A prospective investigation of major depressive disorder and comorbidity in abused and neglected children grown up. *Arch Gen Psychiatry* 64:49–56
38. Spatz Widom C, Raphael KG, DuMont KA (2004) The case for prospective longitudinal studies in child maltreatment research: commentary on Dube, Williamson, Thompson, Felitti, and Anda (2004). *Child Abuse Negl* 28:715–722
39. van Praag HM, de Kloet R, van Os J (eds) (2004) Epilogue. In: *Stress, the brain and depression*. Cambridge University Press, Cambridge, pp 260–264
40. van Praag HM, de Kloet R, van Os J (eds) (2004) Genetics of depression. In: *Stress, the brain and depression*. Cambridge University Press, Cambridge, pp 59–77
41. Vocisano C, Klein DN, Keefe RSE, Dienst ER, Kincaid MM (1996) Demographics, family history, premorbid functioning, developmental characteristics, and course of patients with deteriorated affective disorder. *Am J Psychiatry* 153:248–255
42. Wilhelm K, Niven H, Parker G (2005) The stability of the parental bonding instrument over a 20-year period. *Psychol Med* 35:387–393