

New species of rust fungi (Uredinales) from South Africa and new observations on known species

Reinhard Berndt

Received: 8 September 2008 / Revised: 19 December 2008 / Accepted: 19 December 2008 / Published online: 23 January 2009
© German Mycological Society and Springer-Verlag 2009

Abstract Four new species and a new variety of rust fungi (Uredinales) are described from South Africa: *Puccinia montis-venenosi* on *Galium spurium* ssp. *africanum* and *G. tomentosum*, *P. naufraga* on *Helichrysum* sp., *Uredo fynbosense* on *Phylica oleifolia* and *Ph. buxifolia*, *Uromyces eclipsis* on *Zygophyllum morgsana*, and *Puccinia lycii* var. *bizonata* on *Lycium* sp. *Puccinia austro-africana* is proposed as a new combination for *P. tetragoniae* var. *austro-africana* on *Tetragonia*. New observations are presented about already known species: The uredinial stage of *Uromyces pentaschistidis* is described; *Uredo monechmatis* is newly reported for South Africa on new hosts and its uredinial morphology is described; a detailed description is provided for the poorly known *P. anthospermi*; the identity of *Uredo zygophylli*, *U. zygophyllina* and *Uromyces dinteri* is discussed; a key is presented for the known *Puccinia* species on *Helichrysum* in South Africa.

Keywords *Galium* · *Helichrysum* · *Phylica* · Rust mycobiota · *Zygophyllum*

Introduction

About 522 species of rust fungi have been reported from South Africa (Berndt 2008a), most of them by the South

African mycologist Ethel Doidge (1950, and references therein). After Doidge, new species and reports were only published sporadically (e.g. Morris et al. 1988; Van Reenen 1995). In the last decade, South African rust fungi and their diversity and taxonomy have received more interest again and a considerable number of new species has been described recently (e.g. Berndt and Uhlmann 2006; Mennicken and Oberwinkler 2004; Wood 2006). Besides describing new species and findings, recent publications tended to present more comprehensive treatments of the rust species of certain host groups (e.g. Berndt 2008b; Mennicken et al. 2005; Wood 2007; Wood and Crous 2005).

In the present paper, four new species and a new variety of rust fungi are described and the taxonomic status and new observations of already known species are reported. The work is mainly based on collections made by A. Rössel, E. Uhlmann and the author in the Western Cape Province of South Africa in 2004 and 2005. The region has a prevalent winter rainfall regime and supports Fynbos, a heath-like shrubland, and Nama-karoo, an arid dwarf-shrub savannah, as major vegetation types (Cowling et al. 1997). Most of the collections were made in these vegetation types with some additional ones from coastal scrub and afro-montane forest.

Materials and methods

Infected plant organs were observed with a Zeiss Stemi SV8 stereo microscope. Spore samples and hand sections were obtained from dried host specimens, mounted in lactophenol on a slide glass and gently heated. The preparations were examined either with a Zeiss ‘Axiophot’ compound microscope and photographs taken with a Zeiss

Taxonomical novelties *Puccinia austro-africana* (Doidge) R. Berndt, *Puccinia lycii* Kalchbr. var. *bizonata* R. Berndt, *Puccinia montis-venenosi* R. Berndt, *Puccinia naufraga* R. Berndt, *Uredo fynbosense* R. Berndt, *Uromyces eclipsis* R. Berndt & A. R. Wood

R. Berndt (✉)
ETH Zurich, Institute of Integrative Biology (IBZ),
CHN D-37, Universitätsstr. 16,
8092 Zürich, Switzerland
e-mail: reinhard.berndt@env.ethz.ch

'MC-80' camera on Kodak Ektachrome 64 Professional slide film or with an Olympus 'BX51' equipped with a 'ColorView IIIu' camera. In the latter case, the 'Cell*B' software package (Software Imaging System) was used to capture and edit micrographs and to measure details of spore walls and their ornament. Such measurements are given to 0.1 μm while ordinary measurements made by the use of an ocular micrometer scale are given to 0.5 μm . Measurements were made on freshly embedded spores or sections. At least 30 spores were measured for each spore state; arithmetic means were calculated and are given in brackets (designated 'mean'). Names of herbaria were abbreviated by their acronyms according to Index Herbariorum (Holmgren et al. 1990). 'RSA' numbers are field collection numbers. The occurring spore states were listed designating the sori according to their position in the rust life cycle. Aecia are indicated by the Roman numeral I, uredinia by II, and telia by III.

The rust species are listed with their host families which are arranged alphabetically.

Results and discussion

Acanthaceae

Uredo monechmatis Cummins, new for South Africa on new host genera and species.

(Bull. Torrey bot. Cl. 79: 231. 1952. As *U. monechmae*).

Material examined: South Africa, Western Cape Prov., Knysna, Harkerville Forest, at 'Kranshoek walk', on *Isoglossa ciliata* (Nees) Lindau (RSA 70, II), *Justicia capensis* Thunb. (RSA 68, II) and cf. *Hypoestes aristata* (Vahl.) Sol. ex Roem. & Schult. (without no., II) in coastal scrub forest, 26 Oct 2004, leg. R. Berndt and E. Uhlmann.

Rust fungi collected on members of *Hypoestes*, *Isoglossa* and *Justicia* in the afro-montane and scrub forest of the Knysna region were identified as *U. monechmatis* based on characters of the urediniospores and the presence of a uredinial peridium with apical paraphyses (uredinia of *Malupa*-type). The urediniospores measured 22.5–28.5 \times 16.5–20 μm (mean 25.3 \times 18.6 μm) on *Justicia capensis* and 24–31 \times 17.5–21 μm (mean 27.0 \times 19.3 μm) on *Isoglossa ciliata*. The plant determined as cf. *Hypoestes aristata* bore only a few uredinia with similar urediniospores that were not measured. *U. monechmatis* is only known so far on *Monechma subsessile* C.B. Clarke from Uganda (Laundon 1963a). A uredinial specimen on *Dicliptera* from Ethiopia listed by Laundon (1963a) is very similar to *U. monechmatis* and might belong to the latter. It has slightly smaller (20–25 \times 15–19 μm) and more delicately echinulate urediniospores (Laundon 1963a).

The morphology of the uredinia (*Malupa*-type) and the pallid, thin-walled urediniospores indicate that *U. monechmatis* may be the uredinial state of a phakopsoraceous species. To my knowledge, this would be the first member of Phakopsoraceae on Acanthaceae. *Isoglossa ciliata* and *Justicia capensis* are new host genera and species for the present rust. *Hypoestes* may be an additional new host genus.

Aizoaceae

Puccinia austro-africana (Doidge) R. Berndt, a new combination for *P. tetragoniae* McAlpine var. *austro-africana* Doidge.


Material examined: *P. tetragoniae* var. *tetragoniae*. Australia, Victoria, Sandringham, on *Tetragonia implexicoma* (Miq.) Hook. f., leg. GH Robinson, Aug 1902, det. D. McAlpine (VPRI 4048a).

P. austro-africana (Doidge) R. Berndt, on *Tetragonia* sp. South Africa, Western Cape Prov.: Cape Peninsula, footpath between Cape Point Lighthouse and parking area, leg. E. Uhlmann and R. Berndt, 1 Nov 2004 (RSA 96, II/III). Vanrhynsdorp, at Vanrhynspass, slopes at lower section of pass road, 31°23'54.1"S, 19°01'40.6"E, leg. A. Rössel, E. Uhlmann and R. Berndt, 11 Oct 2005 (RSA 156, II/III). Clanwilliam, road to Pakhuis Pass, 32°08'07.6"S, 18°58'54.5"E, leg. A. Rössel, E. Uhlmann and R. Berndt, 13 Oct 2005 (RSA 194, II). Struisbaai, Cape Agulhas, 'Spookdraai'-hiking trail, coastal scrub within resort, 34°49'24.0"S, 20°01'29.8"E, leg. A. Rössel, E. Uhlmann and R. Berndt, 30 Oct 2005 (RSA 399, II/III).

Puccinia tetragoniae was originally found in Australia on *T. implexicoma* with spermogonia, aecia, uredinia and telia (McAlpine 1906). The *Puccinia* occurring on *Tetragonia* in South Africa was recognized as a distinct variety within the latter species and named var. *austro-africana* by Doidge (1948). Our collections of *P. tetragoniae* made in several locations of the Western Cape Province tallied well with var. *austro-africana* as described by Doidge. A comparison with *P. tetragoniae* var. *tetragoniae* on the type host from Victoria, Australia (Fig. 1a–c), confirmed the differences between the varieties observed by Doidge (1948). Variety *austro-africana* (Fig. 1d–f) has smaller uredinio- and teliospores with thinner spore walls, especially in the urediniospores (Table 1). The known specimens bear only uredinia and telia in contrast to the macrocyclic *P. tetragoniae* var. *tetragoniae*.

In my opinion, the observable differences merit specific delimitation of var. *austro-africana* and I propose the new combination *Puccinia austro-africana* (Doidge) R. Berndt, comb. nov., based on *P. tetragoniae* var. *austro-africana* Doidge (Bothalia 4: 904/905. 1948. Type on *T. expansa* Thunb., PREM 34095).

Fig. 1 a–c *Puccinia tetragoniae* (VPRI 4048a). d–f *Puccinia austro-africana* (RSA 399). Teliospores (a,d), urediniospores in optical section (b,e) and urediniospores with focus on spore surface (c,f). Asterisks indicate urediniospores of *P. tetragoniae* in which the indentation of the spore lumen is especially visible. It is less pronounced in *P. austro-africana* due to its thinner spore wall (arrows). Bars=20 μ m



Asteraceae

Puccinia on *Helichrysum* (*Inuleae*, *Gnaphalinae*)

Doidge (1927) distinguished within *P. kalchbrenneri* De Toni a variety *valida* for specimens with longer teliospores and a more strongly thickened teliospore apex. Berndt and Uhlmann (2006) studied specimens of both varieties and observed that the characters intergraded. They concluded that *P. kalchbrenneri* was a variable species and that var. *valida* was not clearly distinguishable from var. *kalchbrenneri*.

New collections of *P. kalchbrenneri* from the Western Cape Province also varied considerably in several characters of the telio- and urediniospores. Differences were found with regard to the size of the teliospores and traits of their spore wall, the size of the urediniospores, the thickness and echinulation of their spore walls and the number and visibility of the germ pores. Despite the observable differences most of the specimens exhibited an overall similarity that did not allow clear distinguishing of separate morphs. One specimen was sufficiently different to warrant specific delimitation and is described as *P. naufraga*.

Puccinia naufraga R. Berndt, sp. nov.

Figures 2, 3 and 4.

Etymology: Named after the collection site, the ‘Thomas Tucker Shipwreck Trail’, Cape Peninsula.

Aecia singularia vel paucula in pagina abaxiali foliorum, *peridio* albo anguste cylindrico praedita; **aeciosporae** late ellipsoideae ad subgloboseae, saepe leniter deformatae et subangulares, 23–32×20–25 μ m (27.2×22.5 μ m), *pariete* hyalino vel subhyalino ca. 2 μ m crasso regulariter verruculoso, *cellulae peridii* intus verrucis irregularibus delicatis usque ad moderate grossis dense obsitae, extus fere leves vel papillosae verrucis planis et inconspicuis. **Uredinia** in pagina abaxiali foliorum sparsa, singularia vel aliquot aggregata, obscure brunnea basaliter atro-brunnea, cupulata, apertura centrali praedita, *peridio* cupuliformi e cellulis compressis paucistratis brunneis composito, copia sporarum cinnamomea vel ferruginea; **urediniosporae** subgloboseae ad obovoideae, 23–28×21–24 μ m (25.4×22.6 μ m), *pariete* ca. 1.5–2 μ m crasso inconspicue bilaminato, lamina exteriori angustissima subhyalina vel straminea regulariter echinulato spinis delicatis ca. 2–3 μ m distantibus, lamina interiori crassiore dilute brunnea, poris

Table 1 Characters of uredinio- and teliospores of *Puccinia tetragoniae* and *P. austro-africana*

Species, specimen	Urediniospores (means)	Teliospores (means)
<i>P. tetragoniae</i> (from McAlpine 1906)	27–37×22–25 μm (31×24 μm)	40–60×25–35 μm (50×30 μm)
<i>P. tetragoniae</i> (from Laundon 1963b, sub var. <i>tetragoniae</i>)	29–40×27–39 μm, spore wall 3–7 μm thick	40–60×27–40 μm, spore wall 2–5 μm thick, at apex 4–10 μm
<i>P. tetragoniae</i> Australia, Victoria (VPRI 4048a)	30–34×27–32 μm (32.3×29.1 μm), spore wall 3–4 μm thick, at germ pores 7–8 μm, germ pores scattered, 7–10	46–64×29–40 μm (54.1×35.2 μm), spore wall 4–5.5 μm thick, at apex 6–10 μm, pedicel more or less persistent
<i>P. tetragoniae</i> (from Cunningham 1931)	28–36×22–28 μm (30×25 μm), spore wall 3 μm thick, germ pores scattered, 5–6	52–60×32–40 μm (56×36 μm), spore wall to 5 μm thick, at apex to 8 μm or not thickened, pedicel persistent
<i>P. austro-africana</i> (from Doidge 1948, sub var. <i>austro-africana</i>)	20–26×17.5–20 μm (in Latin description), 22.5–32×17.5–25 μm (in English description), spore wall 1.5–2 μm thick, germ pores scattered, 5–6	30–50×22.5–35 μm, spore wall 3–4 μm thick, at apex 5–6(8), pedicel persistent
<i>P. austro-africana</i> (from Laundon 1963b, sub var. <i>austro-africana</i>)	22–25×19–22 μm (23.2×20.8 μm), spore wall 1.5–4 μm thick	Not specified
<i>P. austro-africana</i> South Africa (RSA 96)	23–28×20–23 μm (25.4×21.7 μm), spore wall 1.5–2 μm thick, at germ pores 3–4 μm, germ pores scattered, 6–8	38–50×26–32 μm (43.3×29.3 μm), spore wall ca. 2 μm thick, at apex 3.5–7 μm, pedicel frail
<i>P. austro-africana</i> South Africa (RSA 156)	25–34×17–22.5 μm (28.3×20.1 μm), spore wall ca. 2 μm thick, at germ pores up to 4 μm, germ pores scattered, 5–8	37–45(47)×24–30 μm (42.2×27.3 μm), spore wall 3–3.5 μm thick, at apex 5–8 μm, pedicel frail
<i>P. austro-africana</i> South Africa (RSA 194)	24–28×21–23 μm (26.0×21.8 μm), spore wall ca. 2–2.5 μm thick, at germ pores 3.5–4.5 μm, germ pores scattered, 5–7	Not present
<i>P. austro-africana</i> South Africa (RSA 399)	24–30×20.5–23.5 μm (27.5×22.0 μm), spore wall ca. 2 μm thick, at germ pores 5–6 μm, germ pores scattered, 5–7	39–51×(22.5)26–32 μm (44.8×28.7 μm), spore wall 2.5–3.5 μm thick, at apex 6–8 μm, pedicel frail


germinationis inconspicuis 5–6 sparsis papillis latis humilibus praeditis. **Telia** abaxialia gregaria, mox nuda, rotundata, pulvinata, primum aurantiaco-brunnea deinde saturate ferruginea, textura carnosa vel semi-compacta; **teliosporae** post maturitatem germinantes, ellipsoideae ad subclavatae rariter late ellipsoideae, ad septum leniter constrictae, 44–70×21–27 (56.1×23.9 μm), cellula distali ovoidea late ellipsoidea vel obpyriformi, cellula proximali late ellipsoidea subglobosa vel hilum versus attenuata, *pariete* levi pallide aurantiaco-brunneo ca. 1 μm crasso in cellula proximali, 1–1.5 μm in cellula distali, 4–6 μm in apice, usque ad 4 μm in poro germinationis cellulae proximalis, poris germinationis in apice et septum juxta; *mesosporae* sparsae, 29–43×(17)20–24 μm; *pedicello* leniter incrassato stramineo vel pallide brunneo usque ad 55 μm longo.

In foliis *Helichrysi* sp. (Asteraceae)

Aecia singly or in small groups on abaxial side of leaves, with white slenderly cylindrical peridium; **aeciospores** broadly ellipsoid to subglobose, often slightly deformed and subangular, 23–32×20–25 μm (mean 27.2×22.5 μm), spore wall hyaline to subhyaline, ca. 2 μm thick (incl. ornament), rather evenly and finely verrucose; inner side of peridial cells with irregular, dense, fine to moderately

coarse warts, outer side essentially similar but warts flat and inconspicuous giving the surface an almost smooth to papillose appearance. **Uredinia** scattered, often singly, more rarely in small groups on abaxial side of leaves, dark brown, at the base blackish-brown, cupola-shaped with central aperture, provided with an overarching peridium composed of several layers of compressed, brown cells, spore mass cinnamon to ferruginous; **urediniospores** subglobose to obovoidal, 23–28×21–24 μm (mean 25.4×22.6 μm), spore wall ca. 1.5–2 μm thick, inconspicuously bilaminate with a very thin exterior, subhyaline to straw-coloured layer and a thicker, pale and dull brown inner layer, evenly echinulate with rather fine spines about 2–3 μm apart, germ pores inconspicuous, 5–6, scattered, with broad, flat caps. **Telia** abaxial, in small groups, early naked, round, pulvinate, first orange-brown, later ferruginous, soft to semi-compact; **teliospores** ellipsoid to subclavate, rarely broadly ellipsoid, slightly constricted at septum, 44–70×21–27 (mean 56.1×23.9 μm), distal cell ovoid to broadly ellipsoid or obpyriform, proximal cell broadly ellipsoid, subglobose or almost wedge-shaped, tapering towards the hilum, spore wall smooth, light brown with orange tinge, ca. 1 μm thick in proximal cell, 1–1.5 μm in the distal one, thickening to 4–6 μm at the apex and to 4 μm at the germ pore of the proximal cell, germ pores

Fig. 2 *Puccinia naufraga* (type). **a** Urediniospores in optical section. **b** Lateral part of uredinium showing peridial cells (stars) and the base of the cupola-like cover (arrow). **c** Teliospores and a single one-celled mesospore (star). **d** Teliospores. Arrows in **c** and **d** point to the teliospore apex which is less pigmented than the remaining spore wall and shows an inconspicuous lenticular swelling. Bars=10 μ m



apical and adjacent to the septum, with a thickening that is less pigmented than the remaining wall and most often lenticular in the distal cell, germinating upon maturity; one-celled mesospores scattered, $29\text{--}43 \times (17)20\text{--}24$ μ m, with pale brown, ca. 1 μ m thick wall, slightly thicker at apex; pedicel broadly attached, slightly thick-walled, straw-coloured to pallid brown, up to 55 μ m long.

On leaves of *Helichrysum* sp. (Asteraceae).


Holotype (PREM 60146): South Africa, Western Cape Prov., Cape Peninsula S of Simons' Town, 'Thomas Tucker Shipwreck Trail', $34^{\circ}15'56.3''\text{S}$, $18^{\circ}23'04.5''\text{E}$, alt. ca. 5 m a.s.l., on *Helichrysum* sp., leg. A. Rössel, E. Uhlmann and R. Berndt, 6 Nov 2005 (RSA 431, I/II/III).

P. naufraga differs from *P. kalchbrenneri* by less pigmented, orange-brown teliospore walls, a thinner teliospore apex and the apical germ pore with a small, lenticular papilla-like thickening (Figs. 2c–d, 4) which is less

pigmented than the remaining wall. The thickening is visible until germination of the teliospores. Similar thickenings, though more pronounced, also occur in *P. macowani* but which differs by aeciospores with broad, flat warts and by the lack of a uredinial stage. In *P. kalchbrenneri*, the teliospore apex has a more or less pronounced conical pit in the spore wall probably indicating the location of the germ pore. It has not been proven so far that *P. kalchbrenneri* has an aecial stage though Doidge (1927) assumed that it may be represented by *Aecidium helichrysi* Doidge. Aeciospores of *P. naufraga* are hardly distinguishable from *A. helichrysi* but have slightly thinner spore walls which tend to be more finely verruculose.

Both *P. naufraga* and *P. kalchbrenneri* possess peridiate uredinia (Fig. 2b). The peridium first covers young uredinia like a cupola but later becomes evanescent. It is built up by one to several layers of compressed hyphae that become

Fig. 3 *Puccinia naufraga* (type). **a** Aeciospores in optical section. **b** Aeciospores, focus on spore surface showing verruculose ornament. **c** Peridial cells, surface of inner tangential wall. **d** Peridial cells, surface of outer tangential wall. Bars=20 μ m



very indistinct towards the apex of the peridium which hence appears more or less amorphous in section and face view. The presence of the peridium results in a characteristic bullate to broadly flask-like shape of the uredinia which liberate the spores through an irregular pore or slit at their apex. Among *Puccinia* species on *Helichrysum* the presence of a uredinial peridium was first reported in *P. cornurediata* R. Berndt which is characterised by conspicuous horn-like peridia (uredinia of *Uredostilbe*-type). As discussed by Berndt and Uhlmann (2006), this character has been used to delimit the genus *Miyagia* from *Puccinia*. In young uredinia of *P. kalchbrenneri* and *P. naufraga*, the peridia resembled the similar if more delicate dome-shaped uredinial peridia of members of Pucciniaceae, *Cronartium* and *Melampsora* (Moss 1926, 1928). Such uredinia have been assigned to the anamorph genus *Milesia*. It is evident that the occurrence of morphologically similar

uredinia does not indicate a closer relationship between the discussed *Puccinia* species and members of the named taxa.

The following key comprises *Puccinia* species known from South Africa on *Helichrysum* species. It should be noted that it may be impossible to determine a specimen with only a single spore state.

- 1 Teliospores with very irregularly thickened, knotty or bulged cell wall, germ pores depressed 1/3–2/3 in both cells; uredinia unknown *P. pienarii*
- 1* Teliospores with other characters, germ pores apical and close to septum 2
- 2 Teliospores uniformly thin-walled, straw-coloured to ochraceous; aeciospores with flat, button-like warts; uredinia unknown *P. horti-kirstenboschi*
- 2* Teliospores with apically thickened wall, straw-coloured to chestnut brown 3




Fig. 4 *Puccinia naufraga* (type). Teliospores, among them three one-celled mesospores. Bar=20 µm

- 3 Uredinia with a slenderly cylindrical to subconical peridium up to 1 mm long, composed of linear, golden, thick-walled hyphae *P. cornurediata*
- 3* Uredinia with bullate, cupola-like, delicate peridium, not peridiate (?), or uredinia unknown 4
- 4 Telia ferruginous; teliospores light to orange brown, spore apex with a small to large lenticular to subglobose pallid swelling, only rarely with a conical pit 5
- 5 teliospores ellipsoidal to fusiform, with a subglobose or broadly ellipsoidal thickening at germ pores; aeciospores with broad, flat, button-like warts; uredinia unknown *P. macowani*
- 5* teliospores ellipsoidal, with orange brown wall and a small lenticular swelling at the spore apex; uredinia with cupulate peridium; aeciospores finely and densely verrucose *P. naufraga*
- 4* Telia apricot, ferruginous or blackish brown; teliospores apically thickened, with or without conical pits at the germ pores but not with lenticular or subglobose apical swelling 6
- 6 Telia apricot to ferruginous, erumpent; teliospores fusiform to ellipsoidal with straw-coloured wall and an oblique thickening at the apex, often conspicuously thickened in the angles of the septum; uredinia unknown *P. subindumentana*
- 6* Telia dark brown to blackish brown, erumpent (to crustose); teliospore wall more deeply pigmented,

- apically thickened but generally not obliquely; uredinia known 7
- 7 Urediniospore wall 2–3 µm thick, yellow-brown; teliospores relatively broad (21–37 µm); aecia unknown *P. rocherpaniana*
- 7* Urediniospore wall 1–2 µm thick, straw-coloured to light orange brown; teliospores narrower on average; aecial stage maybe present and is possibly identical with *Aecidium helichrysi* *P. kalchbrenneri* (vars.)

Poaceae


Uromyces on *Pentaschistis*

Figure 5.

Material examined: South Africa, Western Cape Prov., Cederberge Mountains, Driehoek, on *Pentaschistis pallida* (Thunb.) Linder, leg. E. Uhlmann and R. Berndt, 12 Oct 2004 (RSA 5, II/III). South Africa, Western Cape Prov., Vanrhynsdorp, viewpoint above Vanrhynspass, 31°22'19.1"S, 19°01'02.3"E, alt. ca. 825 m a.s.l., on *P. cf. pallida*. leg. A. Rössel, E. Uhlmann and R. Berndt, 11 Oct 2005 (RSA 145, II/III). South Africa, Western Cape Prov., Vanrhynsdorp, parking bay at Vanrhynspass, on *P. airoides* Stapf, leg. A. Rössel, E. Uhlmann and R. Berndt, 11 Oct 2005 (RSA 168, II/III).

Gjærøum (1988) described *U. pentaschistidis* on *P. airoides* from South Africa based on the telial state. We collected three specimens of *Uromyces* on *P. airoides* and *P. pallida* which revealed telio- and urediniospores. The teliospores from both host species were very similar but slightly smaller on *P. airoides* (Table 2). They agreed quite well with the description of *U. pentaschistidis* given by Gjærøum though he described thicker spore walls (3–4 µm at sides vs ca. 2 µm, up to 8 µm at the apex vs 4–7 µm). The urediniospores obtained from the present specimens were similar by their globose to broadly ellipsoid or broadly obovoid shape, by ca. 1.5–2 µm thick walls which are evenly and rather densely echinulate by delicate spines, and by the presence of 9–14 scattered germ pores. The spores differed, however, among the three specimens with regard to their size and the wall thickening at the germ pores (Table 2). The urediniospores taken from *P. airoides* (RSA 168, Fig. 5a,b) were considerably smaller than those from *P. pallida* (Fig. 5c,d) and showed conspicuous thickenings at the germ pores. More material needs to be studied, however, to decide whether the observed differences are constant. Should it turn out that the rusts on *P. pallida* and *P. airoides* are different species it would be difficult to find out which one belongs to *U. pentaschistidis*. The teliospores are too similar in all specimens to allow a safe distinction. Urediniospores, however, which do seem to be

Fig. 5 *Uromyces pentaschistidis*. **a** Urediniospores, focus on spore surface (RSA 168, on *Pentaschistis airoides*). **b** Urediniospores in optical section (RSA 168). Note internal swellings of spore wall at germ pores. **c** Urediniospores, focus on spore surface (RSA 145, on *P. pallida*). **d** Urediniospores in optical section (RSA 145). The spore wall is only slightly swollen at the germ pores. Bars=20 μm



reasonably different are unknown in *U. pentaschistidis*. Unfortunately, type material of *U. pentaschistidis* could not be obtained and searched for the presence of urediniospores. Therefore, I provisionally assign all studied specimens to *U. pentaschistidis*.

Rhamnaceae

Uredo fynbosense R. Berndt, *sp. nov.*

Figure 6.

Etymology: Fynbos, habitat of the known hosts of the present species.

Spermogonia, aecia et telia absentia. **Uredinia** indumento lanoso coperta, solitaria vel laxe sparsa in pagina abaxiali foliorum, subepidermalia, minuta, pallide cinnamomea, pulverulenta, hyphis paucis paraphysiformibus tenue tunicatis praedita sed structuras proprias carentia

quas soros circumdant; **urediniosporae** praecipue obovoideae, magis rariter ellipsoideae ad late ellipsoideae vel subclavatae, 19–28(31)×14–18 μm (23.5×16.1 μm), *pariete* pallide aureo ca. 1 μm crasso non vel lenissime incrassato apicem versus, aequaliter et sate delicate echinulatae spinis inter se 1.5–2.5 μm distantibus, minus delicate et densiore echinulatae hilum versus, poris germinationis obscuris verosimiliter 2–3(4?) approx. aequatorialibus.

In foliis *Phyllicae* specierum.

Spermogonia, aecia and **telia** not present. **Uredinia** singly or sparsely scattered on abaxial side of leaves, hidden under the woolly indument, subepidermal, tiny, pallid cinnamon brown, pulverulent, without clearly differentiated binding structures though a few thin-walled, hyaline, cylindrical paraphysis-like hyphae are sometimes present which do not appear to represent spore pedicels; **urediniospores** mostly obovoid, more rarely ellipsoid to broadly ellipsoid, subclavate, 19–28(31)×14–18 μm (mean

Table 2 Comparison between uredinio- and teliospores of *Uromyces* on *Pentaschistis pallida* and *P. airoides*

Host (specimen)	Urediniospores (means)	Urediniospores, germ pores	Teliospores (means)
On <i>P. pallida</i> (RSA 5)	(25)27–31.5×22.5–26 μm (30.2×24.2 μm)	9–12, scattered, spore wall slightly thickened at pores	21.5–28(30)×17–22 μm (24.7×20.0 μm)
On <i>P. pallida</i> (RSA 145)	27–30×24.5–28 μm (28.7×26.4 μm)	10–12, scattered, spore wall slightly thickened at pores	21–27×(15)17–23 μm (24.2×19.7 μm)
On <i>P. airoides</i> (RSA 168)	21–25(27)×20.5–23 μm (23.4×21.6 μm)	11–14, scattered, spore wall conspicuously thickened at pores	19–24×16.5–21 μm (21.4×19.6 μm)
<i>Uromyces pentaschistidis</i> on <i>P. airoides</i> (from Gjørørum 1988)	Unknown		(19)22–24×19–22 μm (mean not calculated)

23.5×16.1 μm), spore wall pallid golden yellow, ca. 1 μm thick, not or hardly thickened at apex, more or less evenly and rather finely echinulate with spines ca. 1.5–2.5 μm apart, somewhat more closely and less delicately echinulate towards the hilum, germ pores obscure, probably 2–3(4?), approximately equatorial.

On leaves of *Phylica* spp.

Holotype (PREM 60149): South Africa, Western Cape Prov., Vanrhynsdorp, ascent to Gifberg, 31°46'31.1"S, 18°45'51.7"E, ca. 660 m a.s.l., on *Phylica* cf. *oleifolia* Vent., 12 Oct 2005, leg. A. Rössel, E. Uhlmann and R. Berndt (RSA 183, II). Isotype Z+ZT (ZT Myc 1270).

Additional material studied: South Africa, Western Cape Prov., Goukamma Nature Reserve SW of Groenvlei lake E of Lake Pleasant, 34°02'10.4"S, 22°50'14.7"E, alt. ca. 20 m a.s.l., on *Phylica* cf. *buxifolia* L., 26 Oct 2005, leg. A. Rössel, E. Uhlmann and R. Berndt (RSA 372, II).

Phylica is a genus of about 150 species of which 134 occur in the Cape Floristic Kingdom, the vast majority of them endemic to the region (Linder 2003). The only rust fungus described from a member of this genus is *Uredo phylicae* Vienn.-Bourg. from the island of New Amsterdam.


According to the description given by Viennot-Bourgin (1970) it differs from the present species in much broader urediniospores (25–31.5×24–29 μm , mean 28×26 μm) that are closely verruculose and have 5–7 germ pores. The affiliation of *U. fynbosense* with a telial genus is unknown but its general appearance indicates that it may be a member of Phakopsoraceae. Known *Phakopsora* species on Rhamnaceae are *Ph. colubrinae* Viégas and *Ph. zizyphi-vulgaris* Dietel, the latter with peridiate and paraphysate uredinia (Yen 1975), the former with unbounded uredinia and urediniospores similar to the present species (Viégas 1960).

Rubiaceae

Puccinia species on members of Rubioideae, tribes Rubieae and Anthospermeae

Findings of rust fungi on *Galium* species (Rubiaceae) from Western Cape Province revealed the widespread *P. punctata* Link and a similar, yet different *Puccinia* species which is described as *P. montis-venenosi*.

Fig. 6 *Uredo fynbosense* (holotype). **a** Urediniospores in optical section. **b** Urediniospores, focus on spore surface. Bars=20 μm



Puccinia montis-venenosi R. Berndt, *sp. nov.*

(Fig. 7)

Etymology: after the ‘Gifberg’ (= poison mountain), collection site of the type.


Spermogonia et **aecia** ignota. **Uredinia** praecipue in pagina foliorum abaxiali vel caulicola, subepidermalia, ferruginea, pulverulenta, telios evolvere possunt; **urediniosporae** late obovoideae late ellipsoideae, subglobosae, rariter late ovoideae vel globosae, $24\text{--}35 \times 21\text{--}26 \mu\text{m}$ ($29.2 \times 23.2 \mu\text{m}$), *pariete* aurantio-brunneo ca. $1.5(2) \mu\text{m}$ crasso leniter incrassato (usque ad $3 \mu\text{m}$) hilum versus sate delicate echinulato spinis ca. $1.8\text{--}3.2 \mu\text{m}$ inter se distantibus evanescentibus in dimidio inferiore sporarum in quarta vel quinta parte inferiore carentibus, poris germinationis 2–3 plusminusve aequatorialibus et aequidistantibus, papillis parvis et inconspicuis praeditis. **Telia** nigrescentia, subcompacta, pulvinata, rotundata vel elongata in caulibus; **teliosporae** clavatae, ellipsoideae, oblongae vel subfusi-

formes, $(32)36\text{--}70 \times 16\text{--}27 \mu\text{m}$ ($52.9 \times 21.1 \mu\text{m}$), non vel paulum constrictae in septo, apice rotundato conico vel – rariore – subtruncato rostellato vel irregulariter deformato, cellula proximali attenuata hilum versus, *pariete* levi sate aureo ad dilute castaneo (praecipue apicem versus) lateraliter ca. $2 \mu\text{m}$ crasso in cellula proximali, $2\text{--}3 \mu\text{m}$ in cellula distali, in apice $10\text{--}20 \mu\text{m}$ crasso, poris germinationis apicalibus et juxta septum, papillis carentibus, *pedicello* ad hilum late affixo ($7\text{--}10 \mu\text{m}$) tenue tunicato dilute brunneo vel subhyalino fragili sed longitudinem usque ad $70 \mu\text{m}$ attingenti.

In foliis caulibusque *Galii* specierum.

Spermogonia and **aecia** unknown. **Uredinia** predominantly on abaxial side of leaves or on stems, subepidermal, ferruginous, pulverulent, may give rise to telia; **urediniospores** broadly obovoid, broadly ellipsoid, subglobose, rarely broadly ovoid or globose, $24\text{--}35 \times 21\text{--}26 \mu\text{m}$ (mean $29.2 \times 23.2 \mu\text{m}$), spore wall orange brown, ca. $1.5(2) \mu\text{m}$ thick, slightly thicker (up to $3 \mu\text{m}$) towards hilum, rather

Fig. 7 *Puccinia montis-venenosi* (paratype). **a,b** Teliospores showing a part of the variation with regard to teliospore size, shape and thickness of the spore apex. **c** Urediniospores in optical section. Note smooth spore base (arrows). **d** Urediniospores, focus on spore surface. Note smooth spore base (arrows) Bars = $50 \mu\text{m}$ (**a**), $20 \mu\text{m}$ (**b–d**)



finely echinulate with spines ca. 1.8–3.2 μm apart, spines become smaller on proximal half of spores whilst proximal fourth or fifth is smooth, germ pores 2–3, more or less equatorial and equidistant, with very small and inconspicuous caps. **Telia** blackish, subcompact, pulvinate and rounded, generally elongated on stems; **teliospores** clavate, ellipsoid, oblong or subfusiform, (32)36–70 \times 16–27 μm (mean 52.9 \times 21.1 μm), not or hardly constricted at septum, apex rounded, conical, more rarely subtruncate, beaked or irregularly deformed, proximal cell tapering towards the hilum, spore wall smooth, deeply golden brown to light chestnut brown (especially in thickened apex), laterally about 2 μm thick in proximal cell, 2–3 μm in distal cell, apex 10–20 μm thick, germ pores apical and at septum, without papillae, pedicels up to 70 μm long or breaking off shorter, collapsing or not, broadly attached (7–10 μm) to spore base, more or less thin-walled, pale brownish or subhyaline.

On leaves and stems of *Galium* spp.

Holotype (PREM 60147): South Africa, Western Cape Prov., Vanrhynsdorp, gravel road to Gifberg/Matzikamma Mountains. SSE of Vanrhynsdorp, ascent to Gifberg, on *Galium* cf. *spurium* L. ssp. *africanum* Verdc., leg. A. Rössel, E. Uhlmann and R. Berndt, 12 Oct 2005 (RSA 178, II/III). Isotype Z+ZT (ZT Myc 1268). Paratype: South Africa, Western Cape Prov., Vanrhynsdorp, at Vanrhynspass, slopes at parking bay in middle section of pass road, on *Galium* cf. *tomentosum* Thunb. (non-flowering plant), leg. A. Rössel, E. Uhlmann and R. Berndt, 11 Oct 2005 (RSA 170, II/III).

Additional material examined: *Puccinia punctata*: South Africa, Western Cape Prov., Ceres, ascent to Gydopass, on two collections of *Galium capense* Thunb. cf. ssp. *capense* in Fynbos and roadside vegetation, leg. A. Rössel, E. Uhlmann and R. Berndt, 18 Oct 2005 (RSA 249 and 251, each II/III).

Puccinia montis-venenosi resembles *P. punctata* by the appearance of the uredinia and the telia and by similar uredinio- and teliospores. It differs essentially by the echinulation of the urediniospores which becomes finer towards the base of the spores and vanishes entirely on the proximal fourth or fifth of the spore surface (Fig. 7c, d). In *P. punctata*, the urediniospores are evenly echinulate. It should be noted that Klebahn (1914) observed partially smooth urediniospores, too, in specimens tentatively assigned to *P. deminuta* Vleugel on *Galium uliginosum* L. and *G. palustre* L. The smooth patches occurred mainly lateral on the spores, however, and did not comprise their proximal parts. Gäumann (1959) considered *P. deminuta* to be a *forma specialis* of *P. punctata*.

The investigated specimens were collected on two species of *Galium* and differed considerably from each other with regard to teliospore size. In the specimen from

Vanrhynspass, teliospores measured 48–70 \times 19–27 μm (mean 57.2 \times 22.7 μm), in that from Gifberg (32)36–60 \times 16–22(24) μm (mean 48.8 \times 19.7 μm); the respective urediniospores measured 26–35 \times 21–24 μm (mean 30.6 \times 22.4 μm) and 24–31(33) \times 22–26 μm (mean 27.8 \times 24.0 μm). It may be interesting that *P. punctata* also has uredinio- and teliospores of rather variable size (Gäumann 1959).

Puccinia punctata from Gydopass revealed uredinia and telia. Urediniospores measured 22–28 \times 19–24 μm (mean 24.5 \times 21.3 μm), the teliospores (38)40–60 \times 18–25 μm (mean 47.8 \times 20.5 μm). The spore size and other characters tally well with descriptions of *P. punctata* by various authors (e.g. Gäumann 1959; Klebahn 1914). *P. punctata* is known from many different *Galium* species (Gäumann 1959) and has in addition been reported on members of *Asperula*, *Rubia* and *Relbunium*. The rust is autoecious but shows plasticity with regard to its life cycle with macrocyclic and demicyclic variants (Wurth 1905; Gäumann 1959). In South Africa, it was listed by Doidge (1927, 1950) on *G. capense*, with spermogonia, aecia and uredinia present. Only uredinia and telia are known so far for *P. montis-venenosi* and it is uncertain at this stage whether it is macrocyclic.

Puccinia anthospermi Syd.


Figure 8.

Material examined: South Africa, Western Cape Prov., Elim, Geelkop Hills, 34°33'46.7"S, 19°47'41.7"E, on *Anthospermum galioides* Rchb. f. ssp. *galioides*, leg. A. Rössel, E. Uhlmann and R. Berndt, 29 Oct 2005 (RSA 392, III).

Puccinia galopinae Cooke: South Africa, KwaZulu-Natal, Inanda, on *Galopina aspera* Sond., leg. GM Wood (no. 602), Jun 1881 (PREM 11192 and 10474, types, each with III).

Two *Puccinia* species are known on southern African Anthospermeae: *P. anthospermi* on *Anthospermum* species and *P. galopinae* on *Galopina* species. In the original description of *P. anthospermi* on *A. hirtum* Cruse from South Africa, Sydow and Sydow (1904) described the teliospores as 35–46 \times 16–21 μm large, apically thickened to 9 μm , with a pedicel up to 30 μm long. Holm (1973) measured 40–50 \times 12–15 μm for the teliospores, an apical thickening of up to 10 μm and pedicels up to 40 μm long on *A. muricatum* from Tanzania [this name does not occur in 'The International Plant Names Index' (2004) or Puff (1986). In 'World check list of Rubiaceae' (Govaerts et al. n.d.), *A. muriculatum* Hochst. ex A. Rich. (= *A. herbaceum* L. f.) is listed which may be meant]. He re-studied the "very poor" type collection and could not find essential differences between the specimens.

Fig. 8 *Puccinia anthospermi*, teliospores (RSA 392). Arrow shows a single, three-celled spore. Stars indicate germinated teliospore cells Bars=20 μ m



As the available descriptions of *P. anthospermi* are short, a more detailed one is provided based on the specimen on *A. galioides* ssp. *galioides*:

Telia abaxial on leaves, subepidermal, pallid to dark ferruginous, strongly pulvinate, compact, discrete or confluent, up to 2 mm diam., with a white or greyish cast after germination of basidia; **teliospores** 38–55 \times 14–18.5 μ m (mean 46.8 \times 16.4 μ m), ellipsoid, more rarely fusiform or oblong, two-celled (very rarely three-celled), at septum not or hardly constricted, apex conical, narrowly conical, more rarely rounded, proximal cell tapering towards the pedicel, spore wall smooth, pale golden brown to straw-coloured, ca. 1–2 μ m thick at sides, 4–9 μ m at spore apex which is slightly paler than the remaining wall, germ pores apical in distal cell, close to septum in proximal cell, spores germinate upon maturity, pedicels subhyaline, slightly thick-walled, up to 80 μ m long.

A very similar species is *P. galopinae* whose teliospores are described by Doidge (1927) to be 35–50 \times 11–15 μ m, apex thickened to 7 μ m and with pedicels up to 70 μ m long. I measured (36)40–60 \times 12.5–18 μ m (mean 51.1 \times 15.0 μ m), an apical thickening of 5–9 μ m and pedicels up to 50 μ m long. Apart from slightly narrower and longer teliospores there is little to differentiate *P. galopinae* from *P. anthospermi*. Given this similarity and the close relationship of the host genera, one might assume that these rusts are conspecific. In this case, the older name *P. galopinae* would take precedence over *P. anthospermi*. *P. anthospermi* is known from South Africa and Tanzania (Holm 1973), *P. galopinae* from South Africa (Doidge 1950) and Nigeria (Eboh 1984). Eboh (1984) reported the rust on *Victaria* sp. This name does not seem to exist and is most probably an error for *Virectaria* Afzel. ex Sm. em. Bremek. which belongs to tribe Sabiceae of subfam. Ixoroideae (Khan et al. 2008).

Solanaceae

Puccinia lycii Kalchbr. var. *bizonata* R. Berndt, var. nov.

Figure 9.

Ab varietate typica *Puccinae lycii* differt urediniosporis brevioribus latioribus poris germinationis bizonate positis.

In foliis *Lycii* sp.

Uredinia almost entirely replaced by telia; **urediniospores** ferruginous, ellipsoid, more rarely broadly ellipsoid, narrowly obovoid or almost oblong, 37–53 \times 21–24 μ m (mean 43.6 \times 22.3 μ m), spore wall ochraceous, ca. 1.5 μ m thick, apically to 2 μ m, echinulate by spines ca.




Fig. 9 *Puccinia lycii* var. *bizonata* (holotype). Urediniospores with bizonate germ pores. Bar=20 μ m

2–3.5 μm apart, becoming smooth in proximal fifth of spore surface, germ pores bizonate, rings each with 4–6(7) pores, rarely pores almost scattered, with small, flat subhyaline caps and slight internal thickening of the wall. **Telia** amphigenous on leaves, black, pulvinate and pulverulent; **teliospores** broadly ellipsoid, more rarely ellipsoid or obovoid, rounded at apex and towards hilum, sometimes tapering towards hilum, not or hardly constricted at septum, 39–52 \times 24–30 μm (mean 44.3 \times 26.0 μm), spore wall light chestnut brown, densely covered by fine or slightly coarser, flat or hemispherical warts, ca. 1.5–2.5 μm thick, apically hardly thickened, germ pore of distal cell ca. 1/3 (–1/2) shifted towards septum, in the proximal cell ca. 1/2 (–2/3) towards hilum, with broad, flat, yellowish brown caps, pedicel thick-walled, subhyaline, yellowish brown close to hilum, swelling strongly ca. 18–30 μm from hilum to assume a more or less tuberous shape.

On the leaves of *Lycium* sp.

Holotype (PREM 60148). South Africa, Western Cape Prov., Nieuwoudtville, gravel road to Louriesfontein, trails at ‘Windmill Museum’, 31°19′10.5″S, 19°07′03.0″E, alt. ca. 670 m a.s.l., on non-flowering *Lycium* sp., leg. A. Rössel, E. Uhlmann and R. Berndt, 11 Oct 2005 (RSA 137). Isotype Z+ZT (ZT Myc 1269).

In *P. lycii* var. *lycii*, the vast majority of the urediniospores has (sub)equatorial germ pores. The new variety differs from the nominal one by shorter and broader urediniospores whose germ pores are constantly bizonate and generally more numerous within a ring. I would not have hesitated to describe it as a new species had not a study of the type of *P. lycii* var. *lycii* also revealed a few urediniospores with bizonate pores (Berndt and Uhlmann 2006). As it cannot be ruled out at present that *P. lycii* is variable with regard to the position of the germ pores only variety rank is conferred to the form with bizonate pores. For a key of African species of *Puccinia* on *Lycium* see Berndt and Uhlmann (2006).

Zygophyllaceae (Zygophylloideae)

Mennicken et al. (2005) surveyed rust fungi on African Zygophylloideae and distinguished four species of *Uromyces* on *Zygophyllum*. We found an additional *Uromyces* species that is described as *U. eclipsis*.

Uromyces eclipsis R. Berndt & A. R. Wood, sp. nov.

Figure 10.

Etymology: Designating the appearance of the teliospores.

Spermogonia et **uredinia** non visa. **Aecia** aggregata vel singularia amphigena in foliis ramulorum scoparie deformium vel non deformium, inaperta cum peridio anguste conico, aperta cum peridio cylindrico ad anguste

cyathiformi eburneo ca. 0.4–0.7 mm longo, margine integro vel sublacerato, cellulae peridii intus verrucis sate delicatis ad grossis confluentibus obsitae, extus conspicue striatae striis delicatis densis longitudinalibus subundulatis; **aeciosporae** subglobosae ad late ellipsoideae plerumque subangulosae, 25–31 \times 22–26.5 μm (28.0 \times 24.0 μm), pariete hyalino ca. 1–1.5 μm crasso dense delicateque verruculoso. **Telia** in foliis sparsa et amphigena subepidermalia atro-brunnea vel atra pulverulentia aeciis saepe consociata; **teliosporae** ovoideae late ellipsoideae vel subglobosae saepe leniter asymmetricae, 37–47 \times 32–41 μm (pro holotypo 40.0 \times 35.7 μm , pro paratypo 41.5 \times 37.0 μm), pariete distincte bilaminato, lamina interiore castanea ca. (3)4–7 μm crassa aequaliter remoteque echinulata spinis grossis ca. 3–5 μm longis et inter se 5–8 μm distantibus, lamina exteriori straminea vel ochracea ca. 1.5–2.5(4) μm crassa, poro germinationis apicali vel subapicali papilla indistincta praedito vel sine illa, pedicello fragili brevi.


In *Zygophylli morgsanae* L.

Spermogonia and **uredinia** not seen. **Aecia** amphigenous and densely aggregated on leaves of witches’ brooms or scattered and amphigenous on leaves of apparently unaltered branches, sometimes on small, weakly hypertrophied areas of leaves, closed with narrowly conical peridia, opened peridia cylindrical to narrowly cup-shaped, ca. 0.4–0.7 mm long, with margin entire or weakly lacerated, ivory coloured; peridial cells verrucose on inner side with rather delicate to coarse warts which become confluent, on the outside with a conspicuous pattern of fine, longitudinal, often somewhat wavy, densely arranged, shorter or longer striae; **aeciospores** subglobose to broadly ellipsoid, generally subpolyhedral or slightly deformed by mutual pressure, 25–31 \times 22–26.5 μm (mean 28.0 \times 24.0 μm), spore wall hyaline, ca. 1–1.5 μm thick (incl. ornament), finely and densely verruculose, with some coarser warts interspersed. **Telia** scattered and amphigenous on leaves, subepidermal, blackish brown to black, pulverulent, often closely associated with aecia; **teliospores** ovoid, broadly ellipsoid or subglobose, often slightly asymmetrical, 37–47 \times 32–41 μm (mean for holotype 40.0 \times 35.7 μm , for paratype 41.5 \times 37.0 μm), spore wall distinctly two-layered, with an inner, chestnut brown layer ca. (3)4–7 μm thick and a straw-coloured to ochraceous outer layer ca. 1.5–2.5(4) μm thick, inner wall layer evenly and distantly covered by coarse spines ca. 3–5 μm long and 5–8 μm apart, germ pore apical or subapical, visible as a conical or trapezoid pit, sometimes indistinct, sometimes with an ill-defined and inconspicuous papilla, pedicels frail, normally breaking close to the hilum.

On *Zygophyllum morgsana* L.

Holotype (ZT Myc 643): South Africa, Western Cape Prov., Plettenberg Bay, Keurboomstrand, on *Z. morgsana*, leg. H. Schuepp, 22 Oct 1959 (sub *U. trollipi*, I [obsolete] /III).

Fig. 10 *Uromyces eclipsis*. **a** Teliospores (holotype). **b** Aeciospores in optical section (RSA 378). **c** Aeciospores, focus on spore surface (RSA 378). **d** Peridial cells, inner tangential wall with moderately coarse warts (RSA 378). **e** Peridial cells, outer tangential wall with finely striated surface (RSA 378). Bars=10 μ m



Paratype (PREM 60150): South Africa, Western Cape Prov., Groenvlei E of Lake Pleasant, at Goukamma Nature Reserve, scrub along 'Groenvlei trail' at southern side of Groenvlei, 34°02'01.3"S, 22°51'38.1"E, alt. ca. 16 m a.s.l.,

on *Z. morgsana*, leg. A. Rössel, E. Uhlmann and R. Berndt, 26 Oct 2005 (RSA 378, I/III).

Additional material studied by A.R. Wood: South Africa, Western Cape Prov., Koppie Alleen, De Hoop Nature

Table 3 Comparison between uredinio- and teliospores of *Uredo zygophylli* Jacz. (= *U. zygophyllina* Sacc.), *U. zygophylli* Henn. and *Uromyces dinteri* Mennicken, Maier & Oberw

Species (specimen)	Urediniospores (means)	Teliospores (means)
<i>U. zygophylli</i> Jacz. = <i>U. zygophyllina</i> Sacc. (ZT Myc 644)	25–31×22–27 μ m (28.8×24.4 μ m), germ pores 5–8, most often 6, with caps (own observations). 24–36×20–27 μ m (Mennicken et al. 2005)	27–36×27–33 μ m (31.5×29.2 μ m), excl. outer wall layer (own measurements)
<i>U. zygophylli</i> Henn. (holotype, B)	30–36×24–29 μ m (31.0×25.8 μ m), germ pores 6–8, most often 6, with caps (own observations)	Only three teliospores encountered
<i>Urom. dinteri</i> Mennicken, Maier & Oberw.	31–40(42)×(22)24–36 μ m, germ pores 6–9, without caps (Mennicken et al. 2005)	(31)34–44×30–42 μ m, excl. outer wall layer (Mennicken et al. 2005)

Reserve, E of Bredasdorp, 34°28'S, 20°31'E, on *Z. morgsana*, leg. A.R. Wood, 12 Oct 1996 (Wood no. 31, I/III). South Africa, Western Cape Prov., near Homestead, De Hoop Nature Reserve, E of Bredasdorp, 34°27'S, 20°24'E, on *Z. morgsana*, leg. A.R. Wood, 29 Sep 2000 (Wood no. 332, I/III). South Africa, Northern Cape Prov., Niewoudtville Nature Reserve, W of Niewoudtville, 31°21'S, 19°08'E, on *Z. morgsana*, leg. A.R. Wood, 25 Sep 2004 (Wood no. 594, I/III).

Uromyces eclipsis differs from *U. dinteri* Mennicken, Maier & Oberw., *U. paulshoekensis* Mennicken, Maier & Oberw. and *U. trollipi* Kalchbr. & McOwan by its demicyclic life cycle comprising an aecial stage while the latter species are only known with uredinia and telia. In addition, *U. paulshoekensis* and *U. trollipi* have smaller teliospores than *U. eclipsis*. The teliospores of *U. dinteri* are very similar to *U. eclipsis*. *U. namaqualandus* Mennicken, Maier & Oberw. is only known with teliospores which are narrower than in the present species and have considerably thicker inner and outer wall layers.

Observations made by Alan R. Wood (personal communication) in specimens from De Hoop Nature Reserve indicate that the witches' brooms induced by *U. eclipsis* are perennial as new growth bearing rust sori restarts on them after defoliation during the driest season of the year.

The status of Uredo zygophylli Henn. and *U. zygophylli* Jacz. (= *U. zygophyllina* Sacc.)

Material examined: *Uredo zygophylli* Henn. Egypt, on *Zygophyllum decumbens* Delile, leg. Ehrenberg, May of year? (probably between 1820 and 1825, when C.G. Ehrenberg travelled through North Africa and western Asia). (Holotype, B 700007286, II/[III]). *U. zygophyllina* Jacz. Algeria, Biskra, on *Z. cornutum* Coss. Gift from Jaczewski to Ed. Fischer (ZT Myc 644). *U. zygophyllina* Sacc. Algeria, Biskra, on calcareous hills near Hamman Salahin, on *Z. cornutum*, leg. G. Schweinfurth, Apr 1901 (Sydow, Uredineen no. 2296. ZT Myc 645, II).

Uredo zygophylli was described by Hennings (1893) from Egypt. Later in the same year *U. zygophylli* Jacz. was described from Algeria. Saccardo (1895) stated that *U. zygophylli* Jacz. was probably different from *U. zygophylli* Henn. ("videtur diversa ab *U. zygophylli* Henn.") and proposed for it the new name *U. zygophyllina*. Sydow and Sydow (1924) regarded *U. zygophylli* Jacz. and *U. zygophyllina* as synonyms of *U. zygophylli* Henn. Mennicken et al. (2005) followed the view of Saccardo and kept *U. zygophyllina* apart from *U. zygophylli* Henn. which they considered to be the uredinial state of *Urom. dinteri*.

I cannot find important differences between *U. zygophylli* Henn. and *U. zygophylli* Jacz. though the urediniospores are slightly smaller in the latter (Table 3).

Differences occur, however, between *U. zygophylli* Henn. and *Urom. dinteri* whose urediniospores have germ pores "without papillae" (Mennicken et al. 2005) while in *U. zygophylli* the pores are covered by conspicuous bullate caps. Therefore, I agree with Sydow and Sydow (1924) that *U. zygophylli* Henn. and *U. zygophylli* Jacz. represent the same rust species for which *U. zygophyllina* is another synonym.

In the specimen of *U. zygophylli* kept in Z+ZT (ZT Myc 644, sub *U. zygophyllina*), teliospores of *Uromyces* were discovered which were different from *Urom. dinteri* (Table 3). They were smaller and had thinner wall layers (inner layer 4–6 µm vs 5–8(10) µm, outer layer 3–4 vs up to 5 µm). This and the observable differences between the urediniospores indicate that *U. zygophylli* Henn. belongs to a teleomorphic species different from *Urom. dinteri*. As teliospores of *Uromyces* species on *Zygophyllum* appear to be variable and as some of the described species are difficult to distinguish a new species will not be described in ZT Myc 644 without having studied the variability within this group on more material.

Acknowledgements Field work was carried out in collaboration with Anja Rössel and Elisabeth Uhlmann whose companionship and help are gratefully acknowledged. The field stays were funded by the German Ministry of Education and Research (BMBF) in the course of the BIOTA research project (subproject S03b). The Western Cape Nature Conservation Board in South Africa is thanked for issuing collecting and export permits, and the curators and directors of B, PREM, PUR, TUB, VPRI for the loan of specimens. Last but not least I thank two anonymous reviewers for helpful comments and corrections.

References

- Berndt R (2008a) The rust mycobiota of southern Africa: species richness, composition and affinities. *Mycol Res* 112:463–471, doi:10.1016/j.mycres.2007.05.005
- Berndt R (2008b) The rust fungi (Uredinales) on ferns in South Africa. *Mycol Prog* 7:7–19, doi:10.1007/s11557-007-0548-7
- Berndt R, Uhlmann E (2006) New species, reports, observations and taxonomical changes of southern African rust fungi. *Mycol Prog* 5:154–177, doi:10.1007/s11557-006-0510-0
- Cowling RM, Richardson DM, Pierce SM (eds) (1997) *Vegetation of Southern Africa*. Cambridge University Press, Cambridge
- Cunningham GH (1931) *The rust fungi of New Zealand*. John McIndoe, Dunedin, New Zealand
- Doidge EM (1927) A preliminary study of the South African rust fungi. *Bothalia* 2:1–228
- Doidge EM (1948) South African rust fungi, part V. *Bothalia* 4:895–918
- Doidge EM (1950) The South African fungi and lichens to the end of 1945. *Bothalia* 5:1–1094
- Eboh D (1984) A taxonomic survey of Nigerian rust fungi: Uredinales Nigerianensis–III. *Mycologia* 76:179–189, doi:10.2307/3793093
- Gäumann E (1959) *Die Rostpilze Mitteleuropas. Beiträge zur Kryptogamenflora der Schweiz* 12. Büchler, Bern
- Gjærøum HB (1988) Rust fungi (Uredinales) on Poaceae, mainly from Africa. *Mycotaxon* 31:351–378

- Govaerts R, Ruhsam M, Andersson L, Robbrecht E, Bridson D, Davis A, Schanzer I, Sonké B (without date). World Checklist of Rubiaceae. The Board of Trustees of the Royal Botanic Gardens, Kew. Published on the Internet; <http://www.kew.org/wcsp/rubiaceae/> (accessed 17 Nov 2008)
- Hennings P (1893) Fungi aethiopico-arabici I. Bull Herbarii Boissier 1 (3):97–122
- Holm L (1973) Notes on some East African rusts. Sven Bot Tidskr 67:253–256
- Holmgren PK, Holmgren NH, Barnett LC (1990) Index Herbariorum, Part I, 8 edn. The New York Botanical Garden, New York
- Khan SA, Razafimandimbison SG, Bremer B, Liede-Schumann S (2008) Phylogeny and biogeography of the African genus *Virectaria* Bremek. (Sabiceae s. l., Ixoroideae, Rubiaceae). Plant Syst Evol 275:43–58, doi:10.1007/s00606-008-0053-8
- Klebahn H (1914) Kryptogamenflora der Mark Brandenburg, Vol. Va. Pilze III (Uredineen). Gebr. Bornträger, Leipzig
- Laundon GF (1963a) Rust fungi I. On Acanthaceae. Mycol Pap 89:1–89
- Laundon GF (1963b) Rust fungi II. On Aceraceae, Actinidiaceae, Adoxaceae and Aizoaceae. Mycol Pap 91:1–17
- Linder HP (2003) The radiation of the Cape flora, southern Africa. Biol Rev Camb Philos Soc 78:597–638, doi:10.1017/S1464793103006171
- McAlpine D (1906) The rusts of Australia. Dept. of Agriculture, Victoria. The Government Printer, Melbourne
- Mennicken M, Oberwinkler F (2004) A contribution to the rust flora (Uredinales) of southern Africa, with an emphasis on South Africa. Mycotaxon 90:1–28
- Mennicken M, Maier W, Oberwinkler F (2005) A contribution to the rust flora (Uredinales) on Zygothylloideae (Zygothylloaceae) in Africa. Mycotaxon 91:39–48
- Morris MJ, Wingfield MJ, Walker J (1988) First record of a rust on *Acacia mearnsii* in southern Africa. Trans Br Mycol Soc 90:324–327
- Moss EH (1926) The uredo stage of the Pucciniastreae. Ann Bot (Lond) 40:813–849
- Moss EH (1928) The uredinia of *Cronartium comandrae* and *Melampsora medusae*. Mycologia 20:36–40, doi:10.2307/3753804
- Puff C (1986) Rubiaceae, Rubioideae (second part): Paederieae, Anthospermeae, Rubieae. In: Leistner OA (ed) Flora of southern Africa vol. 31, pt. 1, pp 1–79
- Saccardo PA (1895) Sylloge Fungorum vol. XI, suppl. univ. pars III, Padova
- Sydow P, Sydow H (1904) Monographia Uredinearum, Vol. I. *Puccinia*. Gebr. Bornträger, Leipzig
- Sydow P, Sydow H (1924) Monographia Uredinearum, Vol. IV. *Uredineae Imperfectae*. Gebr. Bornträger, Leipzig
- The International Plant Names Index (2004) Published on the Internet <http://www.ipni.org> (accessed 18 Jan 2008)
- Van Reenen M (1995) An annotated list of Urediniomycetes (rust fungi) from South Africa 1: Melampsoraceae and Pucciniaceae, excluding *Puccinia* and *Uromyces*. Bothalia 25:173–181
- Viégas AP (1960) Uma ferrugem do saguaragi. Bragantia 19:101–103
- Viennot-Bourgin G (1970) Une nouvelle espèce d'*Uredo* de l'île de la Nouvelle-Amsterdam. Rev Mycologie 34:340–342
- Wood AR (2006) New and interesting records of southern African rust fungi (Uredinales). S Afr J Bot 72:534–543, doi:10.1016/j.sajb.2006.02.002
- Wood AR (2007) Rust fungi (Uredinales) on *Grewia* species (Tiliaceae) in South Africa, with *Uredopeltis atrides* comb. nov. the new name for *Ravenelia atrides*. Mycol Prog 6:93–99, doi:10.1007/s11557-007-0528-y
- Wood AR, Crous PW (2005) Morphological and molecular characterization of *Endophyllum* species on perennial asteraceous plants in South Africa. Mycol Res 109:387–400, doi:10.1017/S0953756204002175
- Wurth T (1905) Rubiaceen bewohnende Puccinien vom Typus der *Puccinia galii*. Centralblatt für Bakteriologie (II. Abt.) 14: 209–224, 309–320
- Yen J-M (1975) Étude sur les champignons parasites du sud-est asiatique. XXIV. Les Urédinées de Formose. Rev Mycologie 39:251–267