

Apathy: a separate syndrome from depression in dementia? A critical review

Moyra Elizabeth Mortby^{1,2}, Andreas Maercker² and Simon Forstmeier²

¹Centre for Research on Aging, Health and Wellbeing, The Australian National University, Canberra, Australia, ²Division of Psychopathology and Clinical Intervention, University of Zurich, Switzerland

ABSTRACT. *Apathy and depression are the most prevalent neuropsychiatric symptoms in Alzheimer's disease and mild cognitive impairment. Despite much research on apathy and depression in dementia, the nosological position of apathy as a separate syndrome from depression remains debated. This literature review provides a critical analysis of the areas of clinical manifestation, symptomatology, assessment, prevalence and neuropathology. Evidence does not provide a clear view of the nosological position of apathy in dementia for symptoms and neuropathology. However, the ambiguity of the evidence may be attributed in large part to a lack of clarity in definition and etiology, clinical criteria and assessment overlap. Given the evidence, it is concluded that the argument in favor of apathy as a separate syndrome from depression in dementia is persuasive. Reaching a consensus on the definition and nosological position of apathy within dementia is vital to provide patients and caregivers with the support they require, increase understanding of risk factors, and enable comparisons across research and practice.*

(Aging Clin Exp Res 2012; 24: 305-316)

©2012, Editrice Kurtis

INTRODUCTION

Apathy and depression are prevalent neuropsychiatric symptoms in Alzheimer's disease (AD) and mild cognitive impairment (MCI) (1). While closely clinically related, demonstrating co-morbidities and key symptom overlap (e.g., loss of interest and motivation, reduced activity or hedonism, and lack of insight) (2, 3), they are also pathophysiologically distinct (4), and several studies indicate two separate syndromes, with differential prevalence rates, neuropsychiatric symptoms and cognitive functions (2, 4-7). The nosological position of apathy as

a separate syndrome from depression remains debated (3, 8, 9) despite extensive research. Review and research papers mostly address this issue only briefly, not providing in-depth understanding of how and why apathy and depression differ in dementia. For example, Ishii et al. (7) provided a detailed review on apathy as a common psychiatric syndrome in the elderly, but offered only a subsection on the differentiation analysis, both in support of apathy as a separate syndrome from depression and apathy as a symptom of depression. To our knowledge, only Tagariello et al. (9) provide a literature review on this specific subject. Due to its brevity, it only provides a summary of the main issues and does not provide a comprehensive evaluation. The present paper provides an in-depth analysis, examining research both in support of apathy as a separate syndrome from depression and apathy as a symptom of depression.

APATHY IN DEMENTIA

Apathy is a behavioral and personality change prominently observed in AD, and is generally defined as a loss of motivation, manifested in behaviors of diminished initiation, poor persistence, lack of interest, indifference, low social engagement, blunted emotional response, and lack of insight, not attributable to decreased consciousness levels, cognitive impairment or emotional distress (10, 11). Marin et al. (12-14) originally defined apathy as a lack of motivation (i.e., direction, intensity and persistence of goal-directed behavior) relative to the patient's previous level of functioning or standards of age and culture, as indicated either by subjective accounts or observations by others. While their (13, 14) concept of apathy centers on reduced goal-directed behavior, cognition or emotion (15), modern conceptualizations acknowledge apathy as reflecting a multitude of

Key words: Alzheimer disease, apathy, behavioral and psychological symptoms of dementia, dementia, depression.

Correspondence: Dr. Moyra Mortby, Centre for Research on Aging, Health and Wellbeing, ANU College of Medicine, Biology & Environment, The Australian National University, Building 63, Eggleston Road, Canberra ACT 0200, Australia.

E-mail: moyra.mortby@anu.edu.au

Received April 19, 2011; accepted in revised form October 6, 2011.

First published ahead of print November 16, 2011 as DOI: 10.3275/8105.

dimensions (16): a disorder of motivation, with cognitive, sensory, motor and affective subtypes (12); a disorder of interest or motivation, including lack of emotion, initiation and enthusiasm (17); a disorder of initiative, manifested in a lack of self-initiated action (affective, behavioral or cognitive) and including 'social apathy' (a disorder of the sense of self and of social awareness) (18); a disorder of motivation, with emotional blunting, lack of initiative and lack of interest (19); a disorder of intellectual curiosity, action initiation, emotion and self-awareness (20); a disorder of voluntary and goal-directed behaviors, with three subtypes of disrupted "signal" processing: emotional-affective, cognitive and auto-activation (15); and a disorder of motivation, with diminished goal-directed behavior and cognition (21). Despite these differing views on the core features of apathy – namely, whether the central feature of apathy is disturbance of motivation (12) or initiative and self-generated voluntary and purposeful behavior (15, 18) – the consensus is that motivation, interest, action, initiation and emotional reactivity are dimensions of apathy, in which the changes in motivation may relate to either internal (self-conducted behavior) or external stimulation, and that the lack of motivation is central to the disorder (16). A correlation has been found between apathy and increased cognitive impairment, as well as lower cognitive test performance (22, 23), higher conversion rates from MCI to AD (24), depression (25, 26), and increased impairment in functional activities (10, 27). Notably, while Robert et al. (28) found no association between apathy and executive dysfunction, recent studies demonstrate apathy to be linked – especially in the early stages of cognitive dysfunction (i.e., MCI) – with executive dysfunction and especially worse performance in verbal memory, naming, set shifting and verbal fluency (29-31).

Apathy is an important indicator of early AD diagnosis, strongly correlated with more impairments of daily living activities than normally associated with cognitive status, increased dependency on caregivers to initiate activities and provide support and management (e.g. initiation of activities for which the patient is actually still capable of performing independently), reduced quality of life, rapid progression of cognitive degeneration and various psychobehavioral disturbances, which all contribute to heightened caregiver distress and burden (5, 10, 11, 27, 32-36). Assessing apathy in AD is complicated, as loss of motivation must be differentiated from loss of ability (11). This is particularly difficult in the cognitively impaired, as apathy is generally defined as a reduction in behavior in comparison to the patient's pre-morbid state (11). Cognitive function decline complicates such an assessment, as patients face increased difficulty in the organization and conduct of complex behaviors, reduced behavior resulting from impaired motivational abilities, reduced task initiation, and increased

frustration and confusion with hobbies (11, 13, 37). Although apathy is a behavioral sign of cognitive dysfunction in which impairments negatively affect the initiation, planning, and problem-solving of successful task performance (11, 38-40), Marin and Wilkosz (41) observed apathetic individuals generally to be 'able to initiate and sustain behavior, describe their plans, goals and interests, and react emotionally to significant events and experiences' despite diminished motivation associated with apathy (21, pp. 1088-1089). Apathy must therefore be distinguished from cognitive decline, as the dynamic interaction between apathy and cognitive impairment may result in AD patients demonstrating less interest in regular activities as they lose the ability to engage in cognitively complex behaviors and become less likely to initiate regular activities in which they are unsure of the steps required (11). Consequently, apathetic symptoms must be considered when increased functional impairments and reliance on caregiver initiation become more evident than expected for a given level of cognitive impairment (11).

The need for applicable diagnostic criteria for apathy in dementia was originally addressed by Starkstein et al. (10, 21) and recently revised by Robert et al. (16). Starkstein et al. (10, 21) operationalized the definition of apathy of Marin et al. (12) into applicable diagnostic criteria which require: 1) the presence during most of the day for a minimum of 4 weeks of at least one symptom of diminished goal-directed behavior (e.g., lack of effort and initiative, or energy to perform everyday activities; dependency on prompts from others to structure everyday activities); diminished goal-directed cognition (e.g., lack of interest in learning new things or in new experiences; lack of plans and goals, lack of concern about one's own health, functional status or personal problems) or diminished concomitants of goal-directed behavior (e.g., unchanging or flat affect; lack of emotional responsiveness to positive or negative events and restricted responses to important life events); 2) clinically significant distress or impairments in social, occupational or other important areas of functioning caused by apathetic symptoms; and 3) symptoms are not due to a diminished level of consciousness or direct physiological substance effects (e.g., drug abuse, medication) (10, 21). Thus, diminished motivation, initiative and interest, and emotional blunting are at the core of Starkstein et al.'s definition of apathy (16). These criteria have recently been revised into semi-operationalized criteria by Robert et al. (16) based on the synthesis of current concepts of apathy applicable to both research and practice. The revised criteria follow Starkstein et al. (10, 21) structure, and require: 1) loss of or diminished motivation compared with previous functioning levels, not consistent with age or culture, and reported either by the patient or observers; 2) the presence of

impairments (minimum of one symptom) in at least two of the apathetic dimensions of reduced goal-directed behavior, goal-directed cognitive activity, and emotions in which motivational changes are assessed by responsiveness to either internal (self-conducted behavior) or external stimulation (16); 3) functional impairments attributable to apathy; and 4) the exclusion of symptoms and conditions mimicking apathy (42). Future research is required to assess not only the reliability and validity of these criteria, but also the proposed distinction between changes in motivation relating to internal and external stimulation and the impact which such differentiation may have.

DEPRESSION IN DEMENTIA

Depression in AD is characteristically expressed by the high frequency of motivational disturbances (e.g., fatigue, psychomotor slowing and apathy), symptoms of social isolation, withdrawal, irritability or emotional distress, and vegetative symptoms of diminished interest, psychomotor retardation, fatigue, hypersomnia and lack of insight (2, 13, 43-46). While these vegetative symptoms are also key symptoms of apathy (2, 13, 44), depression differs from apathy according to key clinical dysphoric symptoms of sadness, feelings of guilt, self-criticism, helplessness and hopelessness (44), and concurrent reports in the literature propose dysphoria and loss of interest as the most common symptoms of depression in AD (46, 47). However, these findings may be tautological, due to assessment requirements (46, 47). Notably, depressive AD patients demonstrate significant impairments in quality of life, activities of daily living and executive function, although no differences are observed for attention, language, memory and visuospatial functions compared with patients with no changes in mood (48-53). Depression in AD has far-reaching consequences. For AD patients, depression is associated with increased likelihood of physical aggression, being discharged from an assisted living facility and earlier entry into a nursing home, and higher mortality and suicide; for caregivers, it is associated with higher personal depression and burden (44, 52, 54-60). Risk factors for depression in AD include a family history of mood disorders (affective disorders) in first-degree relatives, prior personal depressive history, female gender, and younger age at AD onset (50, 56, 61-65).

Assessing depression in AD is complicated, as symptoms of psychomotor slowing, emotional lability, crying spells, insomnia, weight loss, inability to verbalize affective states and pessimism are prevalent in both depressed and non-depressed AD patients (44, 66). Standardized diagnostic criteria such as the Diagnostic and Statistical Manual of Mental Disorders (DSM) or the International Classification of Diseases (ICD) (67) have been implemented to assess depression in AD. For example, in the DSM-IV

criteria for depression, a principal symptom is loss of interest or pleasure in activities as opposed to depressed mood itself (68). However, in the demented, such a loss of interest or pleasure may reflect more a loss of motivation or ability, rather than a change in affect or depression (5). This symptom overlap (e.g., lack of interest, anxiety, lack of energy, concentration difficulty, agitation or psychomotor retardation, sleep and eating disorders) has not only resulted in blurring between AD and depressive symptomology (46, 47, 60, 67, 69), but also resulted in apathetic individuals meeting the diagnostic criteria for major depression even in the absence of dysphoric symptoms (5, 70, 71).

A further limitation of the use of non-dementia specific assessments is their reliance on ability to report subjective depressive symptoms verbally (e.g., mood changes, loss of interest, hopelessness), a process commonly impaired in AD patients (69). Non-dementia specific diagnostic criteria for depression (e.g., DSM, ICD), developed for use in younger, cognitively unimpaired individuals, lack sensitivity to differentiate accurately between symptoms of cognitive decline and depression, and have decreased validity in AD samples (46, 47, 60, 67, 69, 72). This may be explained by differences in clinical manifestations of late life mood disturbance compared with younger life, and the denial by the depressed elderly of sad mood, who rather report lack of feeling or emotion and acknowledge a loss of interest and pleasure in activities (i.e., the depressive elderly report fewer affective symptoms – known as ‘depression without sadness’) (46, 47, 60, 67, 69, 72-74).

Pertinently, depression and dementia overlap in the under-reporting of depressive symptoms by dementia sufferers and increased reliance on caregiver and observational reports (44, 75). Due to aphasia, many dementia patients lack the ability to express their distress coherently – a further complication when diagnosing depression in AD (44). In addition, the tendency of depressive AD patients to deny the presence of depressed moods and report instead a lack of feeling or emotion, or loss of interest and pleasure in activities, hampers diagnosis with non-dementia specific assessments of depression in the elderly, with and without cognitive impairment (44). This, and the observed difference in depressive symptoms between the elderly with and without dementia, led Olin et al. (46, 69) to propose depression in AD as an atypical form, in which motivational symptoms and delusions are experienced more frequently than by non-cognitively impaired patients (67, 76, 77).

Such difficulties in assessment led Olin et al. (46, 69) to develop the National Institute of Mental Health Provisional Diagnostic Criteria for Depression of Alzheimer’s Disease (NIMH-dAD), based on criteria for major depressive episodes (78), but broadened to include those AD patients who experience ‘clinically significant affective disturbances but do not meet the standardized diagnostic

criteria for depression' (67, p. 590). The NIMH-dAD diagnostic criteria have been modified to be independent of verbal expression (i.e., subjective reporting) and not confounded by cognitive AD symptomology, and are aimed at improving understanding of the nosology, etiology and treatment of AD-related depression (46, 69). The NIMH-dAD criteria require 1) the presence of three (or more) symptoms of clinically significant depressed mood (e.g., depressed, sad, hopeless, discouraged, tearful); decreased positive affect or pleasure as a response to social contact and usual activities; social isolation or withdrawal; appetite disruption; sleep disruption; psychomotor changes (e.g., agitation, retardation); irritability; fatigue or loss of energy; feelings of worthlessness, hopelessness, or excessive or inappropriate guilt; or recurrent thoughts of death or suicidal ideation (plan or attempt); 2) that all criteria for Dementia of Alzheimer Type of the DSM-IV-TR are met; 3) that significant distress or disruption to functioning is caused by the symptoms; 4) that symptoms are not exclusive to the course of delirium, nor the direct result of physiological substance effects (e.g., drug abuse, medication), or that the symptoms may be better explained by other conditions (e.g., major depressive disorder, bipolar disorder, bereavement, schizophrenia, schizoaffective disorder, AD psychosis, anxiety disorder, or substance-related disorder) (69).

The NIMH-dAD proposes that AD-related depression has heterogeneous etiology characterized as one of four subtypes: 1) emotional reaction to cognitive deficits in AD (e.g., adjustment disorder with depressed mood), in which the depressive state may be a reaction to the diagnosis of AD and/or associated with the awareness of loss and disability, or a reaction of certain personality types vulnerable to depressive symptomology in response to negative life events; 2) recurrence of early and midlife major and minor depressive disorders is a risk factor for later life dementia, and may share its etiology with early or midlife depression; 3) vascular diseases associated with AD causing depressive symptoms (so-called vascular depression), in which occurrence of subcortical lesions in depressed elderly patients with vascular disease indicate that vascular depression results from 'critical lesions or an accumulation of lesions leading to disruption of frontostriatal pathways or their modulating systems'; 4) the neurodegenerative process of AD causes depressive symptoms (e.g., mood disorder due to general medical condition), with pathophysiological overlap between AD and the depressive syndrome in AD, in which the 'neurodegenerative process may directly contribute to the development of depressive symptoms' (44, pp. 357-358). This set of criteria has been shown to be particularly sensitive, with higher rates of depression reported in both retrospective (27.4%) (67) and prospective (44%) (43) implementations of the NIMH-dAD compared with other sets of criteria (e.g., DSM-IV, CAMDEX and ICD-10). In support of

Olin et al.'s (46) proposition, Teng et al. (43) demonstrated that an interview using NIMH-dAD criteria more effectively distinguishes significant depressive symptoms in AD patients. Notably, while the symptoms of psychomotor changes, fatigue and guilt/worthlessness were found to predict depression, the newly included symptoms of social isolation/withdrawal and irritability were found not to do so (43). This led the authors to conclude that these symptoms may be more specific for depressive manifestation in AD rather than influence the rate of depression (43). Nonetheless, the NIMH-dAD criteria have face validity (67), and Rosenberg et al. (72) have demonstrated their clinical application.

SYMPTOMATIC OVERLAP OR ASSESSMENT OVERLAP?

Apathy and depression are frequently associated in AD, due to joint key symptoms of diminished motivation and interest, vegetative symptoms of psychomotor retardation, fatigue and hypersomnia, lack of insight and pessimism, diminished ability to concentrate, and weight loss (9, 11, 60). This symptomatic overlap often results in the misinterpretation of apathy as depression, even in the absence of depressed mood or signs of dysphoria according to the DSM criteria (5, 11, 60, 68, 71). According to the DSM-IV criteria for depression, one main symptom is loss of interest or pleasure in activities, as opposed to depressed mood itself (68). However, in the demented, this loss of interest or pleasure may reflect a loss of motivation or ability, rather than a change in affect or depression (5). This both complicates its assessment (44, 47, 60) and results in apathetic individuals meeting the diagnostic criteria for major depression, even in the absence of dysphoric symptoms (5, 11, 70, 71).

The observed overlap may be explained by methodological variations of assessments relating to symptoms common to both syndromes (e.g., closely related subsets of items, such as diminished interest, psychomotor retardation, lack of energy, loss of insight; or dysphoric items, such as depressed mood, guilt, hopelessness and vegetative symptoms) (2, 11, 70, 79). Research has demonstrated high correlations between apathy and depression scores for items corresponding to symptoms commonly observed in the syndrome of apathy (2, 12, 26, 80), as well as categorization of patients into cohorts of "pure apathy", "pure depression" or "apathy and depression" (3, 10, 45, 81). This has led to the conclusion that apathy and depression have divergent natural histories and can be differentiated (7, 82).

Forsell et al. (83, 84) described depressive symptoms in the elderly as falling into two categories: mood problems (symptoms of dysphoria, feelings of guilt and suicidal ideation) and motivational problems (symptoms of lack of interest, low energy and psychomotor slowing). Depression may be distinguished from apathy (characterized

by lack of emotional responsiveness) according to dysphoric symptoms of sadness, feelings of guilt, self-criticism, helplessness and hopelessness (11, 85-87). The observed loss of interest in depression may rather reflect feelings of despair, pessimism and hopelessness which, combined with dysphoria in depression, differentiate depression from apathy (11, 85). Conversely, apathy is distinguishable from depression according to lack of emotional responsiveness, emotional indifference, 'lack or lessening of ability to initiate in multiple domains (motor, gait, cognitive)' and a lack of concern – not all of which are exclusively observed in depressed individuals (11, 13, 71, p. 16, 86, 87).

Differences are also observed for neuropsychiatric symptoms, apathy being associated with disinhibition and aberrant motor behavior, and depression with anxiety, agitation, irritability and hallucinations (9, 45). The occurrence of negative mood and dysphoric symptoms in depression, but not in apathy, provides a further difference (5). Various studies have demonstrated differences between apathy and depression in AD for associated cognitive deficits (11, 30, 88). Research by Kuzis et al. (30) compared only apathetic, only depressive, and co-morbid apathetic and depressive early AD patients. In contrast with only depressive patients, only apathetic patients showed specific cognitive deficits in naming performance, word list learning, verbal fluency and set-shifting (30). The co-morbid apathetic and depressive group had worse impairments in abstract reasoning ability than those with only apathy, while no association with cognitive dysfunction was found for the only depressive group (11, 30).

Although the concept of apathy as a separate syndrome from depression has received much valid support, there are opposing arguments. First, for Marin (13), the diagnosis of apathy should not be made in the context of diminished levels of consciousness, moderate or severe cognitive deficits, or marked emotional distress, as apathy may be an intrinsic symptom of dementia (21). Although this point is useful in identifying a 'pure' apathy syndrome 'uncontaminated by changes in cognitive or affective domains', conceptual and empirical evidence argues against such an approach (21, p. 1089). One conceptual issue is that a syndrome is defined as a constellation of symptoms without reference to a specific etiology, and it is therefore not clear why apathy should be 'considered a syndrome in some contexts (e.g., stroke) but not in others (e.g., dementia)' (21, p. 1089). A second opposing argument is that some psychopathological syndromes (e.g., apathy and depression) co-occur in some neurodegenerative diseases (21, 26). In response, while the apathetic syndrome is observed most frequently in 'individuals with neurological disorders and some degree of cognitive impairment and depression' (e.g., AD, stroke, or Parkinson's disease) (21, p. 1089, 81, 89), this may be due to the inability of assessments to

differentiate between symptoms of apathy, cognitive impairment and depression, as is the case in Parkinson's disease, in which there is an overlap between depression, dementia and apathy (90-92). A third opposing argument is that both DSM-IV and ICD-10 allow for the diagnosis of mild or moderate depression in the absence of depressed mood, while requiring loss of interest or anhedonia ('inability to experience pleasure, as manifested in facial expression, speech, behaviour, lifestyle and the individual's account of personal experience'), or a decrease in energy (21, p. 1090, 93). Starkstein and Leentjens (21, p. 1089) rightly question whether: 1) such a definition adequately explains the high co-occurrence of apathy and depression, 2) such a reliable separation of non-dysphoric depression from apathy may occur, and 3) 'lack of interest and/or anhedonia are core symptoms of diminished motivation', as Starkstein et al.'s (10, 21) apathy diagnostic criteria require diminished goal-directed behavior for apathy (assessed by 'lack of interest'). Accordingly, the above argument for apathy as a symptom of depression must be questioned. On balance, the weight of literary evidence points toward apathy as a separate and distinct syndrome from depression, and this view is further supported by findings on the prevalence of apathy and depression in dementia.

DOES PREVALENCE INDICATE SIMILARITY OR DIFFERENCE?

Prevalence and progression rates further support apathy as a separate syndrome from depression. Apathy is a prominent feature of AD-related behavioral and personality changes (11, 26, 27, 32, 70, 94, 95), appearing early in the course of the disease (reports ranging between 11 and 39% of individuals with MCI) and increasing in prevalence with illness progression until it is universal amongst the severely cognitively impaired (up to 92% of AD patients) (11, 24, 35, 50, 70, 71, 80, 96, 97). Conversely, depression precedes the onset of AD (5, 10, 50, 70, 98, 99) and is often an initial symptom in AD, appearing in mild to moderate stages (including MCI) and becoming less prevalent in more severe stages (46, 56, 100, 101). Prevalence rates of depression (major and minor) in AD reportedly range between 30 and 50% (with more extreme studies reporting rates between 1 and 90%) (44, 46). These inconsistencies may be explained by the large variability in definition, assessment and clinical manifestation in research (44, 46). Notably, as AD progresses, the expression of depressive syndromes changes, resulting in under-recognition of late-life depression due to clinical and nosological ambiguities (83, 102). However, the declining prevalence of depression in more advanced AD stages may be the consequence of assessment difficulties, due to advanced cognitive decline (46, 56).

Apathy also occurs without depression in AD, sug-

gesting that the former is a distinct and distinguishable syndrome from depression (3, 11, 21, 45). Marin et al. (94) demonstrated that 55% of AD patients had high levels of apathy and low levels of depression, whereas 47% of older adults with major depression had low levels of apathy (11). This suggests differential patterns of apathy and depression in AD patients and patients with major depression, 'despite a positive correlation between apathy and depression scores in both groups' (11, p. 1704). These findings of no concomitant depression in AD patients supports apathy as a separate neuropsychiatric syndrome from depression.

Other research demonstrates co-occurrence of apathy and depression varying across neurological conditions, suggesting that these are two neuroanatomically distinct entities (2, 45). Starkstein et al. (26) supported this concept in their longitudinal examination of the association between apathy and depression. Their findings that 25% of non-depressed patients had apathy and that 50% of depressed patients also had apathy led them to conclude that depression is not a requisite for the presence of apathy in AD (26). In addition, they found a significant increase in apathy over time, in that baseline subsyndromal depression was not associated with follow-up apathy levels and baseline apathy levels were a significant predictor of increased follow-up depression (26). This last finding led them to suggest that apathy may be either an early indicator for depression or a prodromal stage of depression (26). However, in the latter case, the prevalence rates of apathy should reflect higher levels of apathy prior to depression. This is not found. While depression is most common in early stages and decreases in prevalence with cognitive decline, apathy increases with increasing cognitive impairment. Therefore, although apathy may be a possible risk factor for the development of later life depression, Starkstein et al.'s (26) proposal of apathy as a prodromal stage of depression requires further investigation. Accordingly, on the basis of current understanding of the prevalence and progression rates of apathy and depression, it is plausible to postulate that apathy and depression are separate syndromes in dementia.

In summary, apathy and depression differ according to prevalence and progression rates. While apathy occurs in early and mild stages of cognitive impairment and increases in prevalence to become universal among the severely cognitively impaired, depression is an initial symptom, already apparent in the pre-clinical stage of MCI and mild to moderate stages of impairment. In contrast to apathy, depression becomes less prevalent with increasing cognitive decline, perhaps as a consequence of methodological limitations (e.g., AD pathology, cognitive decline, and unwillingness or inability to disclose the occurrence of such neuropsychiatric symptoms). The development of more accurate and effective ways of assessing such neuropsychiatric symptoms is required when

diagnosis is not predominantly dependent on caregiver reports or clinical interpretations.

A NEUROPATHOLOGICAL EXPLANATION?

Similarities and differences in apathy and depression have not only been found in symptomatic overlaps, but also on a neuropathological level (11). Apathy is most commonly observed in combination with damage to the frontal lobes or subcortical structures connecting them (e.g., frontal lobe lesion) and impairments in these areas are associated particularly with executive dysfunction and cognitive and behavioral changes (11, 38, 71, 103, 104). In particular, apathy is 'correlated with neuronal loss, higher tangle counts and white-matter hyperintensities in areas that are thought to be essential components' of frontal subcortical circuits (11, p. 1701, 105-110). Notably, patients with moderate or severe apathy demonstrate significantly reduced blood flow in the anterior temporal, orbito-frontal, anterior cingulate and dorso-lateral prefrontal regions, compared with those with no or mild apathy (31, 50, 111, 112). These findings have resulted in the implication that 'apathy is associated with specific cognitive impairment caused by disruption of these brain areas' (29, p. 2) and, specifically, the involvement of frontal-subcortical circuits originating from frontal cortical regions, which mediate executive function and motivation in apathetic patients (30, 31, 113, 114). Thus, apathy in AD has been closely associated with neuropathological impairments in components of the anterior cingulate frontal-subcortical circuit (e.g., anterior cingulate, nucleus basalis of Meynert, hippocampus, medial frontal region) (5, 105-108).

Conversely, the neuropathology of depression in AD is primarily associated with frontal-striatal and subcortical limbic structures (locus ceruleus, substantia nigra, hippocampus and hypothalamus) (5, 9, 50, 115, 116). *Post-mortem* evidence has demonstrated a reduction in the locus ceruleus cell population, the number of substantia nigra cells, and neurochemical changes, i.e., marked reduction of norepinephrine levels in the middle frontal and temporal cortex cortical regions, increased entorhinal cortex dopamine levels, and normal frontal and temporal cortex serotonin levels – even with significant reduction of serotonin uptake sites in the temporal regions (117-121). *Post-mortem* evidence also suggests selective noradrenergic cell loss in the locus ceruleus and loss of dorsal raphe serotonergic nuclei (56, 119, 122-125). Apoptotic processes occurring as a function of AD may also contribute to the loss of aminergic nuclei, in turn affecting mood (46, 126). Interestingly, the presence of depression correlates with elevated global scores of deep white matter lesions, particularly to frontal lobe white matter (127, 128). However, although findings of functional neuroimaging studies are inconsistent, AD-related depression has been associated with reduced parietal lobe

metabolism, cerebral hypoperfusion in temporal-parietal regions, and hemispheric hypoperfusion in the dorsolateral, frontal, temporal and parietal regions (50).

In summary, while apathy in AD is associated with abnormalities of frontal subcortical circuits (specifically, the 'anterior cingulate which is thought to be responsible for motivated behavior') (29, p. 2), AD-related depression is associated with neuropathology of frontal-striatal and subcortical limbic circuits (e.g., locus ceruleus, substantia nigra, hippocampus, hypothalamus) (9).

Apathy and depression are neuropathologically similar, due to hypoperfusion or hypoactivity in frontal, parietal and temporal regions (5, 9, 11, 108, 111, 115, 116, 129-132). Landes et al. (11, p. 1704) suggested that the 'co-occurrence of apathy and depression may relate to involvement of frontal subcortical circuits in both syndromes'. Other research proposes mixed behavioral presentation of apathy and depression, due to impairment of the frontal circuitry in major depression resulting from the proximity of such subcortical circuits (i.e., a subcortical lesion simultaneously affects multiple circuits) (11, 39, 45, 104, 133, 134). However, while the involvement of frontal subcortical circuits has a long-standing tradition as a heuristic explanation of behavioral problems in neuropsychiatry, this hypothesis may lack heuristic value, as neuropathological differences have been observed in which 'AD patients with isolated motivational or mood disturbances reveal differing cortical dysfunction during early disease, confirming a different neuroanatomical basis for the emergence of apathy and depression in AD' (135, p. 418). Holthoff et al.'s (135) positron emission tomography study showed that apathetic early AD patients had significant glucose metabolism decreases in the left orbitofrontal regions, while hypometabolism in the dorsolateral prefrontal regions was associated with depression.

Other research suggests that apathy and depression differ on a neurochemical basis, apathy being associated with cholinergic deficits and depression with serotonergic deficits or a dopamine and norepinephrine imbalance (5, 50, 136-138). Symptoms of apathy in early AD are related to cholinergic input loss to the prefrontal and subcortical structures of the nucleus basalis of Meynert, whereas deepening apathy in later stages results from severe cholinergic denervation, and involvement of the prefrontal cortex and the anterior and medial temporal structures supplying adherent input to this circuit (9, 11). Thus, an interaction between neuropathological frontal brain changes and cholinergic deficiency is reflected in AD-related apathy (9).

The identification of more pronounced apathetic symptoms in AD patients exhibiting extra-pyramidal symptoms and empirical results on modulated dopaminergic neurotransmission treatment further suggest that dopamine plays a mediating role in apathetic AD patients (11, 71, 139). Conversely, serotonergic agents (e.g., se-

lective serotonin re-uptake inhibitors) may worsen apathetic symptoms but relieve depression (45, 86, 140-143). While apathy treatment may be supported by increasing dopaminergic function, serotonergic activity increase is detrimental (33). Thus, cholinomimetic agents reduce apathy but do not affect mood, so that only apathy, and not depression, is relieved by dopaminergic agents (45, 144). One explanation is that depression results from an imbalance in paralimbic neurotransmitter function, whereas apathy may be traced to a functional disconnection of the cortex from the relevant paralimbic input (9). While the treatment of apathy with cholinesterase inhibitors (e.g., tacrine) has demonstrated significant reductions in apathetic symptoms in AD, suggesting a 'cholinergic contribution to the pathophysiology of apathy in AD' (50, p. 852, 145), between 38% (146) and 60% of patients (147) do not respond to this treatment (33). This involvement of differing neurochemical pathways and, to some extent, of neuropathological structures may therefore be a further indication that apathy and depression are two separate syndromes in dementia.

In summary, apathy in AD is associated with bilateral hypoperfusion within the basal ganglia and dorsolateral prefrontal cortex (132), reduction in perfusion of anterior temporal, orbito-frontal, anterior cingulate and dorsolateral prefrontal regions (148, 149), hypometabolism in left orbitofrontal areas (135), and left anterior cingulate neurofibrillary tangle burden (105). Conversely, AD-related depression is associated with hypometabolism in the left prefrontal cortex and superior frontal cortex (135), white matter lesions (127), reduced glucose metabolism in the bilateral anterior cingulate and superior temporal cortex (132) and bilateral superior frontal and left anterior cingulate cortex (131) or parietal lobes (150).

CONCLUSIONS AND FUTURE DIRECTIONS

Although no consensus on the nosological position of apathy in dementia has been reached, apathy, despite some symptomatic overlap with depression, may be viewed as a separate syndrome from depression in dementia. This review provides an overview of the literature focusing on the areas of clinical manifestation, symptomology, assessment, prevalence, and neuropathology. We conclude that, given the evidence, the arguments in support of apathy as a separate syndrome from depression in dementia are persuasive. However, the fact that the definitions of apathy and depression overlap in terms of key symptoms, resulting in a high mis-diagnosis of apathy as depression, raises some concerns. These concerns regard treatment, as medical treatment of depression does not alleviate the negative impact of apathy. To provide sufficient support to individuals suffering from dementia and their caregivers, it is vital that apathy as a syndrome should be recognized and treated. Reaching a

consensus on the definition of apathy within dementia is essential if patients suffering from such neuropsychiatric symptoms are to receive deserved support and attention.

A consensus on the definition of apathy and its nosological position will lead to better development and implementation of diagnostic measures and criteria, more effective research and clinical practice, and better understanding of associated neuropsychiatric disorders in dementia. The identification of apathy as a separate syndrome and its management (whether pharmacologically or non-pharmacologically) requires better understanding of potential risk factors for both apathy and depression in dementia. This is important when considering the direction of future research, specifically on potentially modifiable risk factors for the development of apathy and depression in dementia. While there is ample research on risk factors of AD (151), little research has focused on risk factors for apathy and depression in dementia. Future research should consider the role of behavioral and cognitive models in this context, as these may not only provide further insight and understanding of the multi-dimensionality involved in neuropsychiatric symptoms in dementia, in which depression and/or apathy may not only be characterized as a deficit in motivation or mood, but may also reflect transient modalities to adapt to their environment (152). Future research should also consider the possibility of conversion between apathy and depression, and *vice versa*, as this may provide further understanding of risk factors. Mortby et al. (153) were the first to study midlife motivational abilities as predictors of apathy and depression in dementia, demonstrating that midlife motivational abilities moderate the causal relationship between cognitive impairment and apathy, but not depression. The findings by Mortby et al. (153) – that depression remained stable over time, regardless of severity of impairment, but that apathy increased over time in AD patients – may be interpreted as additional support to the view that apathy and depression are two separate syndromes (11). These findings may provide a new perspective to the ongoing debate on whether behavioral and psychological symptoms of dementia reflect pre-morbid personality traits (e.g., 154-156) preceding apathy and/or depression in dementia. Specifically, these findings may provide stimulation for future research into personality factors or coping styles, to determine whether apathy and/or depression are a state or trait in dementia, as this may provide additional insight into the best approaches for prevention and treatment of both.

REFERENCES

1. Lyketsos CG, Lopez O, Jones B, Fitzpatrick AL, Breitner J, DeKosky S. Prevalence of neuropsychiatric symptoms in dementia and mild cognitive impairment: results from the cardiovascular health study. *JAMA* 2002; 288: 1475-83.
2. Marin RS, Firinciogullari S, Biedrzycki RC. The sources of convergence between measures of apathy and depression. *J Affect Disord* 1993; 28: 117-24.
3. Starkstein SE, Ingram L, Garau ML, Mizrahi R. On the overlap between apathy and depression in dementia. *J Neurol Neurosurg Psychiatry* 2005; 76: 1070-4.
4. Winograd-Gurvich C, Fitzgerald PB, Georgiou-Karistianis N, Bradshaw JL, White OB. Negative symptoms: a review of schizophrenia, melancholic depression and Parkinson's disease. *Brain Res Bull* 2006; 70: 312-21.
5. Landes AM, Sperry SD, Strauss ME. Prevalence of apathy, dysphoria, and depression in relation to dementia severity in Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 2005; 17: 342-9.
6. Benoit M, Andrieu S, Lechowski L, Gillette-Guyonnet S, Robert PH, Vellas B. Apathy and depression in Alzheimer's disease are associated with functional deficit and psychotropic prescription. *Int J Geriatr Psychiatry* 2008; 23: 409-14.
7. Ishii S, Weintraub N, Mervis JR. Apathy: a common psychiatric syndrome in the elderly. *J Am Med Dir Assoc* 2009; 10: 381-93.
8. Aalten P, Verhey FR, Boziki M et al. Neuropsychiatric syndromes in dementia. Results from the European Alzheimer Disease Consortium: Part I. *Dement Geriatr Cogn Disord* 2007; 24: 457-63.
9. Tagariello P, Girardi P, Amore M. Depression and apathy in dementia: same syndrome or different constructs? A critical review. *Arch Gerontol Geriatr* 2009; 49: 246-9.
10. Starkstein SE, Petracca G, Chmerinski E, Kremer J. Syndromic validity of apathy in Alzheimer's disease. *Am J Psychiatry* 2001; 158: 872-7.
11. Landes AM, Sperry SD, Strauss ME, Geldmacher DS. Apathy in Alzheimer's disease. *J Am Geriatr Soc* 2001; 49: 1700-7.
12. Marin RS, Biedrzycki RC, Firinciogullari S. Reliability and validity of the Apathy Evaluation Scale. *Psychiatry Res* 1991; 38: 143-62.
13. Marin RS. Differential diagnosis and classification of apathy. *Am J Psychiatry* 1990; 147: 22-30.
14. Marin RS. Apathy: a neuropsychiatric syndrome. *J Neuropsychiatry Clin Neurosci* 1991; 3: 243-54.
15. Levy R, Dubois B. Apathy and the functional anatomy of the prefrontal cortex-basal ganglia circuits. *Cereb Cortex* 2006; 16: 916-28.
16. Robert P, Onyike CU, Leentjens AF et al. Proposed diagnostic criteria for apathy in Alzheimer's disease and other neuropsychiatric disorders. *Eur Psychiatry* 2009; 24: 98-104.
17. Cummings JL, Mega M, Gray K, Rosenberg-Thompson S, Carusi DA, Gornbein J. The Neuropsychiatric Inventory: comprehensive assessment of psychopathology in dementia. *Neurology* 1994; 44: 2308-14.
18. Stuss DT, Van Reekum R, Murphy KJ. Differentiation of states and causes of apathy. 1 ed. New York: Oxford University Press; 2000.
19. Robert PH, Clairet S, Benoit M, K et al. The apathy inventory: assessment of apathy and awareness in Alzheimer's disease, Parkinson's disease and mild cognitive impairment. *Int J Geriatr Psychiatry* 002; 17: 1099-105.
20. Sockeel P, Dujardin K, Devos D, Deneve C, Destee A, Defebvre L. The Lille apathy rating scale (LARS), a new instrument for detecting and quantifying apathy: validation in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2006; 77: 579-84.
21. Starkstein SE, Leentjens AFG. The nosological position of apa-

- thy in clinical practice. *J Neurol Neurosurg Psychiatry* 2008; 79: 1088-92.
22. Onyike CU, Sheppard JM, Tszchanz JT et al. Epidemiology of apathy in older adults: the Cache County Study. *Am J Geriatr Psychiatry* 2007; 15: 365-75.
 23. Turro-Garriga O, Lopez-Pousa S, Vilalta-Franch J et al. [A longitudinal study of apathy in patients with Alzheimer's disease]. *Rev Neurol* 2009; 48: 7-13.
 24. Robert PH, Berr C, Volteau M et al. Apathy in patients with mild cognitive impairment and the risk of developing dementia of Alzheimer's disease: a one-year follow-up study. *Clin Neurol Neurosurg* 2006; 108: 733-6.
 25. Lavretsky H, Lesser IM, Wohl M, Miller BL, Mehringer CM. Clinical and neuroradiologic features associated with chronicity in late-life depression. *Am J Geriatr Psychiatry* 1999; 7: 309-16.
 26. Starkstein SE, Jorge R, Mizrahi R, Robinson RG. A prospective longitudinal study of apathy in Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 2006; 77: 8-11.
 27. Freels S, Cohen D, Eisdorfer C et al. Functional status and clinical findings in patients with Alzheimer's disease. *J Gerontol* 1992; 47: M177-82.
 28. Robert PH, Berr C, Volteau M et al. Neuropsychological performance in mild cognitive impairment with and without apathy. *Dement Geriatr Cogn Disord* 2006; 21: 192-7.
 29. Drijgers RL, Verhey FR, Leentjens AF, Kohler S, Aalten P. Neuropsychological correlates of apathy in mild cognitive impairment and Alzheimer's disease: the role of executive functioning. *Int Psychogeriatrics* 2011; 28: 1327-33.
 30. Kuzis G, Sabe L, Tiberti C, Dorrego F, Starkstein SE. Neuropsychological correlates of apathy and depression in patients with dementia. *Neurology* 1999; 52: 1403-7.
 31. McPherson S, Fairbanks L, Tiken S, Cummings JL, Back-Madruga C. Apathy and executive function in Alzheimer's disease. *J Int Neuropsychol Soc* 2002; 8: 373-81.
 32. Devanand DP, Brockington CD, Moody BJ et al. Behavioral syndromes in Alzheimer's disease. *Int Psychogeriatr* 1992; 4 (Suppl 2): 161-84.
 33. Lanctôt KL, Herrmann N, Black SE et al. Apathy Associated with Alzheimer disease: use of dextroamphetamine challenge. *Am J Geriatr Psychiatry* 2008; 16: 551-7.
 34. Boyle PA, Malloy PF, Salloway S, Cahn-Weiner DA, Cohen R, Cummings JL. Executive dysfunction and apathy predict functional impairment in Alzheimer disease. *Am J Geriatr Psychiatry* 2003; 11: 214-21.
 35. Strauss ME, Sperry SD. An informant-based assessment of apathy in Alzheimer disease. *Neuropsychiatry Neuropsychol Behav Neurol* 2002; 15: 176-83.
 36. Derouesne CH. Apathy in Alzheimer's disease. *Ann Gen Hosp Psychiatry* 2003; 2 (Suppl 1): S11.
 37. Gilley DW. Behavioral and affective disturbances in Alzheimer's disease. In Parks RW, Zec RF, Wilson RS, Eds. *Neuropsychology of Alzheimer's disease and other dementias*. New York: Oxford University Press, 1993: 112-37.
 38. Fogel BS. The significance of frontal system disorders for medical practice and health policy. *J Neuropsychiatry Clin Neurosci* 1994; 6: 343-7.
 39. Duffy JD, Campbell JJ 3rd. The regional prefrontal syndromes: a theoretical and clinical overview. *J Neuropsychiatry Clin Neurosci* 1994; 6: 379-87.
 40. Royall DR. Precipitous executive dyscontrol as a cause of problem behavior in dementia. *Exp Aging Res* 1994; 20: 73-94.
 41. Marin RS, Wilkosz PA. Disorders of diminished motivation. *J Head Trauma Rehabil* 2005; 20: 377-88.
 42. Mulin E, Leone E, Dujardin K et al. Diagnostic criteria for apathy in clinical practice. *Int J Geriatr Psychiatry* 2011; 26: 158-65.
 43. Teng E, Ringman JM, Ross LK et al. Diagnosing depression in Alzheimer disease with the National Institute of Mental Health provisional criteria. *Am J Geriatr Psychiatry* 2008; 16: 469-77.
 44. Lee HB, Lyketsos CG. Depression in Alzheimer's disease: heterogeneity and related issues. *Biol Psychiatry* 2003; 54: 353-62.
 45. Levy ML, Cummings JL, Fairbanks LA et al. Apathy is not depression. *J Neuropsychiatry Clin Neurosci* 1998; 10: 314-9.
 46. Olin JT, Katz IR, Meyers BS, Schneider LS, Lebowitz BD. Provisional diagnostic criteria for depression of Alzheimer disease: rationale and background. *Am J Geriatr Psychiatry* 2002; 10: 129-41.
 47. Ballard C, Bannister C, Solis M, Oyeboode F, Wilcock G. The prevalence, associations and symptoms of depression amongst dementia sufferers. *J Affect Disord* 1996; 36: 135-44.
 48. Lyketsos CG, Baker L, Warren A et al. Depression, delusions, and hallucinations in Alzheimer's disease: no relationship to apolipoprotein E genotype. *J Neuropsychiatry Clin Neurosci* 1997; 9: 64-7.
 49. Forsell Y, Winblad B. Major depression in a population of demented and nondemented older people: prevalence and correlates. *J Am Geriatr Soc* 1998; 46: 27-30.
 50. Cummings JL. Cognitive and behavioral heterogeneity in Alzheimer's disease: seeking the neurobiological basis. *Neurobiol Aging* 2000; 21: 845-61.
 51. Fitz AG, Teri L. Depression, cognition, and functional ability in patients with Alzheimer's disease. *J Am Geriatr Soc* 1994; 42: 186-91.
 52. Lyketsos CG, Steele C, Baker L et al. Major and minor depression in Alzheimer's disease: prevalence and impact. *J Neuropsychiatry Clin Neurosci* 1997; 9: 556-61.
 53. Royall DR, Mahurin RK, Cornell J. Effect of depression on dementia presentation: qualitative assessment with the Qualitative Evaluation of Dementia (QED). *J Geriatr Psychiatry Neurol* 1995; 8: 4-11.
 54. Gonzalez-Salvador MT, Arango C, Lyketsos CG, Barba AC. The stress and psychological morbidity of the Alzheimer patient caregiver. *Int J Geriatr Psychiatry* 1999; 14: 701-10.
 55. Kopetz S, Steele CD, Brandt J et al. Characteristics and outcomes of dementia residents in an assisted living facility. *Int J Geriatr Psychiatry* 2000; 15: 586-93.
 56. Lyketsos CG, Olin J. Depression in Alzheimer's disease: overview and treatment. *Biol Psychiatry* 2002; 52: 243-52.
 57. Gonzalez-Salvador T, Lyketsos CG, Baker A et al. Quality of life in dementia patients in long-term care. *Int J Geriatr Psychiatry* 2000; 15: 181-9.
 58. Lyketsos CG, Steele C, Galik E et al. Physical aggression in dementia patients and its relationship to depression. *Am J Psychiatry* 1999; 156: 66-71.
 59. Steele C, Rovner B, Chase GA, Folstein M. Psychiatric symptoms and nursing home placement of patients with Alzheimer's disease. *Am J Psychiatry* 1990; 147: 1049-51.
 60. Starkstein SE, Mizrahi R, Power BD. Depression in Alzheimer's disease: phenomenology, clinical correlates and treatment. *Int Rev Psychiatry* 2008; 20: 382-8.
 61. Migliorelli R, Teson A, Sabe L, Petracchi M, Leiguarda R, Starkstein SE. Prevalence and correlates of dysthymia and major

- depression among patients with Alzheimer's disease. *Am J Psychiatry* 1995; 152: 37-44.
62. Lawlor BA, Ryan TM, Schmeidler J, Mohs RC, Davis KL. Clinical symptoms associated with age at onset in Alzheimer's disease. *Am J Psychiatry* 1994; 151: 1646-9.
 63. Pearlson GD, Ross CA, Lohr WD, Rovner BW, Chase GA, Folstein MF. Association between family history of affective disorder and the depressive syndrome of Alzheimer's disease. *Am J Psychiatry* 1990; 147: 452-6.
 64. Strauss ME, Ogrocki PK. Confirmation of an association between family history of affective disorder and the depressive syndrome in Alzheimer's disease. *Am J Psychiatry* 1996; 153: 1340-2.
 65. Harwood DG, Barker WW, Ownby RL, Duara R. Association between premorbid history of depression and current depression in Alzheimer's disease. *J Geriatr Psychiatry Neurol* 1999; 12: 72-5.
 66. McGuire MH, Rabins PV. Mood disorders. In Coffey CE, Cummings JL, Eds. *Textbook of Geriatric Neuropsychiatry*. Washington, DC: American Psychiatric Press, 1994.
 67. Vilalta-Franch J, Garre-Olmo J, Lopez-Pousa S et al. Comparison of different clinical diagnostic criteria for depression in Alzheimer disease. *Am J Geriatr Psychiatry* 2006; 14: 589-97.
 68. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. Washington, DC: APA, 1994.
 69. Mega MS, Cummings JL, Fiorello T, Gornbein J. The spectrum of behavioral changes in Alzheimer's disease. *Neurology* 1996; 46: 130-5.
 70. Lerner AJ, Strauss M, Sami SA. Recognizing apathy in Alzheimer's disease. *Geriatrics* 2007; 62: 14-7.
 71. Olin JT, Schneider LS, Katz IR et al. Provisional diagnostic criteria for depression of Alzheimer disease. *Am J Geriatr Psychiatry* 2002; 10: 125-8.
 72. Rosenberg PB, Onyike CU, Katz IR et al. Depression of Alzheimer's disease study. Clinical application of operationalized criteria for 'Depression of Alzheimer's Disease'. *Int J Geriatr Psychiatry* 2005; 20: 119-27.
 73. Alexopoulos GS, Meyers BS, Young RC et al. Recovery in geriatric depression. *Arch Gen Psychiatry* 1996; 53: 305-12.
 74. Gallo JJ, Rabins PV. Depression without sadness: alternative presentations of depression in late life. *Am Fam Physician* 1999; 60: 820-6.
 75. Teri L, Wagner AW. Assessment of depression in patients with Alzheimer's disease: concordance among informants. *Psychol Aging* 1991; 6: 280-5.
 76. Janzing JG, Hooijer C, van 't Hof MA, Zitman FG. Depression in subjects with and without dementia: a comparison using GMS-AGECAT. *Int J Geriatr Psychiatry* 2002; 17: 1-5.
 77. Zubenko GS, Zubenko WN, McPherson S, et al. A collaborative study of the emergence and clinical features of the major depressive syndrome of Alzheimer's disease. *Am J Psychiatry* 2003; 160: 857-66.
 78. American Psychiatric Association. *Quick Reference to the Diagnostic Criteria from DSM-IV-TR*. Arlington: American Psychiatric Association; 2000.
 79. Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry* 1960; 23: 56-62.
 80. Ready RE, Ott BR, Grace J, Cahn-Weiner DA. Apathy and executive dysfunction in mild cognitive impairment and Alzheimer disease. *Am J Geriatr Psychiatry* 2003; 11: 222-8.
 81. Starkstein SE, Fedoroff JP, Price TR, Leiguarda R, Robinson RG. Apathy following cerebrovascular lesions. *Stroke* 1993; 24: 1625-30.
 82. Starkstein SE, Mizrahi R, Garau L. Specificity of symptoms of depression in Alzheimer disease: a longitudinal analysis. *Am J Geriatr Psychiatry* 2005; 13: 802-7.
 83. Forsell Y, Jorm AF, Fratiglioni L, Grut M, Winblad B. Application of DSM-III-R criteria for major depressive episode to elderly subjects with and without dementia. *Am J Psychiatry* 1993; 150: 1199-202.
 84. Forsell Y, Jorm AF, Winblad B. Association of age, sex, cognitive dysfunction, and disability with major depressive symptoms in an elderly sample. *Am J Psychiatry* 1994; 151: 1600-4.
 85. Marin RS. Differential diagnosis of apathy and related disorders of diminished motivation. *Psychiatric Ann* 1997; 27: 30-3.
 86. Marin RS. Apathy and related disorders of diminished motivation. In Dickstein LJ, Riba MB, Oldham JM, Eds. *Review of Psychiatry*. Washington, DC: American Psychiatric Press, 1996: 205-42.
 87. Royall DR. Some "depressive" symptoms may not imply depression. *J Am Geriatr Soc* 1997; 45: 891-2.
 88. Paulsen JS, Stout JC, De La Pena J et al. Frontal behavioral syndromes in cortical and subcortical dementia. *Assessment* 1996; 3: 327-37.
 89. Starkstein SE, Mayberg HS, Preziosi TJ, Andrezejewski P, Leiguarda R, Robinson RG. Reliability, validity, and clinical correlates of apathy in Parkinson's disease. *J Neuropsychiatry Clin Neurosci* 1992; 4: 134-9.
 90. Oguru M, Tachibana H, Toda K, Okuda B, Oka N. Apathy and depression in Parkinson disease. *J Geriatr Psychiatry Neurol* 2010; 23: 35-41.
 91. Starkstein SE, Merello M, Jorge R, Brockman S, Bruce D, Power B. The syndromal validity and nosological position of apathy in Parkinson's disease. *Mov Disord* 2009; 15; 24: 1211-6.
 92. Pluck GC, Brown RG. Apathy in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2002; 73: 636-42.
 93. Sims A. *Symptoms in the Mind*. London: Saunders, 2003.
 94. Marin RS, Firinciogullari S, Biedrzycki RC. Group differences in the relationship between apathy and depression. *J Nerv Ment Dis* 1994; 182: 235-9.
 95. Bozzola FG, Gorelick PB, Freels S. Personality changes in Alzheimer's disease. *Arch Neurol* 1992; 49: 297-300.
 96. Hwang TJ, Masterman DL, Ortiz F, Fairbanks LA, Cummings JL. Mild cognitive impairment is associated with characteristic neuropsychiatric symptoms. *Alzheimer Dis Assoc Disord* 2004; 18: 17-21.
 97. Geda YE, Smith GE, Knopman DS et al. De novo genesis of neuropsychiatric symptoms in mild cognitive impairment (MCI). *Int Psychogeriatr* 2004; 16: 51-60.
 98. Burns A, Jacoby R, Levy R. Psychiatric phenomena in Alzheimer's disease. III: Disorders of mood. *Br J Psychiatry* 1990; 157: 81-6, 92-4.
 99. Merriam AE, Aronson MK, Gaston P, Wey SL, Katz I. The psychiatric symptoms of Alzheimer's disease. *J Am Geriatr Soc* 1988; 36: 7-12.
 100. Lyketsos CG, Steinberg M, Tschanz JT, Norton MC, Steffens DC, Breitner JC. Mental and behavioral disturbances in dementia: findings from the Cache County Study on Memory in Aging. *Am J Psychiatry* 2000; 157: 708-14.
 101. Forsell Y, Jorm AF, Winblad B. Variation in psychiatric and behavioural symptoms at different stages of dementia: data from physicians' examinations and informants' reports. *Dementia* 1993; 4: 282-6.
 102. Mulsant BH, Ganguli M. Epidemiology and diagnosis of depression in late life. *J Clin Psychiatry* 1999; 60 (Suppl 20): 9-15.

103. Mattson AJ, Levin HS. Frontal lobe dysfunction following closed head injury. A review of the literature. *J Nerv Ment Dis* 1990; 178: 282-91.
104. Mega MS, Cummings JL. Frontal-subcortical circuits and neuropsychiatric disorders. *J Neuropsychiatry Clin Neurosci* 1994; 6: 358-70.
105. Tekin S, Mega MS, Masterman DM et al. Orbitofrontal and anterior cingulate cortex neurofibrillary tangle burden is associated with agitation in Alzheimer disease. *Ann Neurol* 2001; 49: 355-61.
106. Forstl H, Burns A, Levy R, Cairns N, Luthert P, Lantos P. Neuropathological correlates of behavioural disturbance in confirmed Alzheimer's disease. *Br J Psychiatry* 1993; 163: 364-8.
107. Bums A, Forstl H. The Institute of Psychiatry Alzheimer's Disease Cohort: Part 2. Clinicopathological observations. *Int J Geriatr Psychiatry* 1996; 11: 321-7.
108. Starkstein SE, Sabe L, Vazquez S et al. Neuropsychological, psychiatric, and cerebral perfusion correlates of leukoaraiosis in Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 1997; 63: 66-73.
109. Starkstein SE, Mizrahi R, Capizzano AA, Acion L, Brockman S, Power BD. Neuroimaging correlates of apathy and depression in Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 2009; 21: 259-65.
110. Jonsson M, Edman A, Lind K, Rolstad S, Sjogren M, Wallin A. Apathy is a prominent neuropsychiatric feature of radiological white-matter changes in patients with dementia. *Int J Geriatr Psychiatry* 2010; 25: 588-95.
111. Craig AH, Cummings JL, Fairbanks L et al. Cerebral blood flow correlates of apathy in Alzheimer disease. *Arch Neurol* 1996; 53: 1116-20.
112. Chase TN. Apathy in neuropsychiatric disease: diagnosis, pathophysiology, and treatment. *Neurotox Res* 2011; 19: 266-78.
113. Cummings JL. Frontal-subcortical circuits and human behavior. *Arch Neurol* 1993; 50: 873-80.
114. McPherson SE, Cummings JL. The neuropsychology of the frontal lobes. In Ron MA, David RS, Eds. *Disorders of brain and mind*. London: Cambridge University Press, 1998.
115. Mayberg HS. Clinical correlates of PET- and SPECT-identified defects in dementia. *J Clin Psychiatry* 1994; 55 (Suppl): 12-21.
116. Ebmeier KP, Prentice N, Ryman A et al. Temporal lobe abnormalities in dementia and depression: a study using high resolution single photon emission tomography and magnetic resonance imaging. *J Neurol Neurosurg Psychiatry* 1997; 63: 597-604.
117. Zubenko GS, Moosy J, Kopp U. Neurochemical correlates of major depression in primary dementia. *Arch Neurol* 1990; 47: 209-14.
118. Zubenko GS, Moosy J. Major depression in primary dementia. Clinical and neuropathologic correlates. *Arch Neurol* 1988; 45: 1182-6.
119. Forstl H, Burns A, Luthert P, Cairns N, Lantos P, Levy R. Clinical and neuropathological correlates of depression in Alzheimer's disease. *Psychol Med* 1992; 22: 877-84.
120. Chan-Palay V. Alterations in catecholamine neurons of the locus coeruleus in senile dementia of the Alzheimer type and in Parkinson's disease with and without dementia and depression. *J Comp Neurol* 1989; 287: 273-392.
121. Chen CP, Alder JT, Bowen DM et al. Presynaptic serotonergic markers in community-acquired cases of Alzheimer's disease: correlations with depression and neuroleptic medication. *J Neurochem* 1996; 66: 1592-8.
122. Zubenko GS, Moosy J, Martinez AJ et al. Neuropathologic and neurochemical correlates of psychosis in primary dementia. *Arch Neurol* 1991; 48: 619-24.
123. Zubenko GS. Biological correlates of clinical heterogeneity in primary dementia. *Neuropsychopharmacology* 1992; 6: 77-93.
124. Zweig RM, Ross CA, Hedreen JC et al. The neuropathology of aminergic nuclei in Alzheimer's disease. *Ann Neurol* 1988; 24: 233-42.
125. Zweig RM, Ross CA, Hedreen JC et al. Neuropathology of aminergic nuclei in Alzheimer's disease. *Prog Clin Biol Res* 1989; 317: 353-65.
126. Zubenko GS. Neurobiology of major depression in Alzheimer's disease. *Int Psychogeriatr* 2000; 12 (Suppl 1): 217-30.
127. Lopez OL, Becker JT, Reynolds CF 3rd, Jungreis CA, Weinman S, DeKosky ST. Psychiatric correlates of MR deep white matter lesions in probable Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 1997; 9: 246-50.
128. Lopez OL, Kambouh MI, Becker JT, Kaufer DI, DeKosky ST. The apolipoprotein E epsilon 4 allele is not associated with psychiatric symptoms or extrapyramidal signs in probable Alzheimer's disease. *Neurology* 1997; 49: 794-7.
129. Ott BR, Noto RB, Fogel BS. Apathy and loss of insight in Alzheimer's disease: a SPECT imaging study. *J Neuropsychiatry Clin Neurosci* 1996; 8: 41-6.
130. Benoit M, Dygai I, Migneco O et al. Behavioral and psychological symptoms in Alzheimer's disease. Relation between apathy and regional cerebral perfusion. *Dement Geriatr Cogn Disord* 1999; 10: 511-7.
131. Hirono N, Mori E, Ishii K et al. Frontal lobe hypometabolism and depression in Alzheimer's disease. *Neurology* 1998; 50: 380-3.
132. Lopez OL, Zivkovic S, Smith G, Becker JT, Meltzer CC, DeKosky ST. Psychiatric symptoms associated with cortical-subcortical dysfunction in Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 2001; 13: 56-60.
133. Royall DR. Frontal systems impairment in major depression. *Semin Clin Neuropsychiatry* 1999; 4: 13-23.
134. Joseph R. Frontal lobe psychopathology: mania, depression, confabulation, catatonia, perseveration, obsessive compulsions, and schizophrenia. *Psychiatry* 1999; 62: 138-72.
135. Holthoff VA, Beuthien-Baumann B, Kalbe E et al. Regional cerebral metabolism in early Alzheimer's disease with clinically significant apathy or depression. *Biol Psychiatry* 2005; 57: 412-21.
136. Cummings JL, Back C. The cholinergic hypothesis of neuropsychiatric symptoms in Alzheimer's disease. *Am J Geriatr Psychiatry* 1998; 6: S64-78.
137. Meltzer CC, Price JC, Mathis CA et al. PET imaging of serotonin type 2A receptors in late-life neuropsychiatric disorders. *Am J Psychiatry* 1999; 156: 1871-8.
138. Zubenko GS, Hughes HB 3rd, Stiffler JS. Clinical and neurobiological correlates of D10S1423 genotype in Alzheimer's disease. *Biol Psychiatry* 1999; 46: 740-9.
139. Gilley DW, Wilson RS, Bennett DA, Bernard BA, Fox JH. Predictors of behavioral disturbance in Alzheimer's disease. *J Gerontol* 1991; 46: P362-71.
140. Hoehn-Saric R, Lipsey JR, McLeod DR. Apathy and indifference in patients on fluvoxamine and fluoxetine. *J Clin Psychopharmacol* 1990; 10: 343-5.
141. Marin RS, Fogel BS, Hawkins J, Duffy J, Krupp B. Apathy: a treatable syndrome. *J Neuropsychiatry Clin Neurosci* 1995; 7: 23-30.
142. Settle EC, Jr. Antidepressant drugs: disturbing and potentially dangerous adverse effects. *J Clin Psychiatry* 1998; 59 (Suppl 16): 25-30; discussion 40-2.

143. Wongpakaran N, van Reekum R, Wongpakaran T, Clarke D. Selective serotonin reuptake inhibitor use associates with apathy among depressed elderly: a case-control study. *Ann Gen Psychiatry* 2007; 6: 7.
144. Kaufer DI, Cummings JL, Christine D. Effect of tacrine on behavioral symptoms in Alzheimer's disease: an open-label study. *J Geriatr Psychiatry Neurol* 1996; 9: 1-6.
145. Kaufer D, Cummings JL, Christine D. Differential neuropsychiatric symptom responses to tacrine in Alzheimer's disease: relationship to dementia severity. *J Neuropsychiatry Clin Neurosci* 1998; 10: 55-63.
146. Cummings JL, McRae T, Zhang R. Effects of donepezil on neuropsychiatric symptoms in patients with dementia and severe behavioral disorders. *Am J Geriatr Psychiatry* 2006; 14: 605-12.
147. Mega MS, Masterman DM, O'Connor SM, Barclay TR, Cummings JL. The spectrum of behavioral responses to cholinesterase inhibitor therapy in Alzheimer disease. *Arch Neurol* 1999; 56: 1388-93.
148. Benoit M, Koulibaly PM, Migneco O, Darcourt J, Pringuey DJ, Robert PH. Brain perfusion in Alzheimer's disease with and without apathy: a SPECT study with statistical parametric mapping analysis. *Psychiatry Res* 2002; 114: 103-11.
149. Migneco O, Benoit M, Koulibaly PM et al. Perfusion brain SPECT and statistical parametric mapping analysis indicate that apathy is a cingulate syndrome: a study in Alzheimer's disease and nondemented patients. *Neuroimage* 2001; 13: 896-902.
150. Sultzer DL, Mahler ME, Mandelkern MA et al. The relationship between psychiatric symptoms and regional cortical metabolism in Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 1995; 7: 476-84.
151. Forstmeier S, Maercker A. Potentially modifiable risk factors in the development of Alzheimer's disease. *Eur Neurol Rev* 2009; 4: 18-21.
152. Fiske A, Wetherell JL, Gatz M. Depression in older adults. *Annu Rev Clin Psychol* 2009; 5: 363-89.
153. Mortby ME, Maercker A, Forstmeier S. Midlife motivational abilities predict apathy and depression in Alzheimer's disease: the Aging, Demographics and Memory Study. *J Geriatr Psychiatry Neurol* 2011; 24: 151-60.
154. Low LF, Brodaty H, Draper B. A study of premorbid personality and behavioural and psychological symptoms of dementia in nursing home residents. *Int J Geriatr Psychiatry* 2002; 17: 779-83.
155. Kolanowski AM, Whall AL. Life-span perspective of personality in dementia. *Image J Nurs Sch* 1996; 28: 315-20.
156. Strauss ME, Lee MM, DiFilippo JM. Premorbid personality and behavioral symptoms in Alzheimer disease. Some cautions. *Arch Neurol* 1997; 54: 257-9.