



Short Communication

Wide spread of carbapenemase-producing bacterial isolates in a Nigerian environment

Christophe Le Terrier^a, Amandine Masseron^a, Nkolika Stella Uwaezuoke^b,
Chinagozi P. Edwin^c, Agantem E. Ekuma^d, Folake Olugbeminiyi^e, Shuwaram Shettima^f,
Simon Ushie^g, Laurent Poirel^{a,h,i,*}, Patrice Nordmann^{a,h,i,j}

^a Medical and Molecular Microbiology, Section of Medicine, Faculty of Science and Medicine, University of Fribourg, Switzerland

^b Department of Medical Microbiology and Parasitology, National Hospital Abuja, Abuja, Nigeria

^c Dept of Medical Microbiology & Parasitology, Aminu Kano Teaching Hospital, Kano, Nigeria

^d Department of Medical Microbiology, University of Calabar Teaching Hospital, Calabar, Nigeria

^e Department of Medical Microbiology, National Orthopaedic Hospital, Igbobi, Lagos, Nigeria

^f Department of Medical Microbiology, Federal Medical Center, Yola, Nigeria

^g Department of Medical Microbiology & Parasitology, Nnamdi Azikiwe University, Nnewi Campus, Nnewi, Nigeria

^h INSERM European Unit (IAME, France), University of Fribourg, Switzerland

ⁱ Swiss National Reference Center for Emerging Antibiotic Resistance (NARA), University of Fribourg, Switzerland

^j Institute for Microbiology, University of Lausanne and University Hospital Centre, Lausanne, Switzerland

ARTICLE INFO

Article history:

Received 11 July 2019

Received in revised form 11 October 2019

Accepted 11 October 2019

Available online 19 October 2019

Keywords:

Carbapenemase

Environment

Gram-negative bacteria

Nigeria

ABSTRACT

Objectives: The presence of carbapenemase-producing bacterial isolates is found not only in hospital and community settings but also in the environment. Carbapenemase production may be related to acquired, usually plasmid-borne, β -lactamase genes or to chromosomal genes intrinsic to various species. The aim of this study was to evaluate the occurrence of such carbapenemase-producing bacterial isolates among environmental samples from Nigeria.

Methods: A total of 122 environmental samples were plated on carbapenem-containing media. A total of 259 isolates were recovered, among which 124 were carbapenemase-producers according to the results of the Rapidec[®] Carba NP test.

Results: The majority of isolates ($n = 112$) recovered corresponded to natural producers of carbapenemases, i.e. *Stenotrophomonas maltophilia* ($n = 108$), *Burkholderia cepacia* ($n = 1$), *Shewanella* sp. ($n = 1$), *Sphingobacterium* sp. ($n = 1$) and *Chryseobacterium gleum* ($n = 1$). Ten isolates (mainly Enterobacteriaceae and *Acinetobacter baumannii*) produced an acquired carbapenemase, most commonly of the NDM type. In addition, two *Pseudomonas otitidis* isolates were identified as producing the Ambler class B carbapenemase POM-1, further confirming that this carbapenemase is naturally produced in this environmental species. Finally, several isolates co-producing 16S rRNA methylases (ArmA, RmtC) and/or extended-spectrum β -lactamases (CTX-M-9, CTX-M-15) were also identified.

Conclusion: This study revealed the presence and diversity of clinically-relevant antimicrobial-resistant bacteria in the environment in Nigeria.

© 2019 International Society for Antimicrobial Chemotherapy. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Carbapenems are last-resort antibiotics for managing multi-drug-resistant bacterial infections but their effectiveness may be compromised by the production of carbapenemases [1]. Knowing

the extent of the reservoir of carbapenemases is becoming important. The most clinically-relevant carbapenemases are of the KPC type (serine carbapenemase of Ambler class A), VIM-, IMP- and NDM-types (metallo- β -lactamases of Ambler class B) and OXA-48-type (oxacillinases of Ambler class D). Although extensive surveillance studies have been performed in Europe, America and Australia, data from Africa remain scarce. In Nigeria, the most populated country of Africa (ca. 200 million inhabitants), antibiotic use is widespread and largely unregulated [2]. Owing to a lack of sanitation and poor control of public health in many African countries, large amounts of antibiotics may end up in the

* Corresponding author. Present address: Medical and Molecular Microbiology Unit, Department of Medicine, Faculty of Science, University of Fribourg, Chemin du Musée 18, CH-1700 Fribourg, Switzerland.

E-mail address: laurent.poirel@unifr.ch (L. Poirel).

environment and thus contribute to the selection and persistence of antimicrobial resistance. Therefore, the primary aim of this study was to evaluate the spread of multidrug-resistant bacterial strains among environmental samples collected in Nigeria, with a particular focus on carbapenemase-producers.

2. Materials and methods

2.1. Soil and water samples

A total of 122 environmental samples (water and soil) were collected during the same time period in 2017 from six different locations in Nigeria, namely Abuja (North Central), Kano (North West), Yola (North East), Nnewi (South East), Akwa Ibom (South) and Ibadan (South West). Water samples were recovered from different bodies of lakes.

2.2. Selective isolation of carbapenem-resistant Gram-negative bacteria

Selection of carbapenem-resistant isolates was performed following a selective enrichment step by culture in brain–heart infusion broth supplemented with ertapenem (0.25 µg/mL) overnight at 30 °C. Following enrichment, 100 µL of pure and diluted (1/100; 1:1000) cultures was plated onto selective media, including SuperCarba[®] (CHROMagar, Paris, France) [3] and Drigalski agar medium supplemented with imipenem (0.75 µg/mL), daptomycin (10 µg/mL) and amphotericin B (5 µg/mL). Plates were then incubated at 30 °C for 24 h.

2.3. Phenotypic and genotypic characterisation of carbapenem-resistant bacteria

Antimicrobial susceptibility testing was initially performed by the disk diffusion method and was interpreted according to European Committee on Antimicrobial Susceptibility Testing (EUCAST) recommendations (<http://www.eucast.org>), except for colistin for which susceptibility was evaluated by broth micro-dilution, as recommended.

Carbapenemase activity was assessed by the Rapidec[®] Carba NP test (bioMérieux) for each colony type growing on the selective media [4]. Identification of carbapenemase genes was performed by PCR using previously published primers [5]. The following carbapenemase genes were searched for: Ambler class A genes of

*bla*_{KPC}, *bla*_{GES}, *bla*_{IMI} and *bla*_{FRI} types; Ambler class B genes of *bla*_{NDM}, *bla*_{IMB}, *bla*_{VIM}, *bla*_{GIM} and *bla*_{POM} types; and Ambler class D genes of *bla*_{OXA-48}, -181, -23, -40 and -58 types. In addition, the following genes were searched for: extended-spectrum β-lactamases (ESBLs), including *bla*_{CTX-M-1}, -2, -3, -8, -9 and -15, *bla*_{PER}, *bla*_{TEM} and *bla*_{SHV} [6]; aminoglycoside resistance 16S rRNA methylase genes (*armA*, *rmtA*–*rmtH* and *npmA*) [7]; and plasmid-borne colistin resistance *mcr*-like genes (*mcr-1* to *mcr-9*) [8]. All positive PCR amplicons were sent for sequencing (Microsynth AG, Balgach, Switzerland). Bacterial identification was performed by matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry (MALDI-TOF/MS) (VITEK[®] MS; bioMérieux).

Whole-genome sequencing was performed for the two *Pseudomonas* *otitidis* strains for which identification was uncertain. DNA was extracted from isolates using a QIAamp[®] DNA Mini Kit and QIAcube Workstation (QIAGEN, Courtaboeuf, France), according to the manufacturer's instructions. Genomic libraries were generated using a Nextera XT DNA Sample Preparation Kit (Illumina Switzerland GmbH, Zurich, Switzerland) and sequencing was performed on an Illumina MiSeq benchtop sequencer. Reads from sequencing were assembled using CLC Genomics Workbench 7 (QIAGEN). Putative β-lactamase genes were then detected by ResFinder (<https://cge.cbs.dtu.dk/services/ResFinder/>).

3. Results and discussion

From the 122 environmental samples, 259 distinct isolates were isolated on selective carbapenem-containing media. All carbapenem-resistant isolates were identified on both screening media. Of the 259 isolates, 124 were found to be carbapenemase-producers according to the results of the Rapidec[®] Carba NP test. The carbapenemase-negative isolates corresponded to 135 *Pseudomonas* spp. isolates (no acquired resistance genes) that were recovered only on the SuperCarba[®] medium owing to natural ertapenem resistance in this genus. The majority (*n* = 112) of the carbapenemase-producers identified corresponded to bacterial species that naturally produce carbapenemases such as *Stenotrophomonas maltophilia* (L1-like β-lactamase [9]) (*n* = 108), *Chryseobacterium gleum* (CGB-like carbapenemase [10]) (*n* = 1), *Burkholderia cepacia* (PenA chromosomal A penicillinase [11]) (*n* = 1), *Sphingobacterium* sp. (*n* = 1) and *Shewanella* sp. (OXA-181-like carbapenemase [12]) (*n* = 1).

Ten isolates harbouring an acquired carbapenemase gene were also identified (Table 1). A series of enterobacterial strains

Table 1
Information and genetic features of isolates with an acquired carbapenemase gene from environment samples from various geographical locations in Nigeria.

Strain no.	Type of sample ^a	Geographical location ^b	Species	Carbapenemase	ESBL	16S rRNA methyltransferase (s)	Non-susceptible phenotype ^c
1	Soil UO	Abuja (NC)	<i>Acinetobacter baumannii</i>	OXA-23	None	None	CIP, TET, SXT, CHL
2	Soil HE	Nnewi (SE)	<i>A. baumannii</i>	OXA-40	None	ArmA	GEN, AMK, CIP, SXT, CHL
3	Water	Nnewi (SE)	<i>A. baumannii</i>	OXA-40	None	ArmA	GEN, AMK, CIP, SXT, CHL
4	Soil HE	Abuja (NC)	<i>Aeromonas caviae</i>	NDM-1	CTX-M-9	ArmA, RmtC	GEN, AMK, SXT
5	Soil HE	Abuja (NC)	<i>Citrobacter freundii</i>	NDM-5	None	None	GEN, CIP, TET, SXT, CHL
6	Soil HE	Abuja (NC)	<i>Enterobacter cloacae</i>	NDM-5	CTX-M-15	None	GEN, CIP, TET, SXT
7	Soil HE	Akwa Ibom (S)	<i>E. cloacae</i>	NDM-7	None	None	SXT
8	Soil HE	Akwa Ibom (S)	<i>Klebsiella pneumoniae</i>	NDM-1	CTX-M-15	ArmA	GEN, AMK, CIP, TET, SXT
9	Soil HE	Akwa Ibom (S)	<i>K. pneumoniae</i>	NDM-1	CTX-M-15	ArmA	GEN, AMK, CIP, TET, SXT
10	Soil HE	Kano (NW)	<i>Pseudomonas putida</i>	VIM-5	None	None	CIP, TET, SXT, CHL

ESBL, extended-spectrum β-lactamase.

^a UO, urban outskirt; HE, hospital environment.

^b NC, North Central; SE, South East; S, South; SW, South West; NW, North West.

^c CIP, ciprofloxacin; TET, tetracycline; SXT, trimethoprim/sulfamethoxazole; CHL, chloramphenicol; GEN, gentamicin; AMK, amikacin.

producing different NDM-type carbapenemases was found in different locations in Nigeria. This finding highlights the environmental spread of the NDM-type carbapenemase in Nigeria [13]. The large spread of NDM-producers has already been identified in many other African countries [14], including Angola [15] and Togo [16], indicating that this continent may act as a secondary reservoir after the Indian subcontinent for spreading NDM-producing isolates to Europe. The *Acinetobacter baumannii* isolates produced either OXA-23 or OXA-40, which are extensively reported in that species [17]. In addition, a *bla*_{VIM-5}-positive *Pseudomonas putida* isolate was identified, as previously identified from polluted wetlands in Nigeria [18]. Genome sequencing allowed the identification of a *Sphingobacterium* sp. isolate that produced a naturally occurring Ambler class B carbapenemase [19] and two *P. otitidis* isolates that harboured the *bla*_{POM-1} gene. This latter gene encodes a class B carbapenemase, with no associated mobile genetic element, which has been previously identified as intrinsic to that bacterial species [20].

Regarding the non-β-lactamase resistance determinants, it is worth mentioning that the 16S rRNA methylase determinants conferring pan-aminoglycoside resistance were frequently associated with NDM-producers, as previously shown [21]. Noteworthy, the spread of 16S rRNA methylases in Gram-negative bacteria in Africa appears to represent a major problem, with recent reports showing their occurrence in, e.g., Enterobacterales in Angola, Libya, Egypt and Algeria as well as *A. baumannii* in Algeria and Egypt. Finally, three colistin-resistant isolates were recovered among the acquired carbapenem-resistant isolates (*A. baumannii* and *Enterobacter cloacae*) and natural carbapenem-resistant isolates (*Sphingobacterium* spp.) but were not positive for MCR-encoding genes, suggesting that the resistance was mediated by other mechanisms of resistance such as chromosomal mutations. These results may indicate that acquired colistin resistance has not yet spread among environmental isolates in Nigeria.

Overall, this study revealed the presence and diversity of clinically-relevant antimicrobial-resistant bacteria in the environment and demonstrates the need for further investigation.

Competing interests

None declared.

Funding

This work was funded by the University of Fribourg (Fribourg, Switzerland), the Swiss National Science Foundation [projects FNS-31003A_163432 and FNS-407240_177381] and INSERM (Paris, France).

References

- [1] Nordmann P, Poirel L. Epidemiology and diagnostics of carbapenem resistance in Gram-negative bacteria. Clin Infect Dis 2019;69(Suppl 7):S521–8.
- [2] Ogbolu DO, Webber MA. High-level and novel mechanisms of carbapenem resistance in Gram-negative bacteria from tertiary hospitals in Nigeria. Int J Antimicrob Agents 2014;43:412–7.
- [3] Nordmann P, Girlich D, Poirel L. Detection of carbapenemase producers in Enterobacteriaceae by use of a novel screening medium. J Clin Microbiol 2012;50:2761–6.
- [4] Poirel L, Nordmann P, Rapidec Carba NP. test for rapid detection of carbapenemase producers. J Clin Microbiol 2015;23:3003–8.
- [5] Poirel L, Walsh TR, Cuvillier V, Nordmann P. Multiplex PCR for detection of acquired carbapenemase genes. Diagn Microbiol Infect Dis 2011;70:119–23.
- [6] Lartigue MF, Zinsius C, Wenger A, Bille J, Poirel L, Nordmann P. Extended-spectrum β-lactamases of the CTX-M type now in Switzerland. Antimicrob Agents Chemother 2007;51:2855–60.
- [7] Berçot B, Poirel L, Nordmann P. Updated multiplex polymerase chain reaction for detection of 16S rRNA methylases: high prevalence among NDM-1 producers. Diagn Microbiol Infect Dis 2011;71:442–5.
- [8] Lescat M, Poirel L, Nordmann P. Rapid multiplex PCR for detection of *mcr-1* to *mcr-5* genes. Diagn Microbiol Infect Dis 2018;92:267–9.
- [9] Crowder MW, Walsh TR, Banovic L, Pettit M, Spencer J. Overexpression, purification, and characterization of the cloned metallo-β-lactamase L1 from *Stenotrophomonas maltophilia*. Antimicrob Agents Chemother 1998;42:921–6.
- [10] Bellais S, Naas T, Nordmann P. Genetic and biochemical characterization of CGB-1, an Ambler class B carbapenem-hydrolyzing β-lactamase from *Chryseobacterium gleum*. Antimicrob Agents Chemother 2002;46:2791–6.
- [11] Trépanier S, Prince A, Huletsky A. Characterization of the *penA* and *penR* genes of *Burkholderia cepacia* 249 which encode the chromosomal class A penicillinase and its LysR-type transcriptional regulator. Antimicrob Agents Chemother 1997;41:2399–405.
- [12] Potron A, Poirel L, Nordmann P. Origin of OXA-181, an emerging carbapenem-hydrolyzing oxacillinase, as a chromosomal gene in *Shewanella xiamenensis*. Antimicrob Agents Chemother 2011;55:4405–7.
- [13] Oyelade AA, Adelowo OO, Fagade OE. *bla*_{NDM-1}-producing *Vibrio parahaemolyticus* and *V. vulnificus* isolated from recreational beaches in Lagos, Nigeria. Environ Sci Pollut Res Int 2018;25:33538–47.
- [14] Manenzhe RI, Zar HJ, Nicol MP, Kaba M. The spread of carbapenemase-producing bacteria in Africa: a systematic review. J Antimicrob Chemother 2014;70:23–40.
- [15] Kieffer N, Nordmann P, Aires-de-Sousa M, Poirel L. High prevalence of carbapenemase-producing Enterobacteriaceae among hospitalized children in Luanda, Angola. Antimicrob Agents Chemother 2016;60:6189–92.
- [16] Dossim S, Bonnin RA, Salou M, Tanga K, Godonou V, Dagnra AY, et al. Occurrence of carbapenemase-producing Enterobacteriaceae in Togo, West Africa. Int J Antimicrob Agents 2019;53:530–2.
- [17] Potron A, Poirel A, Nordmann P. Emerging broad-spectrum resistance in *Pseudomonas aeruginosa* and *Acinetobacter baumannii*: mechanisms and epidemiology. Int J Antimicrob Agents 2015;45:568–85.
- [18] Adelowo OO, Vollmers J, Mäusezahl I, Kaster AK, Müller JA. Detection of the carbapenemase gene *bla*_{VIM-5} in members of the *Pseudomonas putida* group isolated from polluted Nigerian wetlands. Sci Rep 2018;8:15116.
- [19] Barahona F, Slim J. *Sphingobacterium multivorum*: case report and literature review. New Microbe New Infect 2015;7:33–6.
- [20] Thaller MC, Borgianni L, Di Lallo G, Chong Y, Lee K, Dajcs J, et al. Metallo-β-lactamase production by *Pseudomonas otitidis*: a species-related trait. Antimicrob Agents Chemother 2011;55:118–23.
- [21] Poirel L, Goutines J, Aires-de-Sousa M, Nordmann P. High rate of association of 16S rRNA methylases and carbapenemases in Enterobacteriaceae recovered from hospitalized children in Angola. Antimicrob Agents Chemother 2018;62:e00021–18.